arXiv:hep-th/9901094v1 20 Jan 1999

Supersolutions

Pierre Deligne and Daniel S. Freed

Abstract. We develop classical globally supersymmetric theories. As much as possible, we treat various dimensions and various amounts of supersymmetry in a uniform manner. We discuss theories both in components and in superspace. Throughout we emphasize geometric aspects. The beginning chapters give a general discussion about supersymmetric field theories; then we move on to detailed computations of lagrangians, etc. in specific theories. An appendix details our sign conventions. This text will appear in a two-volume work *Quantum Fields and Strings: A Course for Mathematicians* to be published soon by the American Mathematical Society. Some of the cross-references may be found at `http://www.math.ias.edu/~drm/QFT/`.

Introduction

This text develops the basic classical supersymmetric field theories in all dimensions, but without supergravity. We emphasize at the start that although we hope to provide a bit of a fresh viewpoint in some places, this is standard material for the working quantum field theorist (string theorist, M-theorist, ...)—it was largely developed in the physics literature of the 1970s and 1980s. This account began as the solutions to the Superhomework problems in [I-**Homework**], but then developed into a larger project. Unfortunately, we often sink into an unpalatable morass of indices which may obscure the underlying geometry.[3] So we hope in this introduction to at least provide a roadmap to the more conceptual parts of the text. Certainly some readers may find the plethora of formulas useful. Indeed, we have gone to great lengths[4] to ensure their accuracy, though undoubtedly errors have crept in. Since there are so many formulas, we have boxed the ones we thought most crucial or most likely to be referenced. One contribution here is a consistent choice of conventions in all dimensions and with varying amounts of supersymmetry. Our sign conventions are summarized in the appendix. Throughout we freely use notions of supergeometry as developed in [I-**Supersymmetry**] and notions of classical field theory as developed in [I-**Classical Fields**]. Knowledge of spinors [I-**Spinors**] is also indispensable.

[1]School of Mathematics, Institute for Advanced Study, Olden Lane, Princeton NJ 08540
E-mail address: `deligne@math.ias.edu`
[2]Department of Mathematics, University of Texas, Austin TX 78712
E-mail address: `dafr@math.utexas.edu`
[3]As Mel Brooks famously said, "We mock the thing we are to be."
[4]in units where $c = 1$

The basic Poincaré-invariant classical field theories of a scalar field (spin 0), spinor field (spin 1/2), and gauge field (spin 1) exist in any dimension n. Here we are interested in field theories which are invariant under a supergroup which extends the Poincaré group. Such a *super Poincaré group* is determined by a real spin representation S^* of the Lorentzian spin group $\mathrm{Spin}(1, n-1)$ together with a symmetric pairing $S^* \otimes S^* \to V$ into the vectors, as we explain in §1.1. A theory has *minimal supersymmetry* if S^* is a minimal (that is, irreducible) spin representation; otherwise, it is said to have *extended supersymmetry*. We construct theories on Minkowski space which obey the spin-statistics connection, so that odd symmetries map fields of integral spin to fields of half-integral spin. It is convenient to measure the amount of supersymmetry in terms of $s = \dim S^*$. The larger s is, the greater the mixing of fields of different spin. As we consider theories only[5] with fields of spins at most one, we are restricted to small values of s. Specifically, supersymmetric σ-models—which only have fields of spins at most one-half—must have $s \leq 8$ and supersymmetric gauge theories must have $s \leq 16$. Because of this restriction the theory has a small set of general examples and the study of the theory is centered on the examples. We mostly consider theories with $s = 2, 4, 8, 16$ supersymmetries and these occur in the maximal dimensions $n = 3, 4, 6, 10$. "Maximal" means, for example, that only in dimensions ≤ 6 are there real spin representations of dimension 8. (We also say a bit in §1.3 about theories with a single supersymmetry and there is more in Problem FP2 of [I-**Homework**], which we recommend as a starting point before tackling the more complicated theories with more supersymmetry. Other problems in [I-**Homework**] also deal with formal properties of supersymmetric field theories.) By the exceptional isomorphisms among low dimensional Lie groups, the Lorentzian spin group in dimensions $n = 3, 4, 6$ is $SL(2, \mathbb{F})$ for $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$. (There is also a sense in which the spin group in 10 dimensions is "$SL(2, \mathbb{O})$" for the octonions $\mathbb{O}$: see [I-**Spinors**, §6.5–§6.7].) Our treatment emphasizes this exceptional isomorphism, both in its theoretical and computational aspects. Throughout we focus on theories in the maximal dimension, though in some cases (see Chapter 9, for example) we work out some lower dimensional theories obtained by dimensional reduction. (Not everything in lower dimensions comes by dimensional reduction from the maximal dimension, but we barely mention the exceptions.)

A supersymmetric field theory is first of all an ordinary (lagrangian) field theory on Minkowski space, described by a set of fields and a lagrangian. As usual the Poincaré symmetry is manifest. In addition, there are *nonmanifest* odd infinitesimal symmetries corresponding to the odd generators of the super Poincaré algebra. In this approach one checks by direct computation that the theory is supersymmetric, i.e., that the odd transformation of the fields is a symmetry of the lagrangian. This is the approach we follow in Chapter 3 for σ-models and in Chapter 6 for pure Yang-Mills theories. Although there are heuristic arguments to motivate the form of the odd symmetries, they are not geometric. Moreover, in many examples the bracketing relations of the super Poincaré algebra are only obeyed on-shell. It is more satisfying to use a *superspace formulation* in which the entire super Poincaré symmetry is manifest. In this approach the spacetime M of the classical field theory is *super Minkowski space*, a supermanifold whose underlying ordinary manifold is ordinary Minkowski space. The super Poincaré group acts

[5]For a physical explanation of this restriction, see [II-**Dynamics of QFT**, §2.4]

as symmetry transformations of super Minkowski space. The fields in the theory, usually called *superfields*, are "functions" on super Minkowski space. Then super Poincaré acts on superfields by pullback, and we can write manifestly supersymmetric lagrangians. By restricting superfields and their derivatives to the underlying ordinary Minkowski space, we recover the *component fields* of the ordinary field theory on Minkowski space, and by partial integration we recover the *component lagrangian*. These general ideas are explained in Chapter 1.

Although superspace formulations have the overriding virtue of exhibiting the supersymmetry in a manifest way, there are several drawbacks. First and foremost, there is not always a superspace formulation! Unfortunately, that is the case for theories with the maximal allowed supersymmetry: σ-models with 8 supersymmetries and Yang-Mills theories with 16 supersymmetries. There are good superspace formulations for all theories with at most 4 supersymmetries and also for pure gauge theories with at most 8 supersymmetries. Of course, one can use the superspace formulation with 4 supersymmetries even for theories with more supersymmetry; then some of the supersymmetry is manifest and some is nonmanifest. Another disadvantage of superspace formulations is that in many cases the superfields are constrained. This complicates the classical field theory computations. Also, the constraints vary from case to case and add further to the specificity of the subject. To a physicist the most damning criticism of the superspace formulation is that it obscures the physical intuition, which is based on experience with ordinary fields on Minkowski space. These drawbacks are balanced by the manifest geometric realization of odd symmetries, particularly when applied to the quantum theory. (For example, the possible form of quantum corrections to the classical action is clearer in the superspace formulation.) So we persist in presenting the superspace point of view alongside the component approach.

Chapters 1 and 2 are preliminary. Chapter 1 describes features of classical supersymmetric field theories which go beyond the general ideas of classical field theory. In Chapter 2 we describe the coordinates and bases we use to compute in 3, 4, and 6 dimensions. As mentioned before, they are based on the special form of the spin group in those cases. We list many formulas for easy future reference.

Our real work begins in Chapter 3, where we study supersymmetric σ-models in components. The quantum theories are often discussed in 2 dimensions, but we treat the classical theories in the maximal dimensions $n = 3, 4, 6$ (where they have minimal supersymmetry). In the linear case we give a uniform description of the theory and a uniform proof of supersymmetry. In the nonlinear case the scalar field is a map from ordinary Minkowski space to a curved Riemannian manifold X. Supersymmetry constrains X: in the $n = 4$ theory X must be Kähler and in the $n = 6$ theory X must by hyperkähler. In §3.3 we give a uniform motivation for these constraints. In the nonlinear case there are curvature terms in the lagrangian, and as they vary in the three cases we check supersymmetry on a case-by-case basis. In addition to the kinetic terms, there are supersymmetric potential terms one can add in the $n = 3, 4$ theories; they are summarized in §3.4.

Chapters 4 and 5 present the superspace formulation of supersymmetric σ-models in $n = 3, 4$ dimensions, including potential terms. In Chapter 4 we not only derive the component lagrangian and supersymmetry transformations, but we also discuss the classical field theory computations directly in superspace, including the computation of the Noether current for the supersymmetries. This is the only example we work out in such detail in superspace. The superspace lagrangian of

the $n = 4$ theory is usually given in terms of a local Kähler potential on the target manifold X, and so does not make sense globally. In §5.2 we point out a global lagrangian in case the target manifold is Hodge.

In Chapter 6 we add gauge fields to the mix. We begin in §6.1 with pure supersymmetric Yang-Mills theories in components. The usual argument which verifies supersymmetry relies on a "Fierz identity," something we do not encounter in our treatment. Instead we rely on a geometric property of spinors which only holds in dimensions $n = 3, 4, 6, 10$ with the minimal real spinor representation: the quadratic pairing on spinors takes values on the light cone. (In other cases the values are positive timelike or lightlike, not just lightlike.) That same property enters again in §6.3, where we describe some common features of the superspace formulation of the theory in $n = 3, 4, 6$ dimensions. In each case the basic superfield is a connection on super Minkowski space constrained to have vanishing curvature along a canonical odd distribution. The aforementioned special property of spinors is used to see that this constraint leads to the correct physical component fields. On the other hand, the auxiliary fields[6] differ in each case; they are treated in later chapters.

The most general theory without gravity is a gauge theory with matter—a coupled system consisting of scalar fields, spinor fields, and gauge fields. The general supersymmetric theories are summarized in Theorem 6.33. Of interest in applications are the nontrivial potential energy functions which appear; they lead to nontrivial classical moduli spaces. We describe the coupled systems in dimensions $n = 3, 4, 6$ with minimal supersymmetry. (Recall that there is no possible σ-model with $s = 16$ supersymmetries, only the pure Yang-Mills theory described previously.) Systems with $s = 2, 4, 8$ supersymmetries in lower dimensions are often, but not always, obtained by dimensional reduction from the theories described here. For $n = 3, 4$ there is a good superspace description from which the component description may be derived. This is the subject of Chapters 7 and 8. In the $n = 4$ case the appearance of the moment map in the component lagrangian has a geometric explanation, at least if the target Kähler manifold is Hodge (see §8.3). In Chapters 7 through 11 we only describe the Yang-Mills terms in the lagrangian, having treated the σ-model terms in Chapters 4 and 5. Chapter 10 gives the superspace description of super Yang-Mills theory in $n = 6$ dimensions. (Recall here that there is no superspace description of the σ-model, hence none of the coupled system.) For each of the cases $n = 3, 4, 6$ we prove that the category[7] of constrained connections is equivalent to the category of component fields. For the $n = 6$ case the proof is presented in Chapter 11.

A connection is locally specified by n Lie algebra-valued functions in n dimensions, and the dimensional reduction to $n - k$ dimensions still has n functions: $n - k$ of them transform as a connection and k of them as scalar fields (see §9.1). There are two cases of dimensionally reduced super Yang-Mills theory which are of particular interest—the reduction of the $n = 4$ theory to 2 dimensions and the reduction of the $n = 6$ theory to 4 dimensions. The former is the subject of Chapter 9 and the latter the subject of §10.2.

[6] An *auxiliary field* has an algebraic equation of motion and does not correspond to a physical degree of freedom.

[7] Because of the local gauge transformations, the collection of connections (gauge fields) forms a category, not a set.

Finally, in §5.5 and §10.3 we describe theories in $n = 4$ dimensions with extended supersymmetry on the minimal super Minkowski space. Then four of the supersymmetries are manifest and the extended supersymmetry is nonmanifest.

In several places we refer to the $n = 3, 4, 6$ cases with minimal supersymmetry as the $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ cases.

Writing this has been an absolute *cauchemar de signes*! Standard differential geometry has some bad signs, classical field theory has tricky sign conventions, and odd variables add a whole new level of complication. Our sign conventions are explained in the previous texts in Part 1 and are summarized in [I-**Signs**].

During the preparation of this manuscript we received advice and help from many people. Certainly Edward Witten's Superhomework [I-**Homework**] was our starting point for organizing the material, and Ed was invaluable at many other times. Input from our classmates, notably David Kazhdan and John Morgan, is also reflected here, as is the advice of Joseph Bernstein, Nati Seiberg, and many others. However, they should not be held accountable for any errors.

Contents

Chapter 1. Preliminary Topics

§1.1. Super Minkowski spaces and super Poincaré groups
§1.2. Superfields, component fields, and lagrangians
§1.3. A simple example

Chapter 2. Coordinates on Superspace

§2.1. $M^{3|2}$, $M^{4|4}$, $M^{6|(8,0)}$ and their complexifications
§2.2. Dimensional reduction
§2.3. Coordinates on $M^{3|2}$
§2.4. Coordinates on $M^{4|4}$
§2.5. Coordinates on $M^{6|(8,0)}$
§2.6. Low dimensions

Chapter 3. Supersymmetric σ-Models

§3.1. Preliminary remarks on linear algebra
§3.2. The free supersymmetric σ-model
§3.3. Nonlinear supersymmetric σ-model
§3.4. Supersymmetric potential terms
§3.5. Superspace construction
§3.6. Dimensional reduction

Chapter 4. The Supersymmetric σ-model in dimension 3

§4.1. Fields and supersymmetry transformations on $M^{3|2}$
§4.2. The σ-model action on $M^{3|2}$
§4.3. The potential term on $M^{3|2}$
§4.4. Analysis of the classical theory
§4.5. Reduction to $M^{2|(1,1)}$

Chapter 5. The Supersymmetric σ-Model in dimension 4

§5.1. Fields and supersymmetry transformations on $M^{4|4}$
§5.2. The σ-model action on $M^{4|4}$

§5.3. The superpotential term on $M^{4|4}$
§5.4. Analysis of the classical theory

Chapter 6. Supersymmetric Yang-Mills Theories
§6.1. The minimal theory in components
§6.2. Gauge theories with matter
§6.3. Superspace construction

Chapter 7. $N = 1$ Yang-Mills Theory in Dimension 3
§7.1. Constrained connections on $M^{3|2}$
§7.2. The Yang-Mills action on $M^{3|2}$
§7.3. Gauge theory with matter on $M^{3|2}$

Chapter 8. $N = 1$ Yang-Mills Theory in Dimension 4
§8.1. Constrained connections on $M^{4|4}$
§8.2. The Yang-Mills action on $M^{4|4}$
§8.3. Gauge theory with matter on $M^{4|4}$

Chapter 9. $N = 2$ Yang-Mills in Dimension 2
§9.1. Dimensional reduction of bosonic Yang-Mills
§9.2. Constrained connections on $M^{2|(2,2)}$
§9.3. The reduced Yang-Mills action

Chapter 10. $N = 1$ Yang-Mills in Dimension 6 and $N = 2$ Yang-Mills in Dimension 4
§10.1. Constrained connections on $M^{6|(8,0)}$
§10.2. Reduction to $M^{4|8}$
§10.3. More theories on $M^{4|4}$ with extended supersymmetry

Chapter 11. The Vector Multiplet on $M^{6|(8,0)}$
§11.1. Complements on $M^{6|(8,0}$
§11.2. Constrained connections
§11.3. An auxiliary Lie algebra
§11.4. Components of constrained connections

Appendix: Sign Manifesto

References

CHAPTER 1
Preliminary Topics

A supersymmetric field theory has as symmetry group a *super Poincaré group*. In §1.1 we define these supergroups as symmetry groups of *super Minkowski space*. In many cases supersymmetric theories have a superspace formulation in terms of fields defined on super Minkowski space. Passage from the superspace formulation to the formulation in "components" on ordinary Minkowski space is the subject of §1.2. The simplest example is in one dimension (time) with one odd variable; this is described in §1.3.

§1.1. Super Minkowski spaces and super Poincaré groups

A super Minkowski space M is constructed as follows. The starting data are a Minkowski space $\check{M}$ with vector space of translations V, a positive cone C of time-like vectors in V, a real spinorial representation S of $\mathrm{Spin}(V)$, and a symmetric morphism Γ of representations of $\mathrm{Spin}(V)$:

$$\Gamma\colon S^* \otimes S^* \longrightarrow V. \tag{1.1}$$

which is positive definite in the sense that $\Gamma(s^*, s^*) \in \overline{C}$ for $s^* \in S^*$, with $\Gamma(s^*, s^*) = 0$ only for $s^* = 0$.

Minkowski space is an affine space: it has no origin. By the classification in [I-**Spinors**, §6.3(i)], once the representation S is given, positive definite Γ as in (1.1) exist and are unique up to automorphisms of the representation S. If S is an irreducible real spinorial representation, nonzero symmetric morphisms $S^* \otimes S^* \to V$ are unique up to a factor, are positive or negative definite, and are respected by elements of norm 1 in the field of endomorphisms of S (isomorphic to $\mathbb{R}$, $\mathbb{C}$ or $\mathbb{H}$) [I-**Spinors**, §6.1].

Note that so far we have not used the metric on V, only its conformal structure and the choice of a positive cone.

By [I-**Spinors**, §6.2] there is a unique symmetric morphism

$$\tilde{\Gamma}\colon S \otimes S \to V \tag{1.2}$$

related to Γ by the following formula. With respect to a basis $\{e_\mu\}$ of V and $\{f^a\}$ of S, write[8] $\Gamma(f_a, f_b) = \Gamma^\mu_{ab} e_\mu$ and $\tilde{\Gamma}(f^a, f^b) = \tilde{\Gamma}^{\mu ab} e_\mu$. Then

$$\boxed{\Gamma^\mu_{ab}\,\tilde{\Gamma}^{\nu bc} + \Gamma^\nu_{ab}\,\tilde{\Gamma}^{\mu bc} = 2g^{\mu\nu}\delta^c_a\ .} \tag{1.3}$$

[8] We use the usual summation convention throughout: Repeated indices are summed if one index is "upstairs" and one is "downstairs." Also, the Kronecker δ-symbol has its usual meaning.

If one uses the Minkowski bilinear form $g(\cdot,\cdot) = (\cdot,\cdot)$ on V to convert (1.1) and (1.2) into morphisms

$$\gamma\colon V \to \mathrm{Hom}(S^*, S), \qquad \tilde{\gamma}\colon V \to \mathrm{Hom}(S, S^*) \tag{1.4}$$

i.e., if one defines γ and $\tilde{\gamma}$ by

$$\begin{aligned} t^*\big(\gamma(v)(s^*)\big) &= (\Gamma(s^*,t^*),v) \quad \text{and} \\ \tilde{\gamma}(v)(s)(t) &= (\tilde{\Gamma}(s,t),v), \end{aligned} \tag{1.5}$$

then (1.3) means that for v in V,

$$\begin{aligned} &\tilde{\gamma}(v)\gamma(v)\colon S^* \to S^* \quad \text{and} \\ &\gamma(v)\tilde{\gamma}(v)\colon S \to S \end{aligned}$$

are multiplication by (v,v). In other words, γ and $\tilde{\gamma}$ turn $S \oplus S^*$ into a module over the Clifford algebra $C(V)$. By [I-**Spinors**, §4.9.6] the action of $\mathrm{Spin}(V)$ on S and S^* is induced by this $C(V)$-module structure. Taking the trace in $\tilde{\gamma}(v)\gamma(v) = (v,v)$, i.e., contracting the indices a and c in (1.3), one obtains

$$\Gamma^{\mu}_{ab}\,\tilde{\Gamma}^{\nu ab} + \Gamma^{\nu}_{ab}\,\tilde{\Gamma}^{\mu ab} = 2\,\dim(S) g^{\mu\nu}\ . \tag{1.6}$$

The identity (1.3) also means that if $(v,v) \neq 0$, the inverse of the symmetric bilinear form $(\Gamma(s^*,t^*),v)$ on S^* is the bilinear form $(v,v)^{-1}(\tilde{\Gamma}(s,t),v)$ on S. For v in C, the bilinear form $(\tilde{\Gamma}(s,t),v)$ is hence positive definite, and $\tilde{\Gamma}\colon S \otimes S \to V$ is positive definite.

By the unicity of Γ up to automorphisms of S, it follows that if the representation S of $\mathrm{Spin}(V)$ is isomorphic to its dual, there exists an isomorphism of representations $\alpha\colon S \to S^*$ such that

$$\tilde{\Gamma}(s,t) = \Gamma(\alpha(s),\alpha(t))\ . \tag{1.7}$$

This isomorphism corresponds to a nondegenerate $\mathrm{Spin}(V)$-invariant bilinear form ϵ on S:

$$\epsilon(s,t) = \alpha(s)(t)\ . \tag{1.8}$$

The form $\epsilon(t,s)$ corresponds to the transpose ${}^t\alpha$ of α. One has also $\tilde{\Gamma}(s,t) = \Gamma({}^t\alpha(s), {}^t\alpha(t))$. Indeed, expressing for $(v,v) \neq 0$ that the nondegenerate forms $(v,\tilde{\Gamma}(s,t))$ and $(v,\Gamma(\alpha(s),\alpha(t)))$ have the same inverse, one finds that $\Gamma(s',t') = \tilde{\Gamma}(\alpha^*(s'),\alpha^*(t'))$, with $\alpha^* = {}^t\alpha^{-1}$.

In coordinates, (1.7) reads

$$\boxed{\tilde{\Gamma}^{\mu ab} = \Gamma^{\mu}_{cd}\,\epsilon^{ac}\epsilon^{bd}\ .} \tag{1.9}$$

We now assume that S is irreducible, and is isomorphic to its dual. This excludes the cases where the dimension n of V is congruent to 2 or 6 modulo 8. We treat in turn the cases where the field of endomorphisms Z of S is isomorphic to $\mathbb{R}$ ($n \equiv 1$ or 3 mod 8), $\mathbb{H}$ ($n \equiv 5$ or 7 mod 8) or $\mathbb{C}$ ($n \equiv 0$ or 4 mod 8). Note that (1.7) holds up to a factor for any invariant non-zero bilinear form ϵ.

Case $Z = \mathbb{R}$. Up to a factor, S admits a unique invariant bilinear form, symmetric for $n \equiv 1(\mathrm{mod}8)$ and alternating for $n \equiv 3(\mathrm{mod}8)$. The form ϵ for which (1.7) holds is unique up to sign.

Case $Z = \mathbb{H}$. Up to a factor, S admits a unique invariant bilinear form such that

$$\epsilon(sh, t) = \epsilon(s, t\bar{h}), \qquad h \in H, \quad s, t \in S. \tag{1.10}$$

(We take S to be a *right* $\mathbb{H}$-module.) After extension of scalars from $\mathbb{R}$ to $\mathbb{C}$, S becomes $S_0 \otimes W$, with S_0 irreducible and W of dimension 2, $\mathbb{H}$ becomes $\mathrm{End}(W)$ and (1.10) means that $\epsilon = \epsilon_0 \otimes \epsilon_W$, for ϵ_0 invariant on S_0 and ϵ_W the alternating form on W. The form ϵ for which (1.10) holds is symmetric for $n \equiv 5(\mathrm{mod}8)$ and alternating for $n \equiv 7(\mathrm{mod}8)$, as ϵ_0 is respectively alternating and symmetric [I-**Spinors**, §1.5.1]. The form ϵ for which (1.10) and (1.7) holds is unique up to sign.

Case $Z = \mathbb{C}$. After extension of scalars to $\mathbb{C}$, S becomes the sum of two irreducible representations, orthogonal for $n \equiv 0(\mathrm{mod}8)$ and symplectic for $n \equiv 4(\mathrm{mod}8)$. Over $\mathbb{R}$, the invariant bilinear form will correspondingly be orthogonal or symplectic, with $\epsilon(zs, t) = \epsilon(s, zt)$. The ones for which (1.7) holds form a U_1-orbit under $\epsilon(s, t) \mapsto \epsilon(s, ut)$.

In all cases, ϵ is symmetric or antisymmetric, so that the resulting isomorphism $\alpha\colon S \to S^*$ is unambiguous, at least up to sign. Transporting ϵ by α, we obtain the dual form $\tilde{\epsilon}$ on S^*, which is completely unambiguous. One has[9]

$$\boxed{\epsilon^{ab}\,\tilde{\epsilon}_{cb} = \delta^a_c\ .} \tag{1.11}$$

If the dimension is congruent to 2 or 6 (mod8)—i.e., for the range of dimensions of physical interest, in dimensions 2, 6 and 10—there are super Minkowski spaces based on unequal number of copies of the half spinor representations S^+ and S^-. The invariant pairing in those dimensions is between S^+ and S^-, so that Γ and $\tilde{\Gamma}$ encode information not related by a self-duality pairing.

Let $\mathcal{L}$ be the following super Lie algebra: $\mathcal{L}^0 = V$, $\mathcal{L}^1 = S^*$, the subalgebra $\mathcal{L}^0 = V$ is central, and the Lie bracket on S^* is -2Γ. The corresponding super Lie group $\exp\mathcal{L}$ is defined to be a simply connected super Lie group, given with an isomorphism of $\mathcal{L}$ with its Lie algebra of left-invariant vector fields. It can be constructed as follows. As a space it is

$$\exp\mathcal{L} = V \times \Pi S^*. \tag{1.12}$$

In other words, for any supercommutative ring R the set of its R-points is the even part of $\mathcal{L}_R = \mathcal{L} \otimes R \cong R \otimes \mathcal{L}$. The group law is derived from the Hausdorff formula, which terminates after the quadratic term since $[x, [y, z]] = 0$ for all $x, y, z \in \mathcal{L}$:

$$(v_1, s_1^*)(v_2, s_2^*) = (v_1 + v_2 + \frac{1}{2}[s_1^*, s_2^*], s_1^* + s_2^*), \qquad v_1, v_2 \in R^0 \otimes V, \quad s_1^*, s_2^* \in R^1 \otimes S^*. \tag{1.13}$$

[9] We do not set a convention for raising and lowering a single index with ϵ in the antisymmetric case, since, as was just explained, this involves an arbitrary sign choice we prefer not to make. However, there is no ambiguity in raising and lowering an even number of indices, as for example in (1.10). Thus $\epsilon_{12} = \epsilon^{ab}\epsilon_{1a}\epsilon_{2b} = \epsilon^{21}\epsilon_{12}\epsilon_{21} = -\epsilon^{21} = \epsilon^{12}$. Beware: Many standard physics texts use a different convention here.

The sign rule applied to $[s_1^*, s_2^*] = -2\Gamma(s_1^*, s_2^*)$ gives

$$\frac{1}{2}[b_1 \otimes t_1^*, b_2 \otimes t_2^*] = b_1 b_2 \Gamma(t_1^*, t_2^*), \qquad b_1, b_2 \in R^1, \quad t_1^*, t_2^* \in S^*. \tag{1.14}$$

Fix bases $\{e_\mu\}$ of V and $\{f^a\}$ of $S = (S^*)^*$. As before, write $\Gamma(f_a, f_b) = \Gamma_{ab}^\mu e_\mu$ and $\tilde{\Gamma}(f^a, f^b) = \tilde{\Gamma}^{\mu ab} e_\mu$. In the super Lie algebra $\mathcal{L}$, with the basis $\{e_\mu; f_a\}$, the nontrivial brackets are

$$[f_a, f_b] = -2\Gamma_{ab}^\mu e_\mu. \tag{1.15}$$

In the physics literature it is customary to denote e_μ, f_a by P_μ, Q_a, and we sometimes follow that convention. Thus we write

$$\boxed{[Q_a, Q_b] = -2\Gamma_{ab}^\mu P_\mu.} \tag{1.16}$$

The bases of V^*, S induce a coordinate system (x^μ, θ^a) on $\exp(\mathcal{L}) = V \times \Pi S^*$: the coordinates of $P \in (\mathcal{L} \otimes R)^0$ are $x^\mu(P) = \langle e^\mu, P\rangle$ and $\theta^a(P) = \langle f^a, P\rangle$. If $\{e_\mu\}$ and $\{f_a\}$ are the dual bases of V and S^*, then $P = e_\mu x^\mu + f_a \theta^a$. Let ∂_μ and ∂_a be the corresponding vector fields. In other words,

$$\partial_\mu x^\nu = \delta_\mu^\nu, \qquad \partial_a \theta^b = \delta_a^b, \qquad \partial_\mu \theta^a = \partial_a x^\mu = 0. \tag{1.17}$$

The Lie algebra $\mathcal{L}$ is identified with the tangent space to $\exp \mathcal{L}$ at the origin via $e_\mu \mapsto \partial_\mu \big|_0$ and $f_a \mapsto \partial_a \big|_0$. We shall write D_a for the left-invariant vector field which is ∂_a at 0, and we write τ_{Q_a} for the right-invariant vector field which is ∂_a at 0. Then

$$\begin{aligned} D_a &= \partial_a - \theta^b \Gamma_{ab}^\mu \partial_\mu, \\ \tau_{Q_a} &= \partial_a + \theta^b \Gamma_{ab}^\mu \partial_\mu \end{aligned} \tag{1.18}$$

at a point with coordinates (x^μ, θ^b). The vector fields ∂_μ are both left- and right-invariant. The bracketing relations are

$$\boxed{\begin{aligned} [D_a, D_b] &= -2\Gamma_{ab}^\mu \partial_\mu, \\ [\tau_{Q_a}, \tau_{Q_b}] &= 2\Gamma_{ab}^\mu \partial_\mu, \\ [D_a, \tau_{Q_b}] &= 0. \end{aligned}} \tag{1.19}$$

The D_a, being left-invariant, have the same brackets as the abstract Lie algebra $\mathcal{L}$. (See (1.15).) The vector fields D_a and τ_{Q_b} commute, since right and left translations commute.

Super Minkowski space M is the underlying supermanifold of the super Lie group $\exp(\mathcal{L})$. We let $\exp(\mathcal{L})$ act on the left on M by left translations. Corresponding to (1.12) we have a splitting

$$M = \check{M} \times \Pi S^*. \tag{1.20}$$

It is well-defined since the ambiguity in identifying M with $\exp(\mathcal{L})$ is translation by an element of V, and such translations preserve the splitting (1.12). The *super*

Poincaré group is the semi-direct product $\mathrm{Spin}(V) \ltimes \exp(\mathcal{L})$. It acts on M, and the structures we will consider on M are invariant under the super Poincaré group, with one notable exception—the splitting (1.20). For example, the left translates of $S^* = \mathcal{L}^1 \subset \mathcal{L}$ form a left-invariant distribution τ on M which is preserved by the action of the super Poincaré group.

As we saw, bases $\{e^\mu\}$ of V^* and $\{f^a\}$ of S give a coordinate system (x^μ, θ^a) on $M = \exp(\mathcal{L})$. It will often be more convenient to make computations not in this coordinate system, but rather using the left-invariant moving frame consisting of the vector fields D_a, ∂_μ and/or the dual moving coframe consisting of the 1-forms ρ^a, ω^μ. We fix the sign in the duality by putting vectors to the left of forms:

$$\langle D_a, \rho^b \rangle = \delta_a^b. \tag{1.21}$$

Also, we will sometimes use *complex* bases and so complex vector fields and differential forms. The complex conjugate of a product is the product of the complex conjugates in the *same* order, even for odd elements.[10]

The infinitesimal generators of the action of $\exp(\mathcal{L})$ on M are the right-invariant vector fields. The τ_{Q_a} defined above are a basis of the odd right-invariant vector fields.

A spinor field $\psi = \psi_a f^a$ on $\check{M}$ is a map $\psi\colon \check{M} \to \Pi S$. The form $\tilde{\Gamma}$ then determines a symmetric (in the super sense) bilinear form on spinor fields, which we denote with the usual physicists' notation:

$$\boxed{\psi \not{D} \psi = \tilde{\Gamma}^\mu(\psi, \partial_\mu \psi) = \tilde{\Gamma}^{\mu a b} \psi_a \partial_\mu \psi_b.} \tag{1.22}$$

We call (1.22) the *Dirac form*. One should understand '$\not{D}$' in this formula as the name of the bilinear form, which has been inserted between the arguments. The Dirac form is symmetric up to an exact term, so is exactly symmetric when integrated over $\check{M}$ (assuming no contributions at infinity). A dual spinor field $\lambda = \lambda^a f_a$ is a map $\lambda\colon \check{M} \to \Pi S^*$, and we have the (dual) Dirac form

$$\boxed{\lambda \not{D} \lambda = \Gamma^\mu(\lambda, \partial_\mu \lambda) = \Gamma^\mu_{ab} \lambda^a \partial_\mu \lambda^b.} \tag{1.23}$$

If there is a duality pairing (1.8), then we can use it to identify dual spinors with spinors, in which case (1.23) is identified with (1.22).

We complexify V and S to construct complexified super Minkowski space $M_{\mathbb{C}}$. Inside $\check{M}_{\mathbb{C}}$ we may consider a real affine subspace E of Euclidean signature.[11] However, the complex spinors $S_{\mathbb{C}}$ generally will not carry a real structure under the Euclidean spin group $\mathrm{Spin}(E)$, and so the corresponding superspace $\check{M} \times \Pi S_{\mathbb{C}}$ is complex in odd directions. It is a cs supermanifold [I-**Supersymmetry**, §4.8]. The lagrangians we construct can be rotated to Euclidean space.[12]

[10] See §5 of [I-**Signs**].

[11] With our sign conventions the metric will be *negative* definite; simply change the sign of the metric to obtain a positive definite metric.

[12] For more details on the transition from Minkowski space to Euclidean space, see Chapter 7 of [I-**Classical Fields**].

Remark 1.24. The super Minkowski space is usually called "$N = k$ superspace" if S is the sum of k irreducible real representations of $\mathrm{Spin}(V)$. However, the usage fluctuates somewhat for $n = \dim(\check{M})$ small. For example, if $\dim(\check{M}) = 2$, then $N = 1$ usually means that S is the sum of the two inequivalent real semi-spinorial representations S^+ and S^-. It is more informative to use the notation '$M^{n|s}$' to indicate s supersymmetries in n dimensions, and in dimensions $n \equiv 2(\mathrm{mod}4)$ to use a special notation to indicate the precise spinorial representation: '$M^{n|(s^+,s^-)}$' denotes a super Minkowski space of dimension $n|s^+ + s^-$ built from a spinorial representation which is a sum of copies of S^+, of dimension s^+, and a sum of copies of S^-, of dimension s^-.

The connected group of outer automorphisms of the super Poincaré group which fix the Poincaré subgroup is called the *R-symmetry group*. Dimensional reduction leads to an R-symmetry group (double covering rotations in the extra spatial dimensions), but it is not the only source of R-symmetries.

§1.2. Superfields, component fields, and lagrangians[13]

For many theories we work directly on super Minkowski space M. In such situations we say the theory has a "superspace formulation." Fields on M are called *superfields.* Just as in ordinary nonsupersymmetric field theory, typical examples are: (i) maps $\Phi\colon M \to X$ into an ordinary manifold X; and (ii) a connection A on a principal bundle $\mathcal{P} \to M$. Very often superfields are constrained. For example, in the four-dimensional supersymmetric σ-model we constrain certain derivatives of Φ to vanish, and in all super gauge theories we constrain certain components of the curvature to vanish. The usual elementary facts about functions apply to superfields. Thus elementary operations such as composition, sum, and product produce new superfields. We can also produce new superfields by differentiation, for example by the vector fields D_a, τ_{Q_a} defined in (1.18).

Component fields are ordinary (even and odd) fields on Minkowski space $\check{M}$. We view these as maps from $\check{M}$ into a supermanifold. Given a superfield Φ we define its component fields to be the restriction to $\check{M}$ of certain derivatives of Φ. Let

$$i\colon \check{M} \longrightarrow M \tag{1.25}$$

denote the inclusion of Minkowski space into super Minkowski space. Then for a map $\Phi\colon M \to X$ we typically define a *multiplet* of component fields (ϕ, ψ, F) with formulas like

$$\begin{aligned} \phi &= i^*\Phi, \\ \psi &= i^*D\Phi, \\ F &= i^*DD\Phi. \end{aligned} \tag{1.26}$$

For specific instances, see (4.2) or (5.3). The definition of the multiplet is chosen so that the supermanifold of superfields is isomorphic to the supermanifold of multiplets. (In gauge theory this is an equivalence of categories rather than a bijection.)

[13]Our formulas in this subsection are meant to be suggestive only. Precise versions depend on the particular theory; they appear in subsequent sections. For example, versions of (1.26) for the supersymmetric σ-model appear in (4.7) and (5.7). Also, versions of (1.29) are used in the supersymmetric σ-model to compute (4.14) and (5.10).

In (1.26) ϕ is a map $\phi\colon \check{M} \to X$ and ψ is a spinor field on $\check{M}$ with values in ϕ^*TX. Note that ϕ, F are even, whereas ψ is odd. The field F is *auxiliary* in the sense that for fundamental lagrangians no derivatives of F occur in the lagrangian. The classical field equation for F (keeping the other fields fixed) expresses F in terms of the other fields. In some cases there are no auxiliary fields. If X is flat space, then the formulas (1.26) are well-defined as written. If X is curved, however, then the outermost D in the definition of F acts on a section of the nontrivial bundle Φ^*TX, and we use a covariant derivative to define the action of D. In the simplest supersymmetric σ-models X is Riemannian and we use the pullback of the Levi-Civita connection. In other models different connections on TX (e.g. with torsion) might be more convenient. In gauge theories we meet superfields Λ which are sections of nontrivialized bundles associated to the principal bundle[14] $\mathcal{P}$. A typical example is the adjoint bundle. Since one of the fields in the theory is a connection A on $\mathcal{P}$, we use it to define derivatives of Λ. Our covariant definitions of component fields are global—we never choose coordinate systems or local trivializations.

Component fields are defined in terms of the left-invariant vector fields D. *Lagrangian densities* on superspace are also written in terms of D. Recall that a lagrangian density L is a section of $\operatorname{Dens} M \to \mathcal{F} \times M$, where $\mathcal{F}$ is the supermanifold of superfields and $\operatorname{Dens} M$ the density bundle of M, pulled back over the product. Let η^a be odd parameters. Then the element $\eta^a Q_a$ induces an even vector field ξ on $\mathcal{F} \times M$. The corresponding one-parameter group combines the motion on M with the *inverse*[15] pullback on fields. We now give a general argument that the lagrangians we construct are invariant under ξ. First, note that in our two examples—a σ-model with field $\Phi\colon M \to X$ and a gauge theory with connection A on a principal bundle $\mathcal{P} \to M$ over super Minkowski space—the connection $\mathcal{A}$ we use to define the action of D on superfields depends on the field, so is best thought of as living on $\mathcal{F} \times M$. For example, in the σ-model $\mathcal{A}$ is the pullback of the Levi-Civita connection on TX by the evaluation map $\mathcal{F} \times M \to X$. In the gauge theory $\mathcal{A}$ is the natural "universal connection" on $\mathcal{P} \to \mathcal{F} \times M$ which is flat in the $\mathcal{F}$ direction. Now for the σ-model $\mathcal{A}$ is canonically trivial along the orbits of ξ, since ξ is tangent to the fibers of the evaluation map. Hence the actions of ξ and D on superfields via the connection $\mathcal{A}$ do (super)commute. In the gauge theory case the same statement is true up to gauge transformations, as the reader may verify with some computation (cf. the last remark in [I-**Classical Fields**, §4.2]). Therefore, if L is a gauge-invariant lagrangian density constructed from derivatives of superfields by the D's, then L is invariant under ξ. This is the statement that "L is supersymmetric".

Next, we derive the general shape of the formula for the action of $\eta^a Q_a$ on component fields. We denote a typical component field as

$$f = i^* D^r \Phi. \tag{1.27}$$

Implicit in the notation is that D acts by covariant derivative where necessary. Now $\eta^a Q_a$ generates a diffeomorphism $\exp(\eta^a \tau_{Q_a})$ of M, and we define[16] $\hat{\xi} f$ to be the

[14] In other places we call a 'principal G bundle' a 'G-torsor'.

[15] We use pullback by the *inverse* to achieve a *left* action on functions. This introduces a minus sign into the infinitesimal action of vector fields on functions. For example, ∂_μ acts on functions by *minus* the indicated derivative.

[16] Usually, $\hat{\xi}$ is denoted as 'δ', but since we use 'δ' as the differential along $\mathcal{F}$, we avoid this notation.

change in the component f under this diffeomorphism acting on Φ by the *inverse* pullback:

$$\hat{\xi} f := \frac{D}{Dt}\Big|_{t=0} i^* D^r \big(\exp(-t\eta^a \tau_{Q_a})^* \Phi\big). \tag{1.28}$$

We write 'D/Dt' to indicate that we may have to use a covariant derivative to differentiate in t. To compute (1.28) we use the fact that $\exp(-t\eta^a \tau_{Q_a})$ preserves D and so

$$\begin{aligned} \hat{\xi} f &= \frac{D}{Dt}\Big|_{t=0} i^* \exp(-t\eta^a \tau_{Q_a})^* (D^r \Phi) \\ &= -\eta^a i^* \tau_{Q_a} D^r \Phi \\ &= -\eta^a i^* D_a D^r \Phi. \end{aligned} \tag{1.29}$$

At the last stage we use the fact that τ_{Q_a} and D_a agree when restricted to $\check{M}$. In practice we use (1.29) and the commutation relations among the Ds to express $\hat{\xi} f$ in terms of other component fields.

With our sign conventions, the infinitesimal symmetry P_μ acts on a component field f by $-\partial_\mu f$.

The supersymmetry transformations (1.29) on components obey the commutation relations derived from those of the vector fields Q if there are no covariant derivatives. Explicitly, if η_1, η_2 are odd parameters and $\hat{\xi}_i$ the vector field which corresponds to $\eta_i^a Q_a$, then[17]

$$\boxed{[\hat{\xi}_1, \hat{\xi}_2] f = 2\eta_1^a \eta_2^b \Gamma_{ab}^\mu \partial_\mu f.} \tag{1.30}$$

The same is true for nonlinear σ-models, but in gauge theory these commutation relations may be altered by curvature terms which act as infinitesimal gauge transformations.

The most straightforward way to define a *component lagrangian* for the component fields on $\check{M}$, starting from a lagrangian L in superspace, is to integrate L over the odd variables using the splitting (1.20). In other words, the splitting (1.20) determines a projection

$$\pi\colon M \longrightarrow \check{M} \tag{1.31}$$

and so an integration

$$\pi_*\colon \mathrm{Dens}(M) \longrightarrow \mathrm{Dens}(\check{M}) \tag{1.32}$$

on densities. However, we find better formulas in certain cases by a procedure which differs from this by an exact term. First, the metric on the vector space V determines a canonical positive density $|d^n x|$ on Minkowski space $\check{M}$; it is invariant

[17] In the abstract supersymmetry algebra we have $[\eta_1^a Q_a, \eta_2^b Q_b] = -\eta_1^a \eta_2^b [Q_a, Q_b] = 2\eta_1^a \eta_2^b \Gamma_{ab}^\mu P_\mu$. There are two minus signs which cancel in passing to (1.30): a minus sign for brackets since we use left actions, and the minus sign in the action of P_μ. (See §5 of [I-**Signs**].)

under the Poincaré group. Similarly, if we fix a volume form[18] on S^*, then together with the metric on V this determines a canonical density $|d^n x|\, d^s\theta$ on super Minkowski space M; it is invariant under the super Poincaré group. We define a lagrangian *function* ℓ on M by

$$L = |d^n x|\, d^s\theta\, \ell. \tag{1.33}$$

Then we define the component lagrangian density $|d^n x|\, \check{L}$ on Minkowski space $\check{M}$ by applying some definite combination of D operators, which we symbolically denote 'D^s', to ℓ and then restricting to $\check{M}$:

$$|d^n x|\, \check{L} = |d^n x|\, (i^* D^s \ell). \tag{1.34}$$

Notice that the component lagrangian *function* $\check{L}$ is naturally expressed in terms of component fields. Often i^*D^s is chosen to agree with integration over the odd variables in the splitting (1.20), but in some cases not. For an example of the former (on $M^{3|2}$) see (2.40); for an example of the latter (on $M^{4|4}$) see the discussion following (2.72). In any case we suppose there is a Poincaré invariant differential operator Δ on $\check{M}$ such that

$$\begin{aligned} \pi_* L &= |d^n x|\, (i^* D^s \ell + \Delta i^* \ell) \\ &= |d^n x|\, (\check{L} + \Delta i^* \ell). \end{aligned} \tag{1.35}$$

For integration on $M^{4|4}$ this is formula (2.71).

Theorem 1.36. *The component lagrangian* (1.34) *is supersymmetric.*

Here the supersymmetry is *nonmanifest* so that $\mathrm{Lie}(\hat{\xi})(|d^n x|\, \check{L})$ is a nonzero exact form, where $\hat{\xi}$ is defined in (1.29).

Proof. The verification is easy if the component lagrangian is defined by ordinary integration (1.32), i.e., if $\Delta = 0$ in (1.35):

$$\begin{aligned} \mathrm{Lie}(\hat{\xi})\big(|d^n x|\, \check{L}\big) &= \mathrm{Lie}(\hat{\xi})(\pi_* L) \\ &= \pi_*\big(\mathrm{Lie}(-\eta^a \tau_{Q_a}) L\big) \\ &= \pi_*\big(d\iota(-\eta^a \tau_{Q_a}) L\big) \\ &= d\Big\{\pi_*\big(-\iota(\eta^a \tau_{Q_a}) L\big)\Big\}, \qquad \text{if } \Delta = 0. \end{aligned} \tag{1.37}$$

In this computation 'Lie' denotes the Lie derivative, and we use the Cartan formula $\mathrm{Lie}(\zeta) = d\iota(\zeta) + \iota(\zeta)d$ to define the *integral density*[19] $\iota(\eta^a \tau_{Q_a}) L$. Even if $\Delta \neq 0$ we use (1.35) to rewrite this last expression. First, use (1.31) to lift $|d^n x|$ to M, where

[18] Neither the superspace lagrangian L nor the component lagrangian function $\check{L}$ depends on this volume form; only the superspace lagrangian function ℓ in (1.33) and the choice of D^s in (1.34) depend on it.

[19] See [I-**Supersymmetry**, §§3.9–3.12] for an explanation of integral forms and integral densities.

we view it as a twisted differential form (twisted by the orientation bundle). Then

(1.38)
$$\begin{aligned} d\Big\{\pi_*\big(-\iota(\eta^a Q_a)L\big)\Big\} &= d\Big\{\pi_*\big(-\iota(\eta^a\partial_a + \eta^a\Gamma^\mu_{ab}\theta^b\partial_\mu)\,|d^nx|\,d^s\theta\,\ell\big)\Big\} \\ &= d\Big\{\pi_*\big(-\iota(\eta^a\Gamma^\mu_{ab}\theta^b\partial_\mu)\,|d^nx|\,d^s\theta\,\ell\big)\Big\} \\ &= d\Big\{i^*D^s\big(-\iota(\eta^a\Gamma^\mu_{ab}\theta^b\partial_\mu)\,|d^nx|\,\ell\big) \\ &\qquad\qquad + \Delta i^*\big(-\iota(\eta^a\Gamma^\mu_{ab}\theta^b\partial_\mu)\,|d^nx|\,\ell\big)\Big\} \\ &= d\Big\{i^*D^s\big(-\iota(\eta^a Q_a)\,|d^nx|\,\ell\big)\Big\}. \end{aligned}$$

In the penultimate step we use the fact that $i^*\theta^b = 0$. Now if $\Delta \neq 0$ we write

$$\Delta = \Delta^\mu\partial_\mu \tag{1.39}$$

for some constant differential operators Δ^μ. The expression (1.39) is not in general unique. Now

$$\begin{aligned} \mathrm{Lie}(\hat\xi)\big(|d^nx|\,\Delta i^*\ell\big) &= |d^nx|\,\Delta i^*(-\eta^a\tau_{Q_a}\ell) \\ &= |d^nx|\,\Delta i^*(-\eta^a D_a\ell) \\ &= d\Big\{-\iota(\partial_\mu)\,|d^nx|\,\Delta^\mu i^*(\eta^a D_a\ell)\Big\}. \end{aligned} \tag{1.40}$$

Putting together (1.35), (1.37), (1.38), and (1.40) we have the desired result

(1.41)
$$\mathrm{Lie}(\hat\xi)\big(|d^nx|\,\check L\big) = d\Big\{i^*D^s\big(-\iota(\eta^a Q_a)\,|d^nx|\,\ell\big) - \iota(\partial_\mu)|d^nx|\,\Delta^\mu i^*(\eta^a D_a\ell)\Big\}$$

$$= d\Big\{-\iota(\partial_\mu)|d^nx|\big(\Gamma^\mu_{ab}i^*D^s(\eta^a\theta^b\ell) + \Delta^\mu i^*(\eta^a D_a\ell)\big)\Big\}. \tag{1.42}$$

Equation (1.41) gives one contribution to the *supercurrent*, which is minus the Noether current for the supersymmetry transformation $\hat\xi$. If $\Delta \neq 0$, then the supercurrent we derive from these formulas depends on the choice we make in (1.39).

We sometimes define (a term in) a lagrangian on $\check M$ simply by restricting a scalar superfield S on M to Minkowski space $\check M$:

$$\check L = i^*S. \tag{1.43}$$

Of course, (1.34) is of this form with $S = D^s\ell$. In general (1.43) is *not* supersymmetric. Put differently, only in special cases does the assumption that $\hat\xi S = 0$ on M imply that $\mathrm{Lie}(\hat\xi)(|d^nx|\,\check L)$ is exact.

For some theories there is no (known) superspace formulation and we work directly in components. Then the supersymmetry transformation is defined directly on the component fields and a direct computation is needed to verify that a (component) lagrangian is supersymmetric.

§1.3. A simple example[20]

Here we consider a supersymmetric version of a particle moving on a line. Consider $M^{1|1}$ with coordinates t, θ. The basic odd vector fields are

$$\begin{aligned} D &= \partial_\theta - \theta\partial_t \\ \tau_Q &= \partial_\theta + \theta\partial_t, \end{aligned} \tag{1.44}$$

with $[D, D] = -2\partial_t$ and $[\tau_Q, \tau_Q] = 2\partial_t$. A superfield $\Phi\colon M^{1|1} \to \mathbb{R}$ has components

$$\begin{aligned} \phi &= i^*\Phi, \\ \psi &= i^*D\Phi. \end{aligned} \tag{1.45}$$

Then for $\hat{\xi}$ corresponding to $\eta^a Q_a$ we have, according to (1.29),

$$\begin{aligned} \hat{\xi}\phi &= -\eta i^* D\Phi = -\eta\psi, \\ \hat{\xi}\psi &= -\eta i^* D^2\Phi = \eta\dot{\phi}, \end{aligned} \tag{1.46}$$

where $\dot{\phi} = \partial_t\phi$. We check the bracketing in the simplest case:

$$\begin{aligned} (\hat{\xi}_1\hat{\xi}_2 - \hat{\xi}_2\hat{\xi}_1)\phi &= \hat{\xi}_1(-\eta_2\psi) - \hat{\xi}_2(-\eta_1\psi) \\ &= -\eta_2\hat{\xi}_1\psi + \eta_1\hat{\xi}_2\psi \\ &= -\eta_2\eta_1\dot{\phi} + \eta_1\eta_2\dot{\phi} \\ &= 2\eta_1\eta_2\dot{\phi}. \end{aligned} \tag{1.47}$$

This is consistent with (1.30).[21]

A fundamental lagrangian for the field Φ is

$$L = |dt|\, d\theta \left\{ -\frac{1}{2} D\Phi\partial_t\Phi \right\}. \tag{1.52}$$

[20] Further analysis of this example appears in Problem FP2 of [I-**Homework**].

[21] It is important to use the odd parameter η in these computations. If we write informally

$$\Phi(t, \theta) = \phi(t) + \theta\psi(t), \tag{1.48}$$

then the action of $-\tau_Q$ is

$$-\tau_Q\Phi(t, \theta) = -\psi(t) - \theta\dot{\phi}(t), \tag{1.49}$$

from which we might *erroneously* conclude

$$\begin{aligned} (-\tau_Q)\phi &= -\psi, \\ (-\tau_Q)\psi &= -\dot{\phi}. \end{aligned} \tag{1.50}$$

Multiplying by η we do *not* obtain (1.46). Rather, instead of (1.49) we should write

$$(-\eta\tau_Q)\Phi(t, \theta) = -\eta\psi(t) + \theta\eta\dot{\phi}(t) \tag{1.51}$$

and correctly deduce (1.46). The process of taking components involves odd derivatives; the odd parameter η protects us from making a sign mistake.

The component lagrangian (1.34) is easy to compute:

$$\check{L} = \frac{1}{2}(\dot{\phi})^2 + \frac{1}{2}\psi\dot{\psi}. \tag{1.53}$$

We evaluate the expression in braces on the right hand side of (1.41), noting that $\Delta^i = 0$ in this case:

$$\begin{aligned} i^*D\big(-\iota(\eta Q)\,|dt|\,\frac{-1}{2}D\Phi\partial_t\Phi\big) &= \frac{1}{2}i^*D(\eta\theta D\Phi\partial_t\Phi) \\ &= -\frac{1}{2}\eta\psi\dot{\phi}. \end{aligned} \tag{1.54}$$

We check (1.41) by computing the left hand side directly from (1.53) using (1.46):

$$\begin{aligned} \hat{\xi}\check{L} &= -\big(\partial_t(\eta\psi)\dot{\phi}\big) + \frac{1}{2}\eta\dot{\phi}\dot{\psi} + \frac{1}{2}\psi\partial_t(\eta\dot{\phi}) \\ &= \partial_t(-\frac{1}{2}\eta\psi\dot{\phi}). \end{aligned} \tag{1.55}$$

To compute the Noether current j corresponding to the symmetry $\hat{\xi}$, we first record the variational 1-form computed from (1.53):

$$\gamma = \dot{\phi}\,\delta\phi + \frac{1}{2}\psi\,\delta\psi. \tag{1.56}$$

Then the current is

$$\begin{aligned} j &= \iota(\hat{\xi})\gamma - (-\frac{1}{2}\eta\psi\dot{\phi}) \\ &= (-\eta\dot{\phi}\psi - \frac{1}{2}\eta\psi\dot{\phi}) - (-\frac{1}{2}\eta\psi\dot{\phi}) \\ &= -\eta\dot{\phi}\psi. \end{aligned} \tag{1.57}$$

The *supercharge* is defined to be *minus* the Noether current of Q, which is $\dot{\phi}\psi$.

CHAPTER 2
Coordinates on Superspace

Our task in this chapter is to develop the formalism we need to compute in our main examples: the super Minkowski spaces $M^{3|2}$, $M^{4|4}$, and $M^{6|(8,0)}$. Most of the formulas we derive also apply to differential geometric computations (including spinors) in 3-, 4-, and 6-dimensional ordinary Minkowski space. The entire treatment is based on the exceptional isomorphism of Lie groups in low dimensions, as explained in §2.1. After a brief interlude (§2.2) on dimensional reduction, we flush out the definitions into concrete formulas (§§2.3–2.5). The final section §2.6 records some specializations to dimension 2.

§2.1. $M^{3|2}$, $M^{4|4}$, $M^{6|(8,0)}$ and their complexifications

We consider the cases $n = \dim(V) = 3, 4, 6$ with minimal supersymmetry. The irreducible real spinorial representation S of $\mathrm{Spin}(V)$ is respectively of dimension 2, 4, and 8, with algebra of endomorphisms D respectively reduced to $\mathbb{R}$ ($n = 3$) or isomorphic to $\mathbb{C}$ ($n = 4$) or $\mathbb{H}$ ($n = 6$). Once Γ is chosen, V can be identified with the space of D-hermitian symmetric sesquilinear forms on $\mathrm{Hom}_D(S, D)$, with the positive cone C corresponding to positive forms. Namely, as will be clear from the complexified description, there is up to a real factor a unique symmetric morphism of representations $\Gamma\colon S^* \otimes S^* \to V$, and $\Gamma(\bar{a}s, t) = \Gamma(s, at)$ for $a \in D$. It induces an isomorphism of the symmetric quotient of $S^* \otimes_D S^*$ with V. This symmetric quotient is identified with the space of hermitian symmetric sesquilinear forms on $\mathrm{Hom}_D(S^*, D)$ by letting $s \otimes s$ correspond to the form for which $\langle \alpha, \alpha \rangle = \overline{\alpha(s)}\alpha(s)$.

For V of dimension 4 (resp. 6) we choose an isomorphism of D with $\mathbb{C}$ (resp. $\mathbb{H}$). On the super Minkowski space M, the odd distribution τ then carries a $\mathbb{C}$- (resp left $\mathbb{H}$-) module structure, and this structure is respected by the super Poincaré group.

Over $\mathbb{C}$ one has the exceptional isomorphisms $\mathrm{Spin}(3) \cong SL(2)$, $\mathrm{Spin}(4) \cong SL(2) \times SL(2)$, and $\mathrm{Spin}(6) \cong SL(4)$. They make it convenient to describe the complexification $V_{\mathbb{C}}$ of V starting from that of S. The real form V corresponds to a complex conjugation (antilinear automorphism) of $V_{\mathbb{C}}$.

After complexification $\mathbb{H}$ becomes isomorphic to the 2×2 matrix algebra. This enables us to fix a 2-dimensional complex vector space W, a pseudoreal (or quaternionic) structure on W (that is, an antilinear automorphism j of square -1) and an isomorphism of $\mathbb{H}$ with the ring of real endomorphisms of W, where 'real' means commuting with j. We take these endomorphisms to act on the right. The model

is $W = \mathbb{H}$, the complex structure given by left multiplication by $I \in \mathbb{H}$, the pseudoreal structure given by left multiplication by $J \in \mathbb{H}$. The real endomorphisms are the right multiplications by elements of $\mathbb{H}$. We fix on W a symplectic form ϵ_W which is real, i.e., $\epsilon_W(jx, jy) = \overline{\epsilon_W(x, y)}$, and positive definite, i.e., $\epsilon_W(x, jx) > 0$ for $x \neq 0$. Note that $\epsilon_W(x, jx)$ is real since ϵ_W is. We also fix a basis $\{e', e'' = j(e')\}$ of W for which $\epsilon_W(e', e'') = 1$ and for which the $\mathbb{H}$-module multiplication by $z \in \mathbb{C} \subset \mathbb{H}$ is the multiplication by z on e' and by $\bar{z}$ on e''. In the model $W = \mathbb{H}$ introduced above, one may take $e' = 1$, $e'' = J$.

We now describe, case by case, the complexifications $S_{\mathbb{C}}$ and $V_{\mathbb{C}}$ of S and V, the symmetric morphism $\Gamma\colon S^*_{\mathbb{C}} \otimes S^*_{\mathbb{C}} \to V_{\mathbb{C}}$, the D-module structure of $S_{\mathbb{C}}$, and $\mathrm{Spin}(V)$. For $M^{3|2}$ and $M^{4|4}$ we give a duality pairing $\epsilon\colon S_{\mathbb{C}} \otimes S_{\mathbb{C}} \to \mathbb{C}$. In these cases the form $\tilde{\Gamma}\colon S_{\mathbb{C}} \otimes S_{\mathbb{C}} \to V_{\mathbb{C}}$ is constructed using (1.9). (We write the formulas in coordinates below.) For $M^{6|(8,0)}$ there is no duality pairing and we specify $\tilde{\Gamma}$ explicitly.

3|2 *case.* Here there is no need to complexify to have a nice description, so we directly describe the real superspace. Let S be a 2-dimensional real vector space with volume form ϵ, viewed as a symplectic form on S. Define

$$\begin{aligned} V &= \mathrm{Sym}^2(S^*) \\ \Gamma\colon (s, t) &\longmapsto st \end{aligned} \tag{2.1}$$

There is an isomorphism $SL(S) \xrightarrow{\sim} \mathrm{Spin}(V)$.

4|4 *case.* Let S' and S'' be 2-dimensional complex vector spaces with complex volume forms ϵ', ϵ'', viewed as symplectic forms on S' and S''. Define

$$\begin{aligned} S_{\mathbb{C}} &= S' \oplus S'' \\ V_{\mathbb{C}} &= S'^* \otimes S''^* \\ \Gamma\colon (s'_1 + s''_1,\ s'_2 + s''_2) &\longmapsto (s'_1 \otimes s''_2) + (s'_2 \otimes s''_1) \\ \epsilon &= \epsilon' \oplus \epsilon'' \end{aligned} \tag{2.2}$$

The $\mathbb{C}$-module structure of $S_{\mathbb{C}}$ is multiplication by z on S' and multiplication by $\bar{z}$ on S''. There is an isomorphism $SL(S') \times SL(S'') \xrightarrow{\sim} \mathrm{Spin}(V_{\mathbb{C}})$. The bigger group

$$\{\langle g, g' \rangle\colon \det g' \det g'' = 1\} \subset GL(S') \times GL(S'') \tag{2.3}$$

acts on S' and S'' preserving the bilinear form on $V_{\mathbb{C}}$. The new symmetry in this larger group is called *R-symmetry*; it is the action of unit norm scalars in $\mathbb{C}$ on $S_{\mathbb{C}}$.

6|(8, 0) *case.* Let S_0 be a 4-dimensional complex vector space with a volume form ν. Define

$$\begin{aligned} S_{\mathbb{C}} &= S_0 \otimes W \\ S^*_{\mathbb{C}} &= S_0^* \otimes W^* \\ V_{\mathbb{C}} &= \textstyle\bigwedge^2 S_0^* \\ \Gamma\colon (w' \otimes s'_0, w'' \otimes s''_0) &\longmapsto \epsilon_W(w', w'')\ (s'_0 \wedge s''_0) \end{aligned} \tag{2.4}$$

Then $SL(S_0) \xrightarrow{\sim} \mathrm{Spin}(V_{\mathbb{C}})$. The $\mathbb{H}$-module structure of $S_{\mathbb{C}}$ is induced from that of W. The volume form ν defines a symmetric pairing

$$\nu\colon \textstyle\bigwedge^2 S_0 \otimes \bigwedge^2 S_0 \longrightarrow \mathbb{C} \tag{2.5}$$

via which we identify $V_{\mathbb{C}}$ and $V_{\mathbb{C}}^*$. This pairing is, up to a factor of 2, the inverse metric. Then $\tilde{\Gamma}$ is the composition

$$\tilde{\Gamma}\colon S_{\mathbb{C}} \otimes S_{\mathbb{C}} \longrightarrow V_{\mathbb{C}}^* \longrightarrow V_{\mathbb{C}}, \tag{2.6}$$

where the first arrow is given by a similar formula to that above for Γ and the second arrow is constructed from the duality (2.5). To make (1.3) valid, we define the metric g on $V_{\mathbb{C}}$ to be such that the inverse metric g^{-1} on $\bigwedge^2 S_0$ is $\nu/2$.

In the $6|(8,0)$ and $4|4$ cases we now describe the real spaces V and S as fixed points of an antilinear involution of $V_{\mathbb{C}}$ and $S_{\mathbb{C}}$. The real structures must be compatible with Γ as well as with the symmetric bilinear form g on $V_{\mathbb{C}}$. The induced form on V is to have signature $(1, n-1)$. The real description above of the $3|2$ case does give the required signature $(1,2)$.

$4|4$ *case.* Fix an antilinear involution on $S_{\mathbb{C}} = S' \oplus S''$ which exchanges S' and S'' as well as ϵ' and ϵ''. From this we get real structures on $S_{\mathbb{C}}$ and $V_{\mathbb{C}} = S'^* \otimes S''^*$. The symmetric bilinear form $\epsilon' \otimes \epsilon''$ is real of signature $(1,3)$.

$6|(8,0)$ *case.* A pseudoreal structure j on S_0 induces real structures on $S_{\mathbb{C}} = S_0 \otimes W$ as well as on $V = \bigwedge^2 S_0^*$. The vector $s \wedge j(s) \in V$ is real for any $s \in S_0$, since

$$j\big(s \wedge j(s)\big) = j(s) \wedge (-s) = s \wedge j(s). \tag{2.7}$$

If the volume form ν on S_0 is real, the corresponding quadratic form on the real form of V has signature $(1,5)$ or $(5,1)$. We want (ν, j) to give signature $(1,5)$. This is the case if, for a basis of S_0 of the form $\{s, j(s), t, j(t)\}$, one has $\nu\big(s \wedge j(s) \wedge t \wedge j(t)\big)$ real and positive.

§2.2. Dimensional reduction

The super Lie algebras $V^3 \oplus S^2$ and $V^4 \oplus S^4$ underlying $M^{3|2}$ and $M^{4|4}$ are related as follows. Any (real) isomorphism $\alpha\colon \mathbb{C} \otimes_{\mathbb{R}} S^2 \xrightarrow{\sim} S^4$ extends to $\alpha\colon V^3 \hookrightarrow V^4$. The image $\alpha(V^3)$ is the orthogonal complement to a one-dimensional spacelike subspace $T \subset V^4$. This defines embeddings

$$M^{3|2} \hookrightarrow M^{4|4} \tag{2.8}$$

$$M^{3|2} \hookrightarrow M^{3|4} := M^{4|4}/T. \tag{2.9}$$

Notice that the left translates of S'' form a two-dimensional integrable odd distribution Λ on $M^{3|4}$, and the composition of (2.9) with the quotient map

$$M^{3|2} \hookrightarrow M^{3|4} \longrightarrow M^{3|4}/\Lambda \tag{2.10}$$

is an isomorphism.

Similarly, for the super Lie algebras $V^4 \oplus S^4$ and $V^6 \oplus S^8$ underlying $M^{4|4}$ and $M^{6|(8,0)}$, any isomorphism $\alpha\colon \mathbb{H} \otimes_{\mathbb{C}} S^4 \xrightarrow{\sim} S^8$ extends to $\alpha\colon V^4 \hookrightarrow V^6$. The image $\alpha(V^4) \subset V^6$ is the orthogonal complement to a two-dimensional spacelike subspace T. This defines embeddings

$$\begin{aligned} &M^{4|4} \hookrightarrow M^{6|(8,0)} \\ &M^{4|4} \hookrightarrow M^{4|(8,0)} := M^{6|(8,0)}/T. \end{aligned} \tag{2.11}$$

After complexification we obtain the following picture.

$M^{3|2} \hookrightarrow M^{4|4}$. The map α complexifies to isomorphisms $\alpha'\colon S^2_{\mathbb{C}} \to S'$ and $\alpha''\colon S^2_{\mathbb{C}} \to S''$ which are complex conjugate. The extension to $V^3_{\mathbb{C}} = \mathrm{Sym}^2(S^2_{\mathbb{C}})^*$ is given by $st \mapsto \alpha'(s)\alpha''(t)+\alpha'(t)\alpha''(s)$, with values in $S^4_{\mathbb{C}} = S'\otimes S''$. Identify $S'\otimes S'' \cong S^2_{\mathbb{C}}\otimes S^2_{\mathbb{C}}$ via (α', α''); then the subspace $T \subset S'^* \otimes S''^*$ is identified with the orthogonal complement $\bigwedge^2(S^2_{\mathbb{C}})^*$ to $\mathrm{Sym}^2(S^2_{\mathbb{C}})^*$. The extension $\alpha\colon V^3_{\mathbb{C}} \to V^4_{\mathbb{C}}$ is compatible with the symmetric bilinear forms if, for some λ, we have

$$\alpha'(\epsilon) = \frac{\lambda}{\sqrt{2}}\,\epsilon', \qquad \alpha''(\epsilon) = \frac{\lambda^{-1}}{\sqrt{2}}\,\epsilon''. \tag{2.12}$$

$M^{4|4} \hookrightarrow M^{6|(8,0)}$. The complexified map $\alpha\colon S^4_{\mathbb{C}} = S'\oplus S'' \to S^8_{\mathbb{C}} = S_0\otimes W$ is of the form $\alpha' e' + \alpha'' e''$ for $(\alpha', \alpha'')\colon S'\oplus S'' \to S_0$ an isomorphism. (Recall that $\{e', e''\}$ is a fixed basis of W.) The reality of our starting α means that the pseudoreal structure of $\mathrm{Hom}(S'\oplus S'', S_0)$ transforms α' into α'' and α'' into $-\alpha'$. The extension to $V^4_{\mathbb{C}} = S'^*\otimes S''^*$ is given by $s'\otimes s'' \mapsto \alpha'(s')\wedge\alpha''(s'')$, with values in $\bigwedge^2 S^*_0$. The subspace T of $\bigwedge^2 S^*_0$ is $\bigwedge^2\alpha'(S')\oplus\bigwedge^2\alpha''(S'')$. The extension $\alpha\colon V^4_{\mathbb{C}} \to V^6_{\mathbb{C}}$ will be compatible with the symmetric bilinear forms if

$$\alpha'(\epsilon')\wedge\alpha''(\epsilon'') = -\nu_0. \tag{2.13}$$

§2.3. Coordinates on $M^{3|2}$

A basis $\{f_a\}_{a=1,2}$ of S^* gives a basis $\{e_{ab} = f_a f_b\}$ of $V = \mathrm{Sym}^2(S^*)$. Notice that $e_{ab} = e_{ba}$. We use on V^* not the dual basis, but rather the basis $\{f^a f^b\}$ of $\mathrm{Sym}^2(S)$, in duality with $\mathrm{Sym}^2(S^*)$ by

$$\langle s^*t^*, st\rangle = \frac{1}{2}\big(\langle s^*, s\rangle\langle t^*, t\rangle + \langle s^*, t\rangle\langle t^*, s\rangle\big), \qquad s^*, t^* \in S^*, \quad s, t \in S. \tag{2.14}$$

The chosen bases of V and S^* give us vector fields ∂_a and ∂_{ab}, while the bases of V^* and S give us coordinates y^{ab} and θ^a. We have

$$\begin{aligned} \partial_a\theta^b &= \delta_a^b \\ \partial_{ab}y^{a'b'} &= \frac{1}{2}(\delta_a^{a'}\delta_b^{b'} + \delta_a^{b'}\delta_b^{a'}) \\ \partial_a y^{bc} &= \partial_{ab}\theta^c = 0. \end{aligned} \tag{2.15}$$

The left-invariant vector field D_a corresponding to f_a in S^* is

$$D_a = \partial_a - \theta^b\partial_{ab}, \tag{2.16}$$

and corresponding to the relation $[f_a, f_b] = -2f_{ab}$ in the Lie algebra we have

$$[D_a, D_b] = -2\partial_{ab}. \tag{2.17}$$

The right-invariant vector field τ_{Q_a} corresponding to f_a in S^* is

$$\tau_{Q_a} = \partial_a + \theta^b\partial_{ab} \tag{2.18}$$

with

$$[\tau_{Q_a}, \tau_{Q_b}] = 2\partial_{ab}. \tag{2.19}$$

We use $\{D_a, \partial_{ab}\}$ as a left-invariant moving frame on $M^{3|2}$. Dually, we use the left-invariant moving coframe $\{\rho^a, \omega^{ab}\}$ which has the following duality pairings:[22]

$$\begin{aligned} \langle D_a, \rho^b \rangle &= \delta_a^b \\ \langle \partial_{ab}, \omega^{a'b'} \rangle &= \frac{1}{2}(\delta_a^{a'}\delta_b^{b'} + \delta_a^{b'}\delta_b^{a'}) \\ \langle D_a, \omega^{bc} \rangle &= \langle \partial_{ab}, \rho^c \rangle = 0. \end{aligned} \tag{2.20}$$

This gives

$$\begin{aligned} \rho^a &= d\theta^a \\ \omega^{ab} &= dy^{ab} - (\theta^a d\theta^b + \theta^b d\theta^a) \end{aligned} \tag{2.21}$$

and

$$\begin{aligned} d\rho^a &= 0 \\ d\omega^{ab} &= -2\rho^a \wedge \rho^b. \end{aligned} \tag{2.22}$$

The symmetric pairing Γ is

$$\Gamma(f_a, f_b) = e_{ab} = \Gamma_{ab}^{(a'b')} e_{a'b'} = \frac{1}{2}(\delta_a^{a'}\delta_b^{b'} + \delta_a^{b'}\delta_b^{a'}) e_{a'b'}. \tag{2.23}$$

Similarly,

$$\tilde{\Gamma}(f^a, f^b) = \tilde{\Gamma}^{(a'b')ab} e_{a'b'} = \frac{1}{2}(\epsilon^{a'a}\epsilon^{b'b} + \epsilon^{a'b}\epsilon^{b'a}) e_{a'b'}, \tag{2.24}$$

where ϵ is the volume form on S. The metric on Minkowski space $\check{M}^3$ is the restriction to $\check{M}^3$ of

$$g = 2\epsilon_{aa'}\epsilon_{bb'}\, \omega^{ab} \otimes \omega^{a'b'}, \tag{2.25}$$

and the inverse metric is the restriction to $\check{M}^3$ of

$$g^{-1} = \frac{1}{2}\epsilon^{aa'}\epsilon^{bb'}\, \partial_{ab} \otimes \partial_{a'b'}. \tag{2.26}$$

The reader should be mindful of (2.20), which implies

$$g^{-1}(\omega^{ab}, \omega^{cd}) = \frac{1}{4}(\epsilon^{ac}\epsilon^{bd} + \epsilon^{ad}\epsilon^{bc}). \tag{2.27}$$

The Dirac form (1.22) on a spinor field $\psi = \psi_a f^a$ is

$$\psi \not{D} \psi = -\epsilon^{ab}\epsilon^{cd}\psi_a \partial_{bc} \psi_d. \tag{2.28}$$

[22] Recall the convention (1.21) that we write vector fields to the left of differential forms: $\langle D, df \rangle = \iota(D)df = Df$, where $\iota(D)$ is contraction by the vector field D.

The Dirac form (1.22) on a dual spinor field $\lambda = \lambda^a f_a$ is

$$\lambda \not{D} \lambda = \lambda^a \partial_{ab} \lambda^b. \tag{2.29}$$

We also introduce the Dirac *operator*

$$(\not{D}\psi)_b = -\epsilon^{cd} \partial_{bc} \psi_d, \tag{2.30}$$

which maps sections of S to sections of S. Then (2.28) and (2.30) are compatible in the sense that

$$\epsilon^{ab} \chi_a (\not{D}\psi)_b = \chi \not{D} \psi \tag{2.31}$$

for all spinor fields χ, ψ. The wave operator is

$$\begin{aligned} \Box &= \frac{1}{2} \epsilon^{aa'} \epsilon^{bb'} \partial_{ab} \partial_{a'b'} \\ &= \partial_{11}\partial_{22} - \partial_{12}^2; \end{aligned} \tag{2.32}$$

the sign is chosen so that the continuation to Euclidean space is the nonnegative laplace operator. The reader may check that $\not{D}^2 = -\Box$.

Preferred bases are those for which $\epsilon^{12} = -\epsilon^{21} = 1$ and hence[23] $\epsilon_{12} = -\epsilon_{21} = 1$. Put differently, $\epsilon(f_1, f_2) = 1$. In a preferred basis we introduce standard coordinates $\{x^\mu\}$ on Minkowski space $\check{M}^3$ by

$$\begin{aligned} y^{11} &= \frac{x^0 + x^1}{2}, \\ y^{22} &= \frac{x^0 - x^1}{2}, \\ y^{12} &= \frac{x^2}{2}. \end{aligned} \tag{2.33}$$

The dual vector fields ∂_{ab} and $\partial/\partial x^\mu$ are related by

$$\begin{aligned} \partial_{11} &= \frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^1}, \\ \partial_{22} &= \frac{\partial}{\partial x^0} - \frac{\partial}{\partial x^1}, \\ \partial_{12} &= \frac{\partial}{\partial x^2}. \end{aligned} \tag{2.34}$$

The metric (2.25) is then the standard metric

$$g = (dx^0)^2 - (dx^1)^2 - (dx^2)^2. \tag{2.35}$$

The density $|d^3x|$ in (1.33) is the standard positive density

$$|d^3x| = |dx^0 dx^1 dx^2|. \tag{2.36}$$

[23] See (1.11).

It is convenient in our gauge theory computations to work with the Hodge $*$ operator. For this we stipulate that x^0, x^1, x^2 is a positively oriented coordinate system.[24] Then the volume form is the restriction to $\tilde{M}^3$ of

$$\text{vol} = 4\omega^{11} \wedge \omega^{12} \wedge \omega^{22} \tag{2.37}$$

and the $*$ operator acts by

$$\begin{aligned} *\omega^{ab} &= \epsilon_{a'b'}\, \omega^{aa'} \wedge \omega^{bb'} \\ *(\omega^{aa'} \wedge \omega^{bb'}) &= \frac{1}{4}\left(\epsilon^{ab}\omega^{a'b'} + \epsilon^{ab'}\omega^{a'b} + \epsilon^{a'b}\omega^{ab'} + \epsilon^{a'b'}\omega^{ab}\right). \end{aligned} \tag{2.38}$$

For integration we introduce the notation

$$\int d^2\theta = -\frac{1}{2}\int d\theta^a d\theta^b \epsilon_{ab} = \int d\theta^2 d\theta^1. \tag{2.39}$$

Explicit in (2.39) is our choice of volume form on spinors. Using the identification between integration and differentiation, we write

$$\int d^2\theta = i^*\partial_2\partial_1 = -\frac{1}{2} i^*\epsilon^{ab} D_a D_b = -i^* D^2 = \frac{1}{2} i^*(D_2 D_1 - D_1 D_2), \tag{2.40}$$

where we have introduced the notation

$$D^2 = \frac{1}{2}\epsilon^{ab} D_a D_b. \tag{2.41}$$

The following relations are easy to verify:

$$D_a D_b = -(\partial_{ab} - \epsilon_{ab} D^2) \tag{2.42}$$

$$D_a D^2 = -D^2 D_a = -\epsilon^{bc}\partial_{ab} D_c \tag{2.43}$$

$$D^2 D^2 = -\Box \tag{2.44}$$

§2.4. Coordinates on $M^{4|4}$

We fix a basis $\{f_a\}_{a=1,2}$ of S'^* and let $\{\bar{f}_{\dot{a}}\}$ be the complex conjugate basis of S''^*. As a basis for $V_{\mathbb{C}} = S'^* \otimes S''^*$ we use $\{e_{a\dot{b}} = f_a \bar{f}_{\dot{b}}\}$. Those bases and their duals give us a complex coordinate system $\{y^{a\dot{b}}, \theta^a, \bar{\theta}^{\dot{a}}\}$, with θ^a and $\bar{\theta}^{\dot{a}}$ complex conjugate and $y^{a\dot{b}}$ the complex conjugate to $y^{b\dot{a}}$. We write ∂_a, $\bar{\partial}_{\dot{a}}$, and $\partial_{a\dot{b}}$ for the corresponding vector fields. Note our convention about indices: If both 'a' and '$\dot{a}$' appear in the same equation, then $a = \dot{a}$. This allows us to write reality conditions and equations such as (2.78) and (2.79).

Let D_a and $\overline{D}_{\dot{a}}$ be the left-invariant complex vector fields corresponding to f_a and $\bar{f}_{\dot{a}}$. One has

$$\begin{aligned} D_a &= \partial_a - \bar{\theta}^{\dot{b}}\partial_{a\dot{b}} \\ \overline{D}_{\dot{a}} &= \bar{\partial}_{\dot{a}} - \theta^b \partial_{b\dot{a}}. \end{aligned} \tag{2.45}$$

[24] The orientation makes it easier to write intermediate steps in some of our computations, but the final formulas do not depend on it.

Corresponding to the relations $[f_a, f_b] = [\bar{f}_{\dot{a}}, \bar{f}_{\dot{b}}] = 0$ and $[f_a, \bar{f}_{\dot{b}}] = -2e_{a\dot{b}}$ in the Lie algebra, one has

$$\begin{aligned} [D_a, D_b] &= [\overline{D}_{\dot{a}}, \overline{D}_{\dot{b}}] = 0 \\ [D_a, \overline{D}_{\dot{b}}] &= -2\partial_{a\dot{b}}. \end{aligned} \tag{2.46}$$

The similar right-invariant complex vector fields are

$$\begin{aligned} \tau_{Q_a} &= \partial_a + \bar{\theta}^{\dot{b}}\partial_{a\dot{b}} \\ \tau_{\overline{Q}_{\dot{a}}} &= \bar{\partial}_{\dot{a}} + \theta^b\partial_{b\dot{a}} \end{aligned} \tag{2.47}$$

with

$$\begin{aligned} [\tau_{Q_a}, \tau_{Q_b}] &= [\tau_{\overline{Q}_{\dot{a}}}, \tau_{\overline{Q}_{\dot{b}}}] = 0 \\ [\tau_{Q_a}, \tau_{\overline{Q}_{\dot{b}}}] &= 2\partial_{a\dot{b}}. \end{aligned} \tag{2.48}$$

We use $\{D_a, \overline{D}_{\dot{a}}, \partial_{a\dot{b}}\}$ as a left-invariant moving frame and also use the dual left-invariant moving coframe $\{\rho^a, \overline{\rho}^{\dot{a}}, \omega^{a\dot{b}}\}$. The nontrivial duality pairings are

$$\begin{aligned} \langle D_a, \rho^b\rangle &= \delta_a^b \\ \langle \overline{D}_{\dot{a}}, \overline{\rho}^{\dot{b}}\rangle &= \delta_{\dot{a}}^{\dot{b}} \\ \langle \partial_{a\dot{b}}, \rho^{a'\dot{b}'}\rangle &= \delta_a^{a'}\,\delta_{\dot{b}}^{\dot{b}'}. \end{aligned} \tag{2.49}$$

One has

$$\begin{aligned} \rho^a &= d\theta^a \\ \overline{\rho}^{\dot{a}} &= d\bar{\theta}^{\dot{a}} \\ \omega^{a\dot{b}} &= dy^{a\dot{b}} - (\bar{\theta}^{\dot{b}}d\theta^a + \theta^a d\bar{\theta}^{\dot{b}}) \end{aligned} \tag{2.50}$$

and, dual to the bracket relations,

$$\begin{aligned} d\rho^a = d\overline{\rho}^{\dot{a}} &= 0 \\ d\omega^{a\dot{b}} &= -2\rho^a \wedge \overline{\rho}^{\dot{b}}. \end{aligned} \tag{2.51}$$

The symmetric pairing Γ is

$$\Gamma(f_a, \bar{f}_{\dot{b}}) = e_{a\dot{b}} = \Gamma_{a\dot{b}}^{(a'\dot{b}')}e_{a'\dot{b}'} = (\delta_a^{a'}\delta_{\dot{b}}^{\dot{b}'})e_{a'\dot{b}'}. \tag{2.52}$$

Similarly, the symmetric pairing $\tilde{\Gamma}$ is

$$\tilde{\Gamma}(f^a, \bar{f}^{\dot{b}}) = \tilde{\Gamma}^{(a'\dot{b}')a\dot{b}}e_{a'\dot{b}'} = (\epsilon^{a'a}\epsilon^{\dot{b}'\dot{b}})e_{a'\dot{b}'}, \tag{2.53}$$

where ϵ is the volume form on S', S''. The metric on Minkowski space $\check{M}^4$ is the restriction to $\check{M}^4$ of

$$g = 2\epsilon_{aa'}\epsilon_{\dot{b}\dot{b}'}\,\omega^{a\dot{b}} \otimes \omega^{a'\dot{b}'}, \tag{2.54}$$

and the inverse metric is the restriction to $\check{M}^4$ of

$$g^{-1} = \frac{1}{2}\epsilon^{aa'}\epsilon^{\dot{b}\dot{b}'}\,\partial_{a\dot{b}} \otimes \partial_{a'\dot{b}'}. \tag{2.55}$$

The Dirac form (1.22) pairs a positive chirality spinor field $\psi = \psi_a f^a$ with a negative chirality spinor field $\bar{\psi} = \bar{\psi}_{\dot{a}}\bar{e}^{\dot{a}}$:

$$\bar{\psi}\not{D}\psi = \epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\bar{\psi}_{\dot{d}}\partial_{a\dot{c}}\psi_b. \tag{2.56}$$

The Dirac form (1.22) on dual spinor fields is

$$\bar{\lambda}\not{D}\lambda = \bar{\lambda}^{\dot{b}}\partial_{a\dot{b}}\lambda^a. \tag{2.57}$$

The Dirac operator on spinor fields exchanges sections of S' and S'':

$$\begin{aligned}(\not{D}\psi)_{\dot{c}} &= -\epsilon^{ab}\partial_{a\dot{c}}\psi_b\\ (\not{D}\bar{\psi})_c &= -\epsilon^{\dot{a}\dot{b}}\partial_{c\dot{a}}\bar{\psi}_{\dot{b}}\end{aligned} \tag{2.58}$$

The wave operator is

$$\begin{aligned}\Box &= \frac{1}{2}\epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\partial_{a\dot{c}}\partial_{b\dot{d}}\\ &= \partial_{1\dot{1}}\partial_{2\dot{2}} - \partial_{1\dot{2}}\partial_{2\dot{1}},\end{aligned} \tag{2.59}$$

and $\not{D}^2 = -\Box$.

Preferred bases are those for which $\epsilon'(f_1, f_2) = \epsilon''(\bar{f}_{\dot{1}}, \bar{f}_{\dot{2}}) = 1$. In a preferred basis we introduce standard coordinates $\{x^\mu\}$ on Minkowski space V by

$$\begin{aligned}y^{1\dot{1}} &= \frac{x^0 + x^1}{2},\\ y^{2\dot{2}} &= \frac{x^0 - x^1}{2},\\ y^{1\dot{2}} &= \frac{x^2 + ix^3}{2},\\ y^{2\dot{1}} &= \frac{x^2 - ix^3}{2}.\end{aligned} \tag{2.60}$$

The dual vector fields $\partial_{a\dot{b}}$ and $\partial/\partial x^\mu$ are related by

$$\begin{aligned}\partial_{1\dot{1}} &= \frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^1},\\ \partial_{2\dot{2}} &= \frac{\partial}{\partial x^0} - \frac{\partial}{\partial x^1},\\ \partial_{1\dot{2}} &= \frac{\partial}{\partial x^2} - i\frac{\partial}{\partial x^3},\\ \partial_{2\dot{1}} &= \frac{\partial}{\partial x^2} + i\frac{\partial}{\partial x^3}.\end{aligned} \tag{2.61}$$

The metric (2.54) is then the standard metric

$$g = (dx^0)^2 - (dx^1)^2 - (dx^2)^2 - (dx^3)^2. \tag{2.62}$$

The density $|d^4x|$ in (1.33) is the standard positive density

$$|d^4x| = |dx^0 dx^1 dx^2 dx^3|. \tag{2.63}$$

To define self-duality we need to orient $\check{M}^4$, and so we stipulate that x^0, x^1, x^2, x^3 is an oriented coordinate system.[25] Then for a symmetric tensor F_{ab} the 2-form

$$\frac{1}{2}F_{ab}\epsilon_{\dot{c}\dot{d}}\,\omega^{a\dot{c}} \wedge \omega^{b\dot{d}} \tag{2.64}$$

is self-dual, and for a symmetric tensor $\overline{F}_{\dot{c}\dot{d}}$ the 2-form

$$\frac{1}{2}\epsilon_{ab}\overline{F}_{\dot{c}\dot{d}}\,\omega^{a\dot{c}} \wedge \omega^{b\dot{d}} \tag{2.65}$$

is anti-self-dual. Our convention is that a self-dual 2-form σ satisfies $*\sigma = -\sqrt{-1}\,\sigma$. Then the analytic continuation to Euclidean space gives the usual notion of self-duality. The complex conjugate of a self-dual form (in Minkowski space) is anti-self-dual.

For the 2-form

$$F = \frac{1}{2}(F_{ab}\epsilon_{\dot{c}\dot{d}} + \epsilon_{ab}\overline{F}_{\dot{c}\dot{d}})\,\omega^{a\dot{c}} \wedge \omega^{b\dot{d}} \tag{2.66}$$

we have the formulas

$$\begin{aligned} |F|^2 &= \frac{1}{4}(F_{ab}F_{cd}\epsilon^{ac}\epsilon^{bd} + \overline{F}_{\dot{a}\dot{b}}\overline{F}_{\dot{c}\dot{d}}\epsilon^{\dot{a}\dot{c}}\epsilon^{\dot{b}\dot{d}}), \\ F \wedge F &= \frac{i}{4}(F_{ab}F_{cd}\epsilon^{ac}\epsilon^{bd} - \overline{F}_{\dot{a}\dot{b}}\overline{F}_{\dot{c}\dot{d}}\epsilon^{\dot{a}\dot{c}}\epsilon^{\dot{b}\dot{d}})\,d^4x, \end{aligned} \tag{2.67}$$

where d^4x is the standard volume form

$$d^4x = dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3. \tag{2.68}$$

In general the 2-form F is complex; then '$|F|^2$' in (2.67) is not a norm, but rather is a quadratic form. The 2-form F is real if $\overline{F}_{\dot{a}\dot{b}} = \overline{F_{ab}}$.

We mention the fact that the analytic continuation of the $*$ operator on 2-forms in Minkowski space of *any* dimension is $-\sqrt{-1}$ times the $*$ operator on 2-forms in Euclidean space.

A complex function $\Phi\colon M^{4|4} \to \mathbb{C}$ is *chiral* if $\overline{D}_{\dot{a}}\Phi = 0$ and *antichiral* if $D_a\Phi = 0$.

Following (2.39) we introduce the notations

$$\begin{aligned} \int d^2\bar{\theta} &= -\frac{1}{2}\int d\bar{\theta}^{\dot{a}}d\bar{\theta}^{\dot{b}}\epsilon_{\dot{a}\dot{b}} = \int d\bar{\theta}^{\dot{2}}d\bar{\theta}^{\dot{1}} \\ \int d^4\theta &= \int d^2\theta d^2\bar{\theta}. \end{aligned} \tag{2.69}$$

[25] The orientation only serves to choose which of (2.64) and (2.65) below is self-dual and which is anti-self-dual. It does not enter in a fundamental way in our formulae.

We discuss the meaning of $\int d^2\theta$ and $\int d^2\bar\theta$ in §5.3; in any case we compute them using (2.40) and the analogous formula with the conjugate vector fields. Note that we will use the notation

$$\overline{D}^2 = \frac{1}{2}\epsilon^{\dot a\dot b}\overline{D}_{\dot a}\overline{D}_{\dot b}, \tag{2.70}$$

analogous to (2.41). For integration over all odd coordinates we find the formulas

$$\begin{aligned}\int d^4\theta &= i^*\partial_2\partial_1\bar\partial_{\dot 2}\bar\partial_{\dot 1}\\ &= \frac{1}{2}\, i^*\left\{D_1\overline{D}_{\dot 1}\overline{D}_{\dot 2}D_2 + D_2\overline{D}_{\dot 2}\overline{D}_{\dot 1}D_1\right\} - \Box i^* \qquad (2.71)\\ &= \frac{1}{2}\, i^*\left\{D^2\overline{D}^2 + \overline{D}^2D^2\right\} + \Box i^*. \qquad (2.72)\end{aligned}$$

In each of the last two lines the term in braces disagrees with the integral by a multiple of the wave operator. In a lagrangian this wave operator gives an exact term which is usually omitted anyway. The particular combination of derivatives in the first term of (2.71) has the virtue that it exactly annihilates *chiral* and *antichiral* superfields. This is an advantage over the usual integral and leads to nicer formulas. Hence we compute component lagrangians (1.34) using this improved expression, which is the usual integral plus the wave operator composed with restriction to V. Formula (2.72) is useful in some computations. Both of these integration formulas follow from direct computation.

The following relations are easy to verify.

$$D_aD^2 = D^2D_a = \overline{D}_{\dot a}\overline{D}^2 = \overline{D}^2\overline{D}_{\dot a} = 0 \tag{2.73}$$

$$D_a\overline{D}^2 = \overline{D}^2D_a - 2\epsilon^{\dot b\dot c}\partial_{a\dot b}\overline{D}_{\dot c} \tag{2.74}$$

$$\overline{D}_{\dot a}D^2 = D^2\overline{D}_{\dot a} - 2\epsilon^{bc}\partial_{b\dot a}D_c \tag{2.75}$$

$$\overline{D}^2D^2 = -4\Box \qquad \text{on chiral superfields} \tag{2.76}$$

$$D^2\overline{D}^2 = -4\Box \qquad \text{on antichiral superfields} \tag{2.77}$$

Next, we describe the dimensional reduction map $M^{3|2} \hookrightarrow M^{4|4}$. Decorate the bases (of spinors and vectors) and coordinates in $M^{3|2}$ with a '$\hat{}$'. We take the inclusion to be compatible with the ϵ and Γ tensors:

$$\begin{aligned}\hat f_a &\longmapsto \frac{f_a + \bar f_{\dot a}}{\sqrt{2}}\\ \hat e_{ab} &\longmapsto \frac{e_{a\dot b} + e_{b\dot a}}{2}.\end{aligned} \tag{2.78}$$

The image of the inclusion on spinors is the subspace where $\theta^a = \bar\theta^{\dot a}$ and both are real, described by the equations

$$\theta^a = \bar\theta^{\dot a} = \frac{\hat\theta^a}{\sqrt{2}}. \tag{2.79}$$

The image of the inclusion on vectors is the subspace where $y^{1\dot{2}} = y^{2\dot{1}}$, or equivalently where $x^3 = 0$, described by the equations

$$\begin{aligned} y^{1\dot{1}} &= \hat{y}^{11} \\ y^{2\dot{2}} &= \hat{y}^{22} \\ y^{1\dot{2}} = y^{2\dot{1}} &= \hat{y}^{12}. \end{aligned} \tag{2.80}$$

As an exercise the reader may check that (2.74)–(2.75) reduce to (2.42)–(2.44) under dimensional reduction.

The R-symmetry is generated by an even vector field R with brackets

$$\begin{aligned} [R, D_a] &= D_a \\ [R, \overline{D}_{\dot{a}}] &= -\overline{D}_{\dot{a}} \\ [R, \partial_{a\dot{b}}] &= 0. \end{aligned} \tag{2.81}$$

Note that R is purely imaginary: $\overline{R} = -R$.

§2.5. Coordinates on $M^{6|(8,0)}$

A basis $\{f_a\}_{a=1,2,3,4}$ of S_0^* gives us a basis $e'^* \otimes f_a, e''^* \otimes f_a$ of $S_{\mathbb{C}}^*$ and a basis $\{e_{ab} = f_a \wedge f_b\}_{a<b}$ of $V_{\mathbb{C}}$. For the basis elements of $S_{\mathbb{C}}^*$ we use the notation 'f_{ai}', where $f_{a1} = e'^* \otimes f_a$ and $f_{a2} = e''^* \otimes f_a$. Let y^{ab}, θ^{ai} be the corresponding coordinate system. For $b < a$ define $e_{ba} = -e_{ab}$ and $y^{ba} = -y^{ab}$. Write ∂_{ai} and ∂_{ab} for the corresponding complex vector fields on $M^{6|(8,0)}$.

Let ϵ_{ij} denote the symplectic form on the vector space W^*. We use our standard choice $\epsilon_{12} = 1$.

The left-invariant complex vector field D_{ai} corresponding to e_{ai} is

$$D_{ai} = \partial_{ai} - \epsilon_{ij}\theta^{bj}\partial_{ab} \tag{2.82}$$

with brackets

$$[D_{ai}, D_{bj}] = -2\epsilon_{ij}\partial_{ab}. \tag{2.83}$$

The similar right-invariant vector fields are

$$\tau_{Q_{ai}} = \partial_{ai} + \epsilon_{ij}\theta^{bj}\partial_{ab} \tag{2.84}$$

with

$$[\tau_{Q_{ai}}, \tau_{Q_{bj}}] = 2\epsilon_{ij}\partial_{ab}. \tag{2.85}$$

We will use $\{D_{ai}, \partial_{ab}\}$ as a left-invariant moving frame with dual coframe $\{\rho^{ai}, \omega^{ab}\}$. The nonzero duality pairings are

$$\begin{aligned} \langle D_{ai}, \rho^{bj} \rangle &= \delta_a^b \delta_i^j \\ \langle \partial_{ab}, \omega^{cd} \rangle &= \delta_a^c \delta_b^d - \delta_a^d \delta_b^c. \end{aligned} \tag{2.86}$$

One has

$$\begin{aligned} \rho^{ai} &= d\theta^{ai} \\ \omega^{ab} &= dy^{ab} + \epsilon_{ij}\theta^{ai}d\theta^{bj} - \epsilon_{ij}\theta^{bi}d\theta^{aj} \end{aligned} \tag{2.87}$$

with differentials

$$\begin{aligned} d\rho^{ai} &= 0 \\ d\omega^{ab} &= 2\epsilon_{ij}\,\rho^{ai}\wedge\rho^{bj}. \end{aligned} \tag{2.88}$$

The symmetric pairing Γ is

$$\Gamma(f_{ai}, f_{bj}) = \epsilon_{ij}e_{ab} = \frac{1}{2}\Gamma^{(pq)}_{(ai)(bj)}e_{pq} = \frac{1}{2}\big[\epsilon_{ij}(\delta^p_a\delta^q_b - \delta^q_a\delta^p_b)\big]e_{pq}. \tag{2.89}$$

The symmetric pairing $\tilde{\Gamma}$ is

$$\tilde{\Gamma}(f^{ai}, f^{bj}) = \frac{1}{2}\tilde{\Gamma}^{(pq)(ai)(bj)}e_{pq} = \frac{1}{2}(\epsilon^{ij}\nu^{pqab})e_{pq}, \tag{2.90}$$

where ν is the volume form. The metric on Minkowski space $\check{M}^6$ is the restriction to $\check{M}^6$ of

$$g = \frac{1}{2}\nu_{pqrs}dy^{pq}\otimes dy^{rs} = \sum_{\substack{p<q\\ r<s}} 2\nu_{pqrs}\omega^{pq}\otimes\omega^{rs}, \tag{2.91}$$

and the inverse metric is the restriction to $\check{M}^6$ of

$$g^{-1} = \frac{1}{8}\nu^{pqrs}\,\partial_{pq}\otimes\partial_{rs} = \sum_{\substack{p<q\\ r<s}} \frac{1}{2}\nu^{pqrs}\,\partial_{pq}\otimes\partial_{rs}. \tag{2.92}$$

The volume form and its inverse are related by $\nu^{1234}\nu_{1234} = 1$. The Dirac form (1.22) on spinor fields $\psi = \psi_{ai}f^{ai}$ is

$$\psi\not{D}\psi = \frac{1}{2}\nu^{abcd}\epsilon^{ij}\psi_{ai}\partial_{bc}\psi_{dj} \tag{2.93}$$

On dual spinor fields $\lambda = \lambda^{ai}f_{ai}$ we have the Dirac form (1.23)

$$\lambda\not{D}\lambda = \epsilon_{ij}\lambda^{ai}\partial_{ab}\lambda^{bj}. \tag{2.94}$$

Preferred bases are those for which $\nu^{1234} = -1$. We also use the dual form with $\nu_{1234} = -1$. A real structure is given by the pseudoreal structure on S_0 for which $f_3 = j(f_1)$ and $f_4 = j(f_2)$. Then

$$\begin{aligned} \overline{D_{11}} &= \quad D_{32} \\ \overline{D_{21}} &= \quad D_{42} \\ \overline{D_{31}} &= -D_{12} \\ \overline{D_{41}} &= -D_{22}. \end{aligned} \tag{2.95}$$

In a preferred basis we introduce standard coordinates $\{x^\mu\}$ on Minkowski space $\check{M}^6$ by

$$\begin{aligned} y^{12} &= \frac{x^4 + ix^5}{2}, \\ y^{13} &= \frac{x^0 + x^1}{2}, \\ y^{14} &= \frac{x^2 + ix^3}{2}, \\ y^{23} &= \frac{x^2 - ix^3}{2}, \\ y^{24} &= \frac{x^0 - x^1}{2}, \\ y^{34} &= \frac{x^4 - ix^5}{2}. \end{aligned} \tag{2.96}$$

The dual vector fields ∂_{ab} and $\partial/\partial x^\mu$ are related by

$$\begin{aligned} \partial_{12} &= \frac{\partial}{\partial x^4} - i\frac{\partial}{\partial x^5}, \\ \partial_{13} &= \frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^1}, \\ \partial_{14} &= \frac{\partial}{\partial x^2} - i\frac{\partial}{\partial x^3}, \\ \partial_{23} &= \frac{\partial}{\partial x^2} + i\frac{\partial}{\partial x^3}, \\ \partial_{24} &= \frac{\partial}{\partial x^0} - \frac{\partial}{\partial x^1}, \\ \partial_{34} &= \frac{\partial}{\partial x^4} + i\frac{\partial}{\partial x^5}. \end{aligned} \tag{2.97}$$

The metric (2.91) is then the standard metric

$$g = (dx^0)^2 - (dx^1)^2 - (dx^2)^2 - (dx^3)^2 - (dx^4)^2 - (dx^5)^2. \tag{2.98}$$

Next, we describe the dimensional reduction map $M^{4|4} \hookrightarrow M^{6|(8,0)}$. Denote the bases (of spinors and vectors) and coordinates in $M^{6|(8,0)}$ with a '~'. We take the inclusion maps to be

$$\begin{aligned} f_1 &\longmapsto \tilde{f}_{1(1)} & e_{1\dot{1}} &\longmapsto \tilde{e}_{13} \\ f_2 &\longmapsto \tilde{f}_{2(1)} & e_{2\dot{2}} &\longmapsto \tilde{e}_{24} \\ \bar{f}_{\dot{1}} &\longmapsto \tilde{f}_{3(2)} & e_{1\dot{2}} &\longmapsto \tilde{e}_{14} \\ \bar{f}_{\dot{2}} &\longmapsto \tilde{f}_{4(2)} & e_{2\dot{1}} &\longmapsto \tilde{e}_{23} \end{aligned} \tag{2.99}$$

The image of the inclusion on spinors is described by the equations

$$\begin{aligned} \theta^{1(1)} &= \theta^1 & \theta^{1(2)} &= 0 \\ \theta^{2(1)} &= \theta^2 & \theta^{2(2)} &= 0 \\ \theta^{3(1)} &= 0 & \theta^{3(2)} &= \bar{\theta}^{\dot{1}} \\ \theta^{4(1)} &= 0 & \theta^{4(2)} &= \bar{\theta}^{\dot{2}}. \end{aligned} \tag{2.100}$$

The image of the inclusion on vectors is described by the equations

$$\begin{aligned} \tilde{y}^{12} &= 0 & \tilde{y}^{23} &= y^{2\dot{1}} \\ \tilde{y}^{13} &= y^{1\dot{1}} & \tilde{y}^{24} &= y^{2\dot{2}} \\ \tilde{y}^{14} &= y^{1\dot{2}} & \tilde{y}^{34} &= 0 \end{aligned} \tag{2.101}$$

There is an R-symmetry group $SU(2)_R$ which acts on $M^{6|(8,0)}$. The action of a matrix $S = (S_i^j)$ is

$$D_{ai} \longrightarrow D_{ai'} S_i^{i'}. \tag{2.102}$$

Demanding that this preserve (2.83) requires $S \in SL(2;\mathbb{C})$; asking that the real structure also be preserved reduces us to $SU(2)$. More conceptually, this $SU(2)_R = Sp(1)_R$ symmetry group is the group of real endomorphisms of the complex symplectic vector space W.

There is also a dimensional reduction map $M^{4|8} \hookrightarrow M^{6|(8,0)}$, described simply by equations (2.101). Usually $M^{4|8}$ is described with vector fields $\{D_a^{(i)}, \overline{D}_{\dot{a}(j)}\}_{\substack{a=1,2 \\ i,j=1,2}}$ with bracketing (2.46) supplemented by a δ-function:

$$[D_a^{(i)}, \overline{D}_{\dot{a}(j)}] = -2\delta_j^i \partial_{a\dot{a}}. \tag{2.103}$$

One possible correspondence between these vector fields and the $\{\tilde{D}_{ai}\}_{\substack{a=1,2,3,4 \\ i=1,2}}$ is:

$$\begin{aligned} \tilde{D}_{11} &= D_1^{(1)} & \tilde{D}_{12} &= D_1^{(2)} \\ \tilde{D}_{21} &= D_2^{(1)} & \tilde{D}_{22} &= D_2^{(2)} \\ \tilde{D}_{31} &= -\overline{D}_{\dot{1}(2)} & \tilde{D}_{32} &= \overline{D}_{\dot{1}(1)} \\ \tilde{D}_{41} &= -\overline{D}_{\dot{2}(2)} & \tilde{D}_{42} &= \overline{D}_{\dot{2}(1)} \end{aligned} \tag{2.104}$$

A new feature of $M^{4|8}$ is an enlarged R-symmetry group $U(2)_R$. It acts trivially on the even variables. The representation of $U(2)_R$ on each of the pairs $\{D_a^{(1)}, D_a^{(2)}\}$ of odd vector fields is the standard one, whereas on each of the pairs $\{\overline{D}_{\dot{a}(1)}, \overline{D}_{\dot{a}(2)}\}$ of odd vector fields $U(2)_R$ acts by the conjugate to the standard action. Under the dimensional reduction $M^{4|4} \hookrightarrow M^{4|(8,0)}$, specified by (2.100), only the subgroup $U(1)_R \subset U(2)_R$ given by (2.81) survives. Notice that this subgroup is the induced action by the double cover in the spin group of rotations in the x^4-x^5 plane.

§2.6. Low dimensions

The basic superspace $M^{1|1}$ in dimension one is built starting from a one-dimensional odd space S and letting $V = S^{\otimes(-2)}$. In terms of the obvious left-invariant vector fields D and ∂ we have

$$[D, D] = -2\partial. \tag{2.105}$$

Analogs of other formulas we have developed above are easy to come by. Dimensional reduction from the higher dimensional spaces sets all spacelike coordinates

of Minkowski space to zero leaving just the time coordinate x^0. In this way we also obtain spaces $M^{1|2}$, $M^{1|4}$, and $M^{1|8}$ with more odd variables. As these spaces are reduced from $M^{3|2}$, $M^{4|4}$, and $M^{6|(8,0)}$, we obtain R-symmetry groups Spin(2), Spin(3), and Spin(5) from rotations in the additional spatial directions.

The basic[26] superspace $M^{2|(1,1)}$ in dimension two is simply a product

$$M^{2|(1,1)} = M^{1|1} \times M^{1|1}. \tag{2.106}$$

(Note that the metric (2.109) is not a product, however.) This corresponds to the fact that there are real semi-spinors in 2 dimensions (termed *Majorana-Weyl spinors* in the physics literature). For S^+, S^- real one-dimensional vector spaces, we set $S = S^+ \oplus S^-$ and $V = (S^+)^{\otimes(-2)} \oplus (S^-)^{\otimes(-2)}$ with the obvious bilinear pairing. We can see $M^{2|(1,1)}$ as coming from $M^{3|2}$ by dimensional reduction: in $M^{3|2}$ set $y^{12} = 0$. It is customary to use '+' for the odd index '1' and '−' for the odd index '2'. (These indices appear beginning with formula (2.15).) Also, we set $y^+ = y^{11}$ and $y^- = y^{22}$ with similar notation $\partial_+ = \partial_{11}$ and $\partial_- = \partial_{22}$.

$N = 2$ superspace in 2 dimensions is $M^{2|(2,2)}$, and we view it as embedded in $M^{4|4}$ (dimensional reduction) by setting $y^{1\dot{2}} = y^{2\dot{1}} = 0$, or equivalently $x^2 = x^3 = 0$. The real description of $M^{2|(1,1)}$ makes clear that there is also a real description of $M^{2|(2,2)}$, but we use the complex vector fields and forms induced from $M^{4|4}$ instead. Again we use '+' for the odd index '1' and '−' for the odd index '2' (which appear beginning with formula (2.45).) The '+' and'−' on the bosonic coordinates are the usual lightcone coordinates in two dimensions. Thus the left-invariant odd vector fields are $D_+, \overline{D}_+, D_-, \overline{D}_-$, and we denote the left-invariant even vector fields by ∂_+, ∂_-. The bracketing relations are

$$\begin{aligned} [D_+, D_+] = [D_+, D_-] = [D_+, \overline{D}_-] = [\overline{D}_+, \overline{D}_+] = 0 \\ [D_-, D_-] = [\overline{D}_+, \overline{D}_-] = [D_-, \overline{D}_+] = [\overline{D}_-, \overline{D}_-] = 0 \\ [D_+, \overline{D}_+] = -2\partial_+ \\ [D_-, \overline{D}_-] = -2\partial_-. \end{aligned} \tag{2.107}$$

For convenience we record the nonzero components of $\Gamma, \tilde{\Gamma}$:

$$\begin{aligned} \Gamma^+_{++} = \Gamma^-_{--} = 1 \\ \tilde{\Gamma}^{+--} = \tilde{\Gamma}^{-++} = 1. \end{aligned} \tag{2.108}$$

The formulas in $M^{4|4}$ all apply, though many simplify. For example, the metric (2.54), the inverse metric (2.55), the Dirac form (2.56), the Dirac operator (2.58), and the wave operator (2.59) reduce to

$$g = 2(dy^+ \otimes dy^- + dy^- \otimes dy^+) \tag{2.109}$$

$$g^{-1} = \frac{1}{2}(\partial_+ \otimes \partial_- + \partial_- \otimes \partial_+) \tag{2.110}$$

$$\bar{\psi} \not{D} \psi = \bar{\psi}_+ \partial_- \psi_+ + \bar{\psi}_- \partial_+ \psi_- \tag{2.111}$$

$$(\not{D}\psi)_+ = -\partial_+ \psi_- \qquad (\not{D}\psi)_- = \partial_- \psi_+ \tag{2.112}$$

$$(\not{D}\bar{\psi})_+ = -\partial_+ \bar{\psi}_- \qquad (\not{D}\bar{\psi})_- = \partial_- \bar{\psi}_+ \tag{2.113}$$

$$\Box = \partial_+ \partial_-. \tag{2.114}$$

[26] $M^{2|(1,1)}$ is usually termed $N = 1$ superspace, though it is not the minimal possibility—there is a superspace $M^{2|(s^+,s^-)}$ for any nonnegative integers s^+, s^-.

The relationship to standard coordinates x^0, x^1 is

$$\begin{aligned} y^+ &= \frac{x^0 + x^1}{2}, \\ y^- &= \frac{x^0 - x^1}{2}, \end{aligned} \tag{2.115}$$

and

$$\begin{aligned} \partial_+ &= \frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^1}, \\ \partial_- &= \frac{\partial}{\partial x^0} - \frac{\partial}{\partial x^1}. \end{aligned} \tag{2.116}$$

There is a 2-dimensional space of R-symmetries with basis J_+, J_- described by the brackets

$$\begin{aligned} &[J_+, D_+] = \quad D_+ && [J_-, D_-] = \quad D_- \\ &[J_+, \overline{D}_+] = -\overline{D}_+ && [J_-, \overline{D}_-] = -\overline{D}_- \\ &[J_+, D_-] = [J_+, \overline{D}_-] = 0 && [J_-, D_+] = [J_-, \overline{D}_+] = 0. \end{aligned} \tag{2.117}$$

The R-symmetry (2.81) in $M^{4|4}$ induces $J_+ + J_-$ on $M^{2|(2,2)}$. The double cover in the spin group of rotation in the x^2-x^3 plane in $M^{4|4}$ induces the R-symmetry $J_+ - J_-$ of $M^{2|(2,2)}$.

As in $M^{4|4}$, a complex function $\Phi\colon M^{2|(2,2)} \to \mathbb{C}$ is *chiral* if

$$\overline{D}_+\Phi = \overline{D}_-\Phi = 0; \tag{2.118}$$

then the complex conjugate $\overline{\Phi}$ is *antichiral*. There is a new possibility as well. Namely, a complex function $\Sigma\colon M^{2|(2,2)} \to \mathbb{C}$ is *twisted chiral* if

$$\overline{D}_+\Sigma = D_-\Sigma = 0; \tag{2.119}$$

then the complex conjugate $\overline{\Sigma}$ is *twisted antichiral*.

Formulas (2.71) and (2.72) for computing $\int d^4\theta$ are, of course, valid on $M^{2|(2,2)}$. Again we use $\int d^4\theta + \Box i^*$ to compute component lagrangians when chiral and antichiral fields are involved. When twisted chiral and twisted antichiral fields are involved, it is more convenient to use $\int d^4\theta - \Box i^*$, which equals the first term in (2.72). This has the advantage that the expression $D^2\overline{D}^2 + \overline{D}^2 D^2$ annihilates both twisted chiral and twisted antichiral fields. In a more complicated (term in a) lagrangian involving both chiral and twisted chiral fields one would have to simply choose a procedure and then deal with the exact terms which would most likely appear.

CHAPTER 3
Supersymmetric σ-Models

A scalar field is a map $\phi\colon \check{M}^n \to X$ from ordinary n-dimensional Minkowski space to a manifold X. If X is a linear space, then a theory with the field ϕ is called a linear σ-model; if X is curved, then it is a nonlinear σ-model. In supersymmetric σ-models there is in addition a spinor field. After some linear algebra preliminaries in §3.1, we begin in §3.2 with the linear case, where we show that a supersymmetric extension of the ordinary free σ-model exists in dimensions $n = 3, 4, 6$. We label these three cases according to the ring $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ which underlies the spin group $SL(2, \mathbb{F})$. In the nonlinear case (X a general Riemannian manifold) supersymmetry imposes constraints on the target manifold X: For $\mathbb{F} = \mathbb{C}$ it must be Kähler and for $\mathbb{F} = \mathbb{H}$ it must be hyperkähler. The sufficiency of these constraints is explained at the beginning of §3.3, after which we derive the exact form of the component lagrangian. In the bosonic case any function $V\colon X \to \mathbb{R}$ may serve as a potential energy function, but supersymmetry constrains the form of V. The constraints are described in §3.4, though the proofs are deferred to future chapters. We make some brief remarks about the superspace formulation in §3.5 and about the relationship among the different theories in §3.6.

§3.1. Preliminary remarks on linear algebra

We discuss simultaneously the fields $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$. Since $\mathbb{H}$ is noncommutative, we must be careful about the order of multiplication. Let $^-$ denote the standard conjugation on $\mathbb{F}$, which is trivial for $\mathbb{F} = \mathbb{R}$. For $x, y \in \mathbb{F}$ we have

$$\overline{xy} = \bar{y}\bar{x}. \tag{3.1}$$

The extension of (3.1) to superalgebra has a sign, in accordance with the sign rule. It is useful to note that:

$$\mathrm{Re}(\lambda\mu) = \mathrm{Re}(\mu\lambda) = \mathrm{Re}(\bar{\lambda}\bar{\mu}) = \mathrm{Re}(\bar{\mu}\bar{\lambda}), \qquad \lambda, \mu \in \mathbb{F}. \tag{3.2}$$

Let V be the vector space of translations of $\check{M}^n$ ($n = 3, 4, 6$). If S is an irreducible real spinorial representation of $\mathrm{Spin}(V)$, the field $\mathbb{F}$ of endomorphisms of the representation S is respectively isomorphic to $\mathbb{R}$, $\mathbb{C}$, or $\mathbb{H}$; the vector space S is of dimension 2 over $\mathbb{F}$; and $\mathrm{Spin}(V) \cong SL_{\mathbb{F}}(S)$. We choose S and a *right* vector space structure over $\mathbb{F} = \mathbb{R}, \mathbb{C}$ or $\mathbb{H}$ commuting with the action of $\mathrm{Spin}(V)$. The dual $S^* = \mathrm{Hom}_{\mathbb{F}}(S, \mathbb{F})$ is then a *left* $\mathbb{F}$-vector space, and $\mathrm{Spin}(V) \cong SL_{\mathbb{F}}(S^*)$ as

well. We write $\overline{S}$ for S with the left vector space structure $\lambda s := s\bar\lambda$ and $\overline{S^*}$ for S^* with the right vector space structure $s^*\lambda := \bar\lambda s^*$. The $\mathbb{F}$-dual S^* can be identified with the $\mathbb{R}$-dual $\mathrm{Hom}_{\mathbb{R}}(S,\mathbb{R})$ by $\alpha \mapsto$ real part of the $\mathbb{F}$-linear form α. If $\{f^a\}$ is a basis of S over $\mathbb{F}$, then there are induced bases $\{\bar f^{\dot a}\}$, $\{f_a\}$, $\{\bar f_{\dot a}\}$ of $\overline{S}$, S^*, $\overline{S^*}$.

As explained at the beginning of §2.1 there are invariant symmetric $\mathbb{R}$-bilinear forms $\Gamma\colon S^*\otimes_{\mathbb{R}} S^* \to V$ and $\tilde\Gamma\colon S\otimes_{\mathbb{R}} S\to V$. They factor through $\overline{S^*}\otimes_{\mathbb{F}} S^*$ and $S\otimes_{\mathbb{F}}\overline{S}$, respectively. In general, if A, B, C are respectively $\mathbb{F}$ right, $\mathbb{F}$ left, $\mathbb{R}$ vector spaces, then an $\mathbb{R}$-linear map $T\colon A\otimes_{\mathbb{F}} B\to C$ gives rise to an $(\mathbb{F},\mathbb{F})$-bilinear form $T_1\colon A\otimes_{\mathbb{R}} B\to C\otimes_{\mathbb{R}}\mathbb{F}$, characterized by

$$T(a,b) = \mathrm{Re}\big(T_1(a,b)\big). \tag{3.3}$$

Conversely, the real part of any $(\mathbb{F},\mathbb{F})$-bilinear T_1 obeys

$$\mathrm{Re}\,T_1(a\lambda,b) = \mathrm{Re}\,T_1(a,b)\lambda = \mathrm{Re}\,\lambda T_1(a,b) = \mathrm{Re}\,T_1(a,\lambda b), \tag{3.4}$$

so factors through $A\otimes_{\mathbb{F}} B$. We apply this construction to Γ and $\tilde\Gamma$, and denote by $\Gamma^\mu_{a\dot b}$ and $\tilde\Gamma^{\mu\dot a b}$ the coordinates of Γ_1 and $\tilde\Gamma_1$ in bases $\{e_\mu\}$ of V and $\{f^a\}$ of S, up to a factor:

$$\begin{aligned}\Gamma(x^a f_a, y^b f_b) &= \frac{1}{\kappa^2}\,\mathrm{Re}(y^b\Gamma^\mu_{b\dot a}\overline{x^a})\,e_\mu,\\ \tilde\Gamma(f^a x_a, f^b y_b) &= \quad\ \mathrm{Re}(\overline{y_b}\tilde\Gamma^{\mu\dot b a}x_a)\,e_\mu.\end{aligned} \tag{3.5}$$

Here κ is the factor

$$\kappa = \begin{cases} 1, & \mathbb{F}=\mathbb{R};\\ \sqrt{2}, & \mathbb{F}=\mathbb{C},\mathbb{H}.\end{cases} \tag{3.6}$$

On the left hand side of these equations, $\Gamma,\tilde\Gamma$ denote the real forms acting on elements of the underlying real vector space. Thus, for example, we might write $\Gamma\big(\mathrm{Re}(x^a f_a),\mathrm{Re}(y^b f_b)\big)$ to distinguish it from (2.52). The factor of κ^2 is introduced to make these formulas compatible with (2.52) and similar formulas of Chapter 2. The symmetry of Γ and $\tilde\Gamma$ becomes the hermiticity property

$$\begin{aligned}\overline{\Gamma^\mu_{a\dot b}} &= \Gamma^\mu_{b\dot a},\\ \overline{\tilde\Gamma^{\mu\dot a b}} &= \tilde\Gamma^{\mu\dot b a}.\end{aligned} \tag{3.7}$$

The Clifford relation (1.3) with this normalization is

$$\Gamma^\mu_{a\dot b}\tilde\Gamma^{\nu\dot b c} + \Gamma^\nu_{a\dot b}\tilde\Gamma^{\mu\dot b c} = 2g^{\mu\nu}\delta^c_a \tag{3.8}$$

and its conjugate. We emphasize that $\Gamma^\mu_{a\dot b}, \tilde\Gamma^{\mu\dot a b}$ lie in $\mathbb{F}$, whereas $g_{\mu\nu}, g^{\mu\nu}$ are real.

For the quaternion case $\mathbb{F}=\mathbb{H}$ we can choose bases so that the pairing Γ is given by

$$\begin{aligned}&\Gamma^0_{1\dot1}=1, \quad \Gamma^1_{1\dot1}=1,\\ &\Gamma^0_{2\dot2}=1, \quad \Gamma^1_{2\dot2}=-1,\\ &\Gamma^2_{1\dot2}=1, \quad \Gamma^3_{1\dot2}=-i, \quad \Gamma^4_{1\dot2}=j, \quad \Gamma^5_{1\dot2}=k,\end{aligned} \tag{3.9}$$

where i, j, k are the usual quaternions. Then the pairing $\tilde{\Gamma}$ is given by

$$\begin{aligned} &\Gamma^{0\dot{1}1} = 1, \qquad \Gamma^{1\dot{1}1} = -1, \\ &\Gamma^{0\dot{2}2} = 1, \qquad \Gamma^{1\dot{2}2} = 1, \\ &\Gamma^{2\dot{2}1} = -1, \quad \Gamma^{3\dot{2}1} = -i, \quad \Gamma^{4\dot{2}1} = j, \quad \Gamma^{5\dot{2}1} = k. \end{aligned} \tag{3.10}$$

In this basis the metric has the standard form (2.98). It is useful to note that the vector $\Gamma(f_a, f_a)$ is lightlike (has vanishing norm square) for all a; this property will be important in Chapter 6. The pairings for the complex case $\mathbb{F} = \mathbb{C}$ are obtained by omitting the entries with j, k; those for the real case $\mathbb{F} = \mathbb{R}$ by taking the real part of all entries. Then the chosen bases agree with those in §2.3, §2.4. Recall from (1.9) that for $\mathbb{F} = \mathbb{R}, \mathbb{C}$ we have a skew tensor ϵ, and the pairings $\tilde{\Gamma}$ and Γ are related by

$$\tilde{\Gamma}^{\mu\dot{b}a} = \Gamma^{\mu}_{c\dot{d}} \epsilon^{ac} \epsilon^{\dot{b}\dot{d}}, \qquad \mathbb{F} = \mathbb{R}, \mathbb{C}. \tag{3.11}$$

In §3.3 we have use for the following. Suppose V_R is a right $\mathbb{H}$ vector space and V_L a left $\mathbb{H}$ vector space. Suppose $\tilde{\Gamma}$ is a symmetric real-valued bilinear form on V_R which factors through $V_R \otimes_{\mathbb{H}} \overline{V_R}$ and h a symmetric real-valued bilinear form on V_L which factors through $\overline{V_L} \otimes_{\mathbb{H}} V_L$. Then we claim that there is a well-defined symmetric real bilinear form B on the real vector space $V = V_R \otimes_{\mathbb{H}} V_L$ constructed from $\tilde{\Gamma}, h$. In fact, if $\tilde{\Gamma}_1, h_1$ are the $\mathbb{H}$-valued pairings constructed as in (3.3), then

$$B(f \otimes v, f' \otimes v') = \mathrm{Re}\Big\{ \tilde{\Gamma}_1(f, \bar{f}')\, h_1(\bar{v}, v) \Big\}, \qquad f, f' \in V_R, \quad v, v' \in V_L. \tag{3.12}$$

We easily check that this factors through the quaternionic tensor product using (3.2). Relative to bases $\{f^a\}$ of V_R and v_i of V_L, any vector in V can be written

$$f^a \psi^i_a \otimes v_i = f^a \otimes \psi^i_a v_i, \qquad \psi^i_a \in \mathbb{H}. \tag{3.13}$$

Then the formula for B is

$$B(f^b \psi^j_b v_j, f^a \lambda^i_a v^a) = \mathrm{Re}(\overline{\psi^j_b} \tilde{\Gamma}^{ab} \lambda^i_a h_{ij}). \tag{3.14}$$

§3.2. The free supersymmetric σ-model

The bosonic σ-model lagrangian in n-dimensional Minkowski space $\check{M}^n$ for a scalar field $\phi\colon \check{M}^n \to X$ with values in a Riemannian manifold X is

$$L = \frac{1}{2} |d\phi|^2 \, |d^n x|. \tag{3.15}$$

We look for supersymmetric extensions with various amounts of supersymmetry. With a single supersymmetry the supersymmetric extension for $n = 1$ is given in (1.53). (That lagrangian is in $\check{M}^1$, but in fact it is the dimensional reduction of a lagrangian in $\check{M}^2$; see Problem FP6 of [I-**Homework**].) To construct models with more supersymmetry we first consider the linear case $X = \mathbb{R}^k$ for some k; in the next subsection we treat nonlinear σ-models. The linear model is free, and

to predict which supersymmetric extensions exist we invoke an argument from the quantum theory. Namely, upon quantization we obtain a representation of the supersymmetry algebra, and the representation theory shows that models are possible with $s = 2, 4, 8$ supersymmetries for $k = 1, 2, 4$. That is, the smallest model has a scalar field in $\mathbb{R}^k$ for $k = 1, 2, 4$ and a single spinor field. The maximal dimension in which such models can occur is $n = 3, 4, 6$, and the models in these dimensions have minimal supersymmetry.

We refer to the theories in $n = 3, 4, 6$ according to the field $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$.

To treat the various cases simultaneously, we introduce constants κ in (3.6) and

$$A = \begin{cases} \frac{1}{2}, & \mathbb{F} = \mathbb{R}; \\ 1, & \mathbb{F} = \mathbb{C}, \mathbb{H}. \end{cases} \tag{3.16}$$

(The constant A enters the lagrangian (3.19) and the constant κ enters the supersymmetry transformation laws (3.23), as well as (3.5) above.)

The fields in the theory are

$$\begin{aligned} \phi\colon \check{M}^n &\longrightarrow \mathbb{F} \\ \psi\colon \check{M}^n &\longrightarrow \Pi S, \end{aligned} \tag{3.17}$$

where 'Π' denotes parity reversal. (The spinor field ψ is odd.)

Theorem 3.18. *The σ-model lagrangian*

$$\boxed{L = A\Big\{\langle d\phi, d\bar{\phi}\rangle + \mathrm{Re}(\psi \not{D} \bar{\psi})\Big\}\,|d^n x|} \tag{3.19}$$

is supersymmetric in the three cases $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$.

Proof. The lagrangian (3.19) in more detail is[27]

$$\boxed{L = A\Big\{g^{\mu\nu}\partial_\mu\phi\cdot\partial_\nu\bar{\phi} + \frac{1}{2}\psi_a\cdot\tilde{\Gamma}^{\mu\dot{b}a}\cdot\partial_\mu\bar{\psi}_{\dot{b}} - \frac{1}{2}\partial_\mu\psi_a\cdot\tilde{\Gamma}^{\mu\dot{b}a}\cdot\bar{\psi}_{\dot{b}}\Big\}\,|d^n x|,} \tag{3.21}$$

where '$\cdot$' denotes multiplication in $\mathbb{F}$. We usually omit '$\cdot$'. In case $\mathbb{F} = \mathbb{R}, \mathbb{C}$ we can combine the two fermion terms at the cost of an exact term. For odd parameters $\eta^a \in \mathbb{F}$ we let $\hat{\xi}$ denote the *real* vector field on the space of fields which corresponds to the abstract supersymmetry transformation

$$\frac{\kappa^2}{2}(\eta^a Q_a - \overline{Q}_{\dot{a}}\bar{\eta}^{\dot{a}}), \tag{3.22}$$

where $\bar{\eta}^{\dot{a}} = \overline{\eta^a}$. (Recall the abstract supersymmetry algebra (1.16).) We postulate

$$\boxed{\begin{aligned} \hat{\xi}\phi &= \kappa\psi_a\,\eta^a \\ \hat{\xi}\psi_a &= \kappa\partial_\mu\phi\,\bar{\eta}^{\dot{b}}\,\Gamma^\mu_{a\dot{b}}, \end{aligned}} \tag{3.23}$$

[27] The fermion kinetic term is A times

$$\tilde{\Gamma}(f^a\psi_a, f^b\partial_\mu\psi_b) = \tilde{\Gamma}(f^b\partial_\mu\psi_b, f^a\psi_a) = \mathrm{Re}(\bar{\psi}_{\dot{a}}\tilde{\Gamma}^{\mu\dot{a}b}\partial_\mu\psi_b) = \mathrm{Re}(\psi_a\tilde{\Gamma}^{\mu\dot{b}a}\partial_\mu\bar{\psi}_{\dot{b}}). \tag{3.20}$$

and of course we have the conjugate equations

$$\begin{aligned}\hat{\xi}\bar{\phi} &= -\kappa\bar{\eta}^{\dot{a}}\,\bar{\psi}_{\dot{a}}\\ \hat{\xi}\bar{\psi}_{\dot{a}} &= \quad \kappa\Gamma^{\mu}_{b\dot{a}}\,\eta^{b}\,\partial_{\mu}\bar{\phi}\,.\end{aligned} \tag{3.24}$$

There is a minus sign in passing from $\hat{\xi}\phi$ to $\hat{\xi}\bar{\phi}$ since we commute the odd quantities η^a and ψ_a. These formulas are motivated by Lorentz invariance, parity, and power counting.

We must check that $\mathrm{Lie}(\hat{\xi})L$ is exact and that the transformations (3.23) generate the supersymmetry algebra on-shell.[28] The variation of the bosonic term in L is

$$\hat{\xi}(Ag^{\mu\nu}\,\partial_{\mu}\phi\cdot\partial_{\nu}\bar{\phi}) = \kappa A g^{\mu\nu}\{\partial_{\mu}\psi_{a}\,\eta^{a}\,\partial_{\nu}\bar{\phi} - \partial_{\mu}\phi\,\bar{\eta}^{\dot{a}}\,\partial_{\nu}\bar{\psi}_{\dot{a}}\}. \tag{3.25}$$

The second term in (3.21) has variation

(3.26)

$$\begin{aligned}\hat{\xi}(\frac{A}{2}\,\psi_{a}\cdot\tilde{\Gamma}^{\mu\dot{b}a}\cdot\partial_{\mu}\bar{\psi}_{\dot{b}}) &= \frac{\kappa A}{2}\{\partial_{\nu}\phi\,\bar{\eta}^{\dot{c}}\,\Gamma^{\nu}_{a\dot{c}}\tilde{\Gamma}^{\mu\dot{b}a}\,\partial_{\mu}\bar{\psi}_{\dot{b}} + \psi_{a}\tilde{\Gamma}^{\mu\dot{b}a}\Gamma^{\nu}_{c\dot{b}}\eta^{c}\partial_{\mu}\partial_{\nu}\bar{\phi}\}\\ &= \frac{\kappa A}{2}\{\partial_{\nu}\phi\,\bar{\eta}^{\dot{c}}\,\Gamma^{\nu}_{a\dot{c}}\tilde{\Gamma}^{\mu\dot{b}a}\,\partial_{\mu}\bar{\psi}_{\dot{b}} - \partial_{\nu}\psi_{a}\,\tilde{\Gamma}^{\mu\dot{b}a}\Gamma^{\nu}_{c\dot{b}}\,\eta^{c}\,\partial_{\mu}\bar{\phi}\}\\ &\qquad + \partial_{\nu}\Big\{\frac{\kappa A}{2}\,\psi_{a}\,\tilde{\Gamma}^{\mu\dot{b}a}\Gamma^{\nu}_{c\dot{b}}\,\eta^{c}\,\partial_{\mu}\bar{\phi}\Big\}.\end{aligned}$$

The variation of the third term in (3.21) is the conjugate

(3.27)

$$\begin{aligned}\hat{\xi}(-\frac{A}{2}\,\partial_{\mu}\psi_{a}\cdot\tilde{\Gamma}^{\mu\dot{b}a}\cdot\bar{\psi}_{\dot{b}}) = \frac{\kappa A}{2}\{&-\partial_{\mu}\psi_{a}\,\tilde{\Gamma}^{\mu\dot{b}a}\Gamma^{\nu}_{c\dot{b}}\,\eta^{c}\,\partial_{\nu}\bar{\phi} + \partial_{\mu}\phi\,\bar{\eta}^{\dot{c}}\,\Gamma^{\nu}_{a\dot{c}}\tilde{\Gamma}^{\mu\dot{b}a}\,\partial_{\nu}\bar{\psi}_{\dot{b}}\}\\ &- \partial_{\nu}\{\frac{\kappa A}{2}\partial_{\mu}\phi\,\bar{\eta}^{\dot{c}}\,\Gamma^{\nu}_{a\dot{c}}\tilde{\Gamma}^{\mu\dot{b}a}\bar{\psi}_{\dot{b}}\}.\end{aligned}$$

Adding (3.26) and (3.27) and using the Clifford relation (3.8) we see

$$\hat{\xi}(\text{fermion term}) = \frac{\kappa A}{2}g^{\mu\nu}\{\partial_{\mu}\phi\,\bar{\eta}^{\dot{a}}\,\partial_{\nu}\bar{\psi}_{\dot{a}} - \partial_{\mu}\psi_{a}\,\eta^{a}\,\partial_{\nu}\bar{\phi}\} + EXACT, \tag{3.28}$$

where

$$EXACT = \partial_{\nu}\{\frac{\kappa A}{2}\psi_{a}\,\tilde{\Gamma}^{\mu\dot{b}a}\Gamma^{\nu}_{c\dot{b}}\,\eta^{c}\,\partial_{\mu}\bar{\phi}\ -\ \frac{\kappa A}{2}\partial_{\mu}\phi\,\bar{\eta}^{\dot{c}}\,\Gamma^{\nu}_{a\dot{c}}\tilde{\Gamma}^{\mu\dot{b}a}\,\bar{\psi}_{\dot{b}}\}. \tag{3.29}$$

The nonexact term in (3.28) cancels against (3.25), which proves that $\hat{\xi}$ is a nonmanifest symmetry of L.

We now verify that the supersymmetry algebra holds on-shell. The vector field (3.22) is $\kappa^2\,\mathrm{Re}(\eta^a Q_a)$, and in the abstract supersymmetry algebra (1.16) we have, according to (3.5),

$$\begin{aligned}\big[\mathrm{Re}(\eta_1^a Q_a), \mathrm{Re}(\eta_2^b Q_b)\big] &= -\big[\mathrm{Re}(\eta_2^a Q_a), \mathrm{Re}(\eta_1^b Q_b)\big]\\ &= 2\,\Gamma\big(\mathrm{Re}(\eta_2^a Q_a), \mathrm{Re}(\eta_1^b Q_b)\big)\\ &= \frac{2}{\kappa^2}\,\mathrm{Re}(\eta_1^b\Gamma^{\mu}_{b\dot{a}}\bar{\eta}_2^{\dot{a}})P_{\mu}\\ &= \frac{2}{\kappa^2}\,\mathrm{Re}(\bar{\eta}_1^{\dot{b}}\Gamma^{\mu}_{a\dot{b}}\eta_2^{a})P_{\mu}.\end{aligned} \tag{3.30}$$

[28] By contrast, in one dimension (1.47) holds everywhere, that is, off-shell. We can achieve off-shell supersymmetry for $\mathbb{F} = \mathbb{R}, \mathbb{C}$ by introducing an auxiliary field; see §4.1 and §5.1.

So on a field f we must verify (see (1.30))

$$[\hat{\xi}_1, \hat{\xi}_2]f = 2\kappa^2 \,\mathrm{Re}(\bar{\eta}_1^{\dot{b}}\Gamma^{\mu}_{a\dot{b}}\eta_2^a)\,\partial_\mu f. \tag{3.31}$$

The check for ϕ is straightforward:

$$\hat{\xi}_1\hat{\xi}_2\phi = \hat{\xi}_1(\kappa\psi_a\eta_2^a) = \kappa^2\partial_\mu\phi\,\bar{\eta}_1^{\dot{b}}\Gamma^{\mu}_{a\dot{b}}\eta_2^a, \tag{3.32}$$

and so

$$\begin{aligned}[\hat{\xi}_1, \hat{\xi}_2]\phi &= \kappa^2\partial_\mu\phi(\bar{\eta}_1^{\dot{b}}\Gamma^{\mu}_{a\dot{b}}\eta_2^a - \bar{\eta}_2^{\dot{b}}\Gamma^{\mu}_{a\dot{b}}\eta_1^a)\\ &= 2\kappa^2\,\mathrm{Re}(\bar{\eta}_1^{\dot{b}}\Gamma^{\mu}_{a\dot{b}}\eta_2^a)\,\partial_\mu\phi,\end{aligned} \tag{3.33}$$

as desired. For ψ we must use the equation of motion (3.39), which appears below. Using (3.10) we write this equation as

$$\begin{aligned}\partial_0\psi_1 - \partial_1\psi_1 &= \partial_2\psi_2 - \partial_3\psi_2 i + \partial_4\psi_2 j + \partial_5\psi_2 k,\\ \partial_0\psi_2 + \partial_1\psi_2 &= \partial_2\psi_1 + \partial_3\psi_1 i - \partial_4\psi_1 j - \partial_5\psi_1 k.\end{aligned} \tag{3.34}$$

The action of the supersymmetry transformation on ψ is

$$\hat{\xi}_1\hat{\xi}_2\psi_a = \hat{\xi}_1(\kappa\partial_\mu\phi\,\bar{\eta}_2^{\dot{b}}\Gamma^{\mu}_{a\dot{b}}) = \kappa^2\partial_\mu\psi_c\,\eta_1^c\bar{\eta}_2^{\dot{b}}\Gamma^{\mu}_{a\dot{b}}. \tag{3.35}$$

So

$$[\hat{\xi}_1, \hat{\xi}_2]\psi_a = \kappa^2\partial_\mu\psi_c(\eta_1^c\bar{\eta}_2^{\dot{b}} - \eta_2^c\bar{\eta}_1^{\dot{b}})\Gamma^{\mu}_{a\dot{b}}. \tag{3.36}$$

Checking explicitly for $a = 1, 2$ using (3.9) and (3.34) we find the desired result

$$[\hat{\xi}_1, \hat{\xi}_2]\psi_a = 2\kappa^2\,\mathrm{Re}(\bar{\eta}_1^{\dot{b}}\Gamma^{\mu}_{c\dot{b}}\eta_2^c)\,\partial_\mu\psi_a. \tag{3.37}$$

This completes the proof.

We compute the equations of motion and the variational 1-form. Write the lagrangian (3.21) as $L = \check{L}\,|d^n x|$. Then

$$\begin{aligned}\frac{1}{A}\delta\check{L} = {}&\delta\phi(-g^{\mu\nu}\partial_\mu\partial_\nu\bar{\phi}) + (-g^{\mu\nu}\partial_\mu\partial_\nu\phi)\,\delta\bar{\phi} + \delta\psi_a(-\tilde{\Gamma}^{\mu\dot{b}a}\partial_\mu\bar{\psi}_{\dot{b}}) + (\partial_\mu\psi_a\,\tilde{\Gamma}^{\mu\dot{b}a})\,\delta\bar{\psi}_{\dot{b}}\\ &+ \partial_\nu\{g^{\mu\nu}\delta\phi\,\partial_\mu\bar{\phi} + g^{\mu\nu}\partial_\mu\phi\,\delta\bar{\phi} + \frac{1}{2}\delta\psi_a\,\tilde{\Gamma}^{\nu\dot{b}a}\,\bar{\psi}_{\dot{b}} - \frac{1}{2}\psi_a\,\tilde{\Gamma}^{\nu\dot{b}a}\,\delta\bar{\psi}_{\dot{b}}\}.\end{aligned} \tag{3.38}$$

This leads to the equations of motion

$$\boxed{\begin{aligned}g^{\mu\nu}\partial_\mu\partial_\nu\phi &= 0\\ \partial_\mu\psi_a\tilde{\Gamma}^{\mu\dot{b}a} &= 0\end{aligned}} \tag{3.39}$$

and the variational 1-form

(3.40)
$$\boxed{\gamma = A(g^{\mu\nu}\delta\phi\,\partial_\mu\bar\phi + g^{\mu\nu}\partial_\mu\phi\,\delta\bar\phi + \frac{1}{2}\delta\psi_a\,\tilde\Gamma^{\nu\dot b a}\,\bar\psi_{\dot b} - \frac{1}{2}\psi_a\,\tilde\Gamma^{\nu\dot b a}\,\delta\bar\psi_{\dot b})\,\frac{\partial}{\partial x^\nu}\otimes|d^n x|.}$$

In coordinate-free notation the equations of motion are

(3.41)
$$\boxed{\begin{aligned}\Delta\phi &= 0\\ \not{D}\psi &= 0.\end{aligned}}$$

The Noether current associated to $\hat\xi$ is computed from (3.40) and (3.29) to be

(3.42)
$$\begin{aligned}\iota(\hat\xi)\gamma - (\frac{\kappa A}{2}\psi_a\,\tilde\Gamma^{\mu\dot b a}\Gamma^\nu_{c\dot b}\,\eta^c\,\partial_\mu\bar\phi - \frac{\kappa A}{2}\partial_\mu\phi\,\bar\eta^{\dot c}\,\Gamma^\nu_{a\dot c}\tilde\Gamma^{\mu\dot b a}\,\bar\psi_{\dot b}\,)\,\frac{\partial}{\partial x^\nu}\otimes|d^n x|\\ = \kappa A(\psi_a\,\tilde\Gamma^{\nu\dot b a}\Gamma^\mu_{c\dot b}\,\eta^c\,\partial_\mu\bar\phi - \partial_\mu\phi\,\bar\eta^{\dot c}\,\Gamma^\mu_{a\dot c}\tilde\Gamma^{\nu\dot b a}\bar\psi_{\dot b})\,\frac{\partial}{\partial x^\nu}\otimes|d^n x|.\end{aligned}$$

(We use the Clifford identity (3.8) in this computation.) Thus for $\mathbb{F} = \mathbb{R}, \mathbb{C}$ the *supercurrent*, which is *minus* the Noether current for Q_a, is

(3.43)
$$\boxed{j_a = \frac{2A}{\kappa}\psi_b\,\tilde\Gamma^{\nu\dot c b}\Gamma^\mu_{a\dot c}\,\partial_\mu\bar\phi\,\frac{\partial}{\partial x^\nu}\otimes|d^n x|, \qquad \mathbb{F} = \mathbb{R}, \mathbb{C}.}$$

(Recall the factor $\kappa^2/2$ in (3.22).) As a check, in the case $\mathbb{F} = \mathbb{R}$ this reduces in one dimension with one supersymmetry to (1.57). Other formulas from that section also check against those here.

§3.3. Nonlinear supersymmetric σ-model

Now consider the supersymmetric σ-model of the previous subsection where we replace the target X by a curved Riemannian manifold. Then ψ transforms as an element of the pullback tangent bundle, and it is easy to see that $\dim X$ is divisible by $\dim \mathbb{F}$ if the linearization is to reduce to the linear σ-model. But more is true: To write formulas (3.23) we need to be able to multiply tangent vectors in X by elements of $\mathbb{F}$. Thus we assume that TX is endowed with a *left* $\mathbb{F}$-structure. Of course, this is no extra condition for $\mathbb{F} = \mathbb{R}$. For $\mathbb{F} = \mathbb{C}, \mathbb{H}$ the $\mathbb{F}$ multiplication is a tensor field on X (which is called an almost complex structure for $\mathbb{F} = \mathbb{C}$). Let T denote the covariant derivative of this tensor. Then X is *Kähler* ($\mathbb{F} = \mathbb{C}$) or *hyperkähler* ($\mathbb{F} = \mathbb{H}$) if and only if T vanishes. The fields in the theory are now

(3.44)
$$\begin{aligned}\phi &\in C^\infty(\check M^n, X)\\ \psi &\in C^\infty(\check M^n, \Pi S \otimes_{\mathbb{F}} \phi^* TX).\end{aligned}$$

The lagrangian (3.19) continues to make sense if we use the Riemannian metric h on TX as well as the Levi-Civita covariant derivative ∇. But for ∇ to be well-defined on TX as an $\mathbb{F}$-bundle, we need T to vanish. Therefore, we now assume that X is Kähler for $\mathbb{F} = \mathbb{C}$ and that X is hyperkähler for $\mathbb{F} = \mathbb{H}$. For $\mathbb{F} = \mathbb{H}$ we explained above that if $\{v_i\}$ is a basis of TX at some point, then the Riemannian

metric is represented by a quaternionic matrix (h_{ij}). Using (3.14) we write the lagrangian (3.21) as

$$L = \mathrm{Re}\Big\{ g^{\mu\nu}\partial_\mu\phi^j \cdot h_{ij} \cdot \partial_\nu\bar\phi^i + \bar\psi^j_{\dot b}\cdot\tilde\Gamma^{\mu\dot b a}\cdot\nabla_\mu\psi^i_a\cdot h_{ij}\Big\}\,|d^6x|. \tag{3.45}$$

In (3.57) below we abbreviate the second term as '$\langle\bar\psi_{\dot b}\tilde\Gamma^{\mu\dot b a}\nabla_\mu\psi_a\rangle$'. In all cases the lagrangian (3.19) ceases to be supersymmetric if X is not flat. Rather, we have the following result. Denote the metric h on X as '$\langle\cdot,\cdot\rangle$' and let R be the Riemann curvature tensor.

Theorem 3.46. (*i*) ($\mathbb{F}=\mathbb{R}$) *The nonlinear σ-model lagrangian*

$$\boxed{L = \Big\{\frac{1}{2}|d\phi|^2 + \frac{1}{2}\langle\psi\not{D}_\phi\psi\rangle + \frac{1}{12}\epsilon^{ab}\epsilon^{cd}\langle\psi_a,\phi^*R(\psi_b,\psi_c)\psi_d\rangle\Big\}\,|d^3x|} \tag{3.47}$$

is supersymmetric for any Riemannian manifold X.

(*ii*) ($\mathbb{F}=\mathbb{C}$) *The nonlinear σ-model lagrangian*

$$\boxed{L = \Big\{\langle\overline{d\phi},d\phi\rangle + \mathrm{Re}\langle\bar\psi\not{D}_\phi\psi\rangle - \frac{1}{4}\epsilon^{ac}\epsilon^{\dot b\dot d}\langle\psi_a,\phi^*R(\psi_c,\bar\psi_{\dot d})\bar\psi_{\dot b}\rangle\Big\}\,|d^4x|} \tag{3.48}$$

is supersymmetric if and only if X is Kähler.

(*iii*) ($\mathbb{F}=\mathbb{H}$) *The nonlinear σ-model lagrangian*

$$\boxed{\begin{aligned} L = \Big\{&\frac{1}{16}\tilde h_{\alpha\beta}\epsilon_{ij}\nu^{abcd}(\partial_{ab}\phi)^{\alpha i}(\partial_{cd}\phi)^{\beta j} + \frac{1}{4}\tilde h_{\alpha\beta}\nu^{abcd}\psi^\alpha_a(\phi^*\nabla_{cd})\psi^\beta_b \\ &- \frac{1}{24}\nu^{abcd}\phi^*\Omega_{\alpha\beta\gamma\delta}\psi^\alpha_a\psi^\beta_b\psi^\gamma_c\psi^\delta_d\Big\}\,|d^6x|. \end{aligned}} \tag{3.49}$$

is supersymmetric if and only if X is hyperkähler.

In all cases the Dirac form involves the covariant derivative ∇, and so it also depends on ϕ. For example, in the case $\mathbb{F}=\mathbb{R}$ we have

$$\frac{1}{2}\langle\psi\not{D}_\phi\psi\rangle = \frac{1}{2}\tilde\Gamma^{\mu ab}\langle\psi_a,(\phi^*\nabla_\mu)\psi_b\rangle. \tag{3.50}$$

Note the use of angle brackets on the left hand side to denote the inner product on TX. The supersymmetry transformation laws are the same as for the linear case.[29] We often omit 'ϕ' and 'ϕ^*' from the notation.

We explain (3.49), where we have switched from the quaternionic notation used previously to a complex notation. The complexification of the tangent bundle of a hyperkähler manifold X of real dimension $4n$ can be written

$$T_{\mathbb{C}}X \cong V\otimes W^*, \tag{3.51}$$

where V is a complex vector bundle of complex rank $2n$ and W^* is the dual of the fixed complex symplectic vector space described at the beginning of §2.1. Furthermore, V carries a pseudoreal (quaternionic) structure J and a skew-symmetric

[29] Physicists often use a spinor field ψ which is not intrinsic, so have different formulas.

bilinear form $\tilde{h}$ which encodes the metric. The complexified Riemann curvature tensor, viewed as a functional on $(T_{\mathbb{C}}X)^{\otimes 4} \cong V^{\otimes 4}$, is a totally symmetric tensor Ω. Let S be the real spin representation of $\mathrm{Spin}(1,5)$. As described in §2.1, its complexification is $S_0 \otimes W$ for a 4-dimensional complex vector space S_0. The spinor field ψ is a real odd section of $\phi^*V \otimes_{\mathbb{C}} S_0$. Note there is a trace map

$$\phi^*(T_{\mathbb{C}}X) \otimes S_{\mathbb{C}} \cong (\phi^*V \otimes W^*) \otimes (S_0 \otimes W) \longrightarrow \phi^*V \otimes S_0 \tag{3.52}$$

which expresses the quaternionic tensor product in (3.44). (All tensor products in (3.52) are over $\mathbb{C}$.) The pseudoreal structures on S_0 and ϕ^*V give a real structure on $\phi^*V \otimes S_0$. Recall that S_0 carries a volume form ν. Now fix a local framing of V (indices $\alpha, \beta, \dots = 1, \dots, 2n$), a basis of W (indices $i, j, \dots = 1, 2$), and a basis of S_0 (indices $a, b, \dots = 1, 2, 3, 4$). The inner product of two tangent vectors ξ_1, ξ_2 is

$$\langle \xi_1, \xi_2 \rangle = \tilde{h}_{\alpha\beta}\epsilon_{ij}\xi_1^{\alpha i}\xi_2^{\beta j}. \tag{3.53}$$

We will find it useful to lift a spinor field ψ to an odd section of $\phi^*(T_{\mathbb{C}}X) \otimes S_{\mathbb{C}}$ via (3.52):

$$\psi_{aj}^{\alpha i} = \psi_a^\alpha \delta_j^i. \tag{3.54}$$

Our normalization of the tensor Ω in terms of the Riemann curvature R is

$$R_{(\alpha i)(\beta j)(\gamma k)(\delta \ell)} = \epsilon_{ij}\epsilon_{k\ell}\Omega_{\alpha\beta\gamma\delta}. \tag{3.55}$$

In this notation the supersymmetry variation of the fields (3.23) is

$$\boxed{\begin{aligned} (\hat{\xi}\phi)^{\alpha i} &= -\sqrt{2}\eta^{ai}\psi_a^\alpha \\ \hat{\xi}\psi_a^\alpha &= \sqrt{2}\,\epsilon_{ij}\eta^{bj}(\partial_{ab}\phi)^{\alpha i} \end{aligned}} \tag{3.56}$$

for $\eta^{ai} \in \mathbb{C}$.

Proof of Theorem 3.46. We review the proof of Theorem 3.18 to determine what changes when X is curved. (In this paragraph only we use quaternionic notation for the $\mathbb{F} = \mathbb{H}$ case.) The variation of the lagrangian L has a new contribution from varying the covariant derivative in the fermion kinetic term, and this cancels the variation of the curvature term. We do this computation presently, but first we examine changes in the previous computations. For $\mathbb{F} = \mathbb{R}$ they are unaltered except for the substitution of covariant derivatives. But for $\mathbb{F} = \mathbb{C}, \mathbb{H}$ our assumption that X is Kähler, hyperkähler is crucial. For in the "integration by parts" formula (3.26) there is an extra term from differentiating the $\mathbb{F}$-structure:

(3.57)

$$\begin{aligned} \hat{\xi}_\psi\big(\frac{A}{2}\langle \psi_a \cdot \tilde{\Gamma}^{\mu\dot{b}a} \cdot \nabla_\mu \bar{\psi}_{\dot{b}} \rangle\big) = \frac{\kappa A}{2}\big\{\langle \partial_\nu \phi\, \bar{\eta}^{\dot{c}}\, \Gamma^\nu_{a\dot{c}} \tilde{\Gamma}^{\mu\dot{b}a}\, \nabla_\mu \bar{\psi}_{\dot{b}} \rangle - \langle \partial_\nu \psi_a\, \tilde{\Gamma}^{\mu\dot{b}a} \Gamma^\nu_{c\dot{b}}\, \eta^c\, \partial_\mu \bar{\phi} \rangle\big\} \\ + \partial_\nu\big\{\frac{\kappa A}{2}\langle \psi_a\, \tilde{\Gamma}^{\mu\dot{b}a} \Gamma^\nu_{c\dot{b}}\, \eta^c\, \partial_\mu \bar{\phi} \rangle\big\} \\ - \frac{\kappa A}{2}\big\{\langle \psi_a\, T_\nu\, \tilde{\Gamma}^{\mu\dot{b}a} \Gamma^\nu_{c\dot{b}}\, \eta^c\, \partial_\mu \bar{\phi} \rangle + \langle \psi_a\, \tilde{\Gamma}^{\mu\dot{b}a} \Gamma^\nu_{c\dot{b}}\, \eta^c\, T_\nu\, \partial_\mu \bar{\phi} \rangle\big\}. \end{aligned}$$

The subscript on $\hat{\xi}$ indicates that we only vary the fermion; the variation of ϕ is computed below in (3.60). We also pick up the conjugate terms in (3.27). Our assumption on X is that $T = 0$, so that this extra term does not appear. With this assumption the verification of the supersymmetry algebra also goes through as before.

We turn to the variation of the covariant derivative and the curvature term. Here it is easier to proceed on a case-by-case basis. First, consider $\mathbb{F} = \mathbb{R}$ so that X is any Riemannian manifold. Let R denote the Riemann curvature tensor. Fix a basis $\{v_i\}$ of TX at a point. Then as usual we write

$$R_{ijk\ell} = \langle v_i, R(v_k, v_\ell)v_j\rangle. \tag{3.58}$$

The curvature tensor obeys certain symmetries and satisfies the Bianchi identity:

$$\begin{aligned} &R_{ijk\ell} = -R_{jik\ell} = -R_{ij\ell k} = R_{k\ell ij} \\ &R_{ijk\ell} + R_{jki\ell} + R_{kij\ell} = 0. \end{aligned} \tag{3.59}$$

Now the variation in ϕ of (3.50) is

$$\begin{aligned} \hat{\xi}_\phi\big(\frac{1}{2}\tilde{\Gamma}^{\mu ab}\langle\psi_a, (\phi^*\nabla_\mu)\psi_b\rangle\big) &= \frac{1}{2}\tilde{\Gamma}^{\mu ab}\langle\psi_a, R(\hat{\xi}\phi, \partial_\mu\phi)\psi_b\rangle \\ &= \frac{1}{2}\tilde{\Gamma}^{\mu ab}\eta^c\langle\psi_a, R(\psi_c, \partial_\mu\phi)\psi_b\rangle \\ &= -\frac{1}{2}\tilde{\Gamma}^{\mu ab}\eta^c R_{ijk\ell}\psi_a^i\psi_b^j\psi_c^k(\partial_\mu\phi)^\ell. \end{aligned} \tag{3.60}$$

The last minus sign comes from commuting ψ_b and ψ_c. We claim that this cancels against the variation in ψ of the curvature term in the lagrangian (3.47). That variation is

(3.61)

$$\begin{aligned} &\hat{\xi}_\psi\big(\frac{1}{12}\epsilon^{ab}\epsilon^{cd}R_{ijk\ell}\psi_a^i\psi_d^j\psi_b^k\psi_c^\ell\big) \\ &\quad= \frac{1}{12}\epsilon^{ab}\epsilon^{cd}R_{ijk\ell}\eta^e\Big[\Gamma^\mu_{ae}(\partial_\mu\phi)^i\psi_d^j\psi_b^k\psi_c^\ell - \Gamma^\mu_{de}\psi_a^i(\partial_\mu\phi)^j\psi_b^k\psi_c^\ell \\ &\qquad\qquad + \Gamma^\mu_{be}\psi_a^i\psi_d^j(\partial_\mu\phi)^k\psi_c^\ell - \Gamma^\mu_{ce}\psi_a^i\psi_d^j\psi_b^k(\partial_\mu\phi)^\ell\Big] \\ &\quad= \frac{1}{12}\eta^e\Gamma^\mu_{ae}\Big[\epsilon^{ab}\epsilon^{cd}R_{ijk\ell} - \epsilon^{db}\epsilon^{ca}R_{jik\ell} + \epsilon^{da}\epsilon^{cb}R_{jki\ell} - \epsilon^{dc}\epsilon^{ab}R_{jk\ell i}\Big](\partial_\mu\phi)^i\psi_d^j\psi_b^k\psi_c^\ell \\ &\quad= \frac{1}{12}\eta^e\Gamma^\mu_{ae}\epsilon^{ab}\epsilon^{cd}\Big[R_{ijk\ell} + R_{ijk\ell} + R_{kji\ell} + R_{kji\ell}\Big](\partial_\mu\phi)^i\psi_d^j\psi_b^k\psi_c^\ell \\ &\quad= \frac{1}{6}\eta^e\Gamma^\mu_{ae}\epsilon^{ab}\epsilon^{cd}(R_{ijk\ell} + R_{kji\ell})(\partial_\mu\phi)^i\psi_d^j\psi_b^k\psi_c^\ell \\ &\quad= \frac{1}{6}\Gamma^\mu_{ae}\epsilon^{ab}(-\eta^d\epsilon^{ec} - \eta^c\epsilon^{de})(R_{ijk\ell} + R_{kji\ell})(\partial_\mu\phi)^i\psi_d^j\psi_b^k\psi_c^\ell \\ &\quad= \frac{1}{6}(-\tilde{\Gamma}^{\mu bc}\eta^d + \tilde{\Gamma}^{\mu bd}\eta^c)(R_{ijk\ell} + R_{kji\ell})(\partial_\mu\phi)^i\psi_d^j\psi_b^k\psi_c^\ell \\ &\quad= \frac{1}{3}\tilde{\Gamma}^{\mu bd}\eta^c(R_{ijk\ell} + R_{kji\ell})(\partial_\mu\phi)^i\psi_d^j\psi_b^k\psi_c^\ell. \end{aligned}$$

In the third to last step we use the fact that $\mathbb{F}^2$ is two-dimensional, so that the cyclic sum $\eta^e\epsilon^{cd} + \eta^d\epsilon^{ec} + \eta^c\epsilon^{de}$ vanishes. To compare (3.61) to (3.60) we need to move around indices:

$$\hat{\xi}_\psi\big(\frac{1}{12}\epsilon^{ab}\epsilon^{cd}R_{ijk\ell}\psi_a^i\psi_d^j\psi_b^k\psi_c^\ell\big) = \frac{1}{3}\tilde{\Gamma}^{\mu ab}\eta^c(R_{\ell ijk} + R_{ijk\ell})\psi_a^i\psi_b^j\psi_c^k(\partial_\mu\phi)^\ell. \tag{3.62}$$

Using the Bianchi identity (3.59) and other symmetries we find

$$\begin{aligned}\tilde{\Gamma}^{\mu ab}\eta^c R_{\ell ijk}\psi_a^i\psi_b^j\psi_c^k(\partial_\mu\phi)^\ell &= -\tilde{\Gamma}^{\mu ab}\eta^c(R_{ij\ell k} + R_{j\ell ik})\psi_a^i\psi_b^j\psi_c^k(\partial_\mu\phi)^\ell \\ &= \ \ \tilde{\Gamma}^{\mu ab}\eta^c(R_{ijk\ell} + R_{i\ell jk})\psi_a^i\psi_b^j\psi_c^k(\partial_\mu\phi)^\ell\end{aligned} \tag{3.63}$$

from which

$$\tilde{\Gamma}^{\mu ab}\eta^c R_{\ell ijk}\psi_a^i\psi_b^j\psi_c^k(\partial_\mu\phi)^\ell = \frac{1}{2}\tilde{\Gamma}^{\mu ab}\eta^c R_{ijk\ell}\psi_a^i\psi_b^j\psi_c^k(\partial_\mu\phi)^\ell. \tag{3.64}$$

Substituting into (3.62) we find

$$\hat{\xi}_\psi\big(\frac{1}{12}\epsilon^{ab}\epsilon^{cd}R_{ijk\ell}\psi_a^i\psi_d^j\psi_b^k\psi_c^\ell\big) = \frac{1}{2}\tilde{\Gamma}^{\mu ab}\eta^c R_{ijk\ell}\psi_a^i\psi_b^j\psi_c^k(\partial_\mu\phi)^\ell \tag{3.65}$$

which cancels (3.60).

To complete the verification that (3.47) is supersymmetric, we check that the variation in ϕ of the curvature term vanishes:

$$\hat{\xi}_\phi\big(\epsilon^{ab}\epsilon^{cd}\langle\psi_a, R(\psi_b,\psi_c)\psi_d\rangle\big) = \epsilon^{ab}\epsilon^{cd}\eta^e\langle\psi_a, (\nabla_{\psi_e}R)(\psi_b,\psi_c)\psi_d\rangle. \tag{3.66}$$

Now the symmetries of R and the (second) Bianchi identity imply that the expression $\langle\psi_a, (\nabla_{\psi_e}R)(\psi_b,\psi_c)\psi_d\rangle$ is symmetric in a,d and in b,c,e. In addition, it is symmetric under the interchange $a,d \leftrightarrow b,c$. Altogether this implies that it is totally symmetric in a,b,c,d,e, whence (3.66) vanishes. This completes the proof for $\mathbb{F} = \mathbb{R}$.

Next, consider $\mathbb{F} = \mathbb{C}$. Thus X is Kähler and we work with the complexified tangent bundle as usual. We consider ψ_a to be a complex vector of type $(1,0)$. The Riemann curvature tensor is of type $(1,1)$. If $\{v_i\}$ is a basis of type $(1,0)$ vectors at a point of X, we write

$$R_{i\bar{j}k\bar{\ell}} = \langle v_i, R(v_k, \overline{v_\ell})\overline{v_j}\rangle. \tag{3.67}$$

Then the symmetries of the curvature tensor are

$$R_{i\bar{j}k\bar{\ell}} = R_{k\bar{j}i\bar{\ell}} = R_{i\bar{\ell}k\bar{j}} = -R_{\bar{j}ik\bar{\ell}} = -R_{i\bar{j}\bar{\ell}k}. \tag{3.68}$$

Now the variation in ϕ of the fermion kinetic term in (3.48) is

$$\begin{aligned}&\hat{\xi}_\phi\big(\frac{1}{2}\langle\psi\not{D}\psi\rangle - \frac{1}{2}\langle(\not{D}\bar{\psi})\psi\rangle\big) \\ &\qquad = \frac{1}{2}\tilde{\Gamma}^{\mu a\dot{b}}\langle\bar{\psi}_{\dot{b}}, R(\hat{\xi}\phi,\partial_\mu\phi)\psi_a\rangle - \frac{1}{2}\tilde{\Gamma}^{\mu a\dot{b}}\langle R(\hat{\xi}\phi,\partial_\mu\phi)\bar{\psi}_{\dot{b}}, \psi_a\rangle \\ &\qquad = \tilde{\Gamma}^{\mu a\dot{b}}\langle\psi_a, R(\hat{\xi}\phi,\partial_\mu\phi)\bar{\psi}_{\dot{b}}\rangle \\ &\qquad = -\kappa\tilde{\Gamma}^{\mu a\dot{b}}R_{i\bar{j}k\bar{\ell}}\psi_a^i\bar{\psi}_{\dot{b}}^{\bar{j}}\big[\eta^e\psi_e^k(\partial_\mu\bar{\phi})^{\bar{\ell}} - \bar{\eta}^{\dot{e}}(\partial_\mu\phi)^k\bar{\psi}_{\dot{e}}^{\bar{\ell}}\big],\end{aligned} \tag{3.69}$$

where $\kappa = \sqrt{2}$. We claim this cancels against the variation in ψ of the curvature term in the lagrangian (3.48). That variation is

(3.70)

$$\begin{aligned}
&\hat{\xi}_\psi\big(-\frac{1}{4}\epsilon^{ac}\epsilon^{\dot b\dot d}R_{i\bar\jmath k\bar\ell}\psi^i_a\bar\psi^{\bar\jmath}_{\dot b}\psi^k_c\bar\psi^{\bar\ell}_{\dot d}\big)\\
&\quad= -\frac{1}{4}\kappa\epsilon^{ac}\epsilon^{\dot b\dot d}R_{i\bar\jmath k\bar\ell}\bar\eta^{\dot e}[\Gamma^\mu_{a\dot e}(\partial_\mu\phi)^i\bar\psi^{\bar\jmath}_{\dot b}\psi^k_c\bar\psi^{\bar\ell}_{\dot d}+\Gamma^\mu_{c\dot e}\psi^i_a\bar\psi^{\bar\jmath}_{\dot b}(\partial_\mu\phi)^k\bar\psi^{\bar\ell}_{\dot d}]\\
&\qquad\quad+\frac{1}{4}\kappa\epsilon^{ac}\epsilon^{\dot b\dot d}R_{i\bar\jmath k\bar\ell}\eta^e[\Gamma^\mu_{e\dot b}\psi^i_a(\partial_\mu\bar\phi)^{\bar\jmath}\psi^k_c\bar\psi^{\bar\ell}_{\dot d}+\Gamma^\mu_{e\dot d}\psi^i_a\bar\psi^{\bar\jmath}_{\dot b}\psi^k_c(\partial_\mu\bar\phi)^{\bar\ell}]\\
&\quad= -\frac{1}{2}\kappa\Gamma^\mu_{a\dot e}\epsilon^{ac}\epsilon^{\dot b\dot d}\bar\eta^{\dot e}R_{i\bar\jmath k\bar\ell}(\partial_\mu\phi)^i\bar\psi^{\bar\jmath}_{\dot b}\psi^k_c\bar\psi^{\bar\ell}_{\dot d}+\frac{1}{2}\kappa\Gamma^\mu_{e\dot b}\epsilon^{ac}\epsilon^{\dot b\dot d}\bar\eta^{\dot e}R_{i\bar\jmath k\bar\ell}\psi^i_a(\partial_\mu\bar\phi)^{\bar\jmath}\psi^k_c\bar\psi^{\bar\ell}_{\dot d}\\
&\quad= -\frac{1}{2}\kappa R_{i\bar\jmath k\bar\ell}\Big[-\Gamma^\mu_{a\dot e}\epsilon^{ac}(-\epsilon^{\dot d\dot e}\bar\eta^{\dot b}-\epsilon^{\dot e\dot b}\bar\eta^{\dot d})(\partial_\mu\phi)^i\bar\psi^{\bar\jmath}_{\dot b}\\
&\qquad\qquad+\Gamma^\mu_{e\dot b}(-\epsilon^{cd}\eta^a-\epsilon^{ea}\eta^c)\epsilon^{\dot b\dot d}\psi^i_a(\partial_\mu\bar\phi)^{\bar\jmath}\Big]\psi^k_c\bar\psi^{\bar\ell}_{\dot d}\\
&\quad= -\frac{1}{2}\kappa R_{i\bar\jmath k\bar\ell}\big[(-\tilde\Gamma^{\mu c\dot d}\bar\eta^{\dot b}+\tilde\Gamma^{\mu c\dot b}\bar\eta^{\dot d})(\partial_\mu\phi)^i\bar\psi^{\bar\jmath}_{\dot b}\\
&\qquad\qquad+(\tilde\Gamma^{\mu c\dot d}\eta^a-\tilde\Gamma^{\mu a\dot d}\eta^c)\psi^i_a(\partial_\mu\bar\phi)^{\bar\jmath}\big]\psi^k_c\bar\psi^{\bar\ell}_{\dot d}\\
&\quad= -\kappa\tilde\Gamma^{\mu c\dot d}R_{i\bar\jmath k\bar\ell}\big[\eta^e\psi^i_e(\partial_\mu\bar\phi)^{\bar\jmath}-\bar\eta^{\dot e}(\partial_\mu\phi)\bar\psi^{\bar\jmath}_{\dot e}\big]\psi^k_c\bar\psi^{\bar\ell}_{\dot d}\\
&\quad= \kappa\tilde\Gamma^{\mu a\dot b}R_{i\bar\jmath k\bar\ell}\psi^i_a\bar\psi^{\bar\jmath}_{\dot b}\big[\eta^e\psi^k_e(\partial_\mu\bar\phi)^{\bar\ell}-\bar\eta^{\dot e}(\partial_\mu\phi)^k\bar\psi^{\bar\ell}_{\dot e}\big].
\end{aligned}$$

This cancels (3.69) as claimed.

To complete the proof that (3.48) is supersymmetric, we verify that the variation in ϕ of the curvature term vanishes:

$$\begin{aligned}
(3.71)\quad \hat{\xi}_\phi\big(\epsilon^{ac}\epsilon^{\dot b\dot d}\langle\psi_a,R(\psi_c,\bar\psi_{\dot d})\bar\psi_{\dot b}\rangle\big)&=\kappa\epsilon^{ac}\epsilon^{\dot b\dot d}\eta^e\langle\psi_a,(\nabla_{\psi_e}R)(\psi_c,\bar\psi_{\dot d})\bar\psi_{\dot b}\rangle\\
&\quad-\kappa\epsilon^{ac}\epsilon^{\dot b\dot d}\bar\eta^{edd}\langle\psi_a,(\nabla_{\bar\psi_{\dot e}}R)(\psi_c,\bar\psi_{\dot d})\bar\psi_{\dot b}\rangle.
\end{aligned}$$

The Bianchi identity implies that the factor $\langle\psi_a,(\nabla_{\psi_e}R)(\psi_c,\bar\psi_{\dot d})\bar\psi_{\dot b}\rangle$ in the first term is symmetric in e,c. Combining with (3.68) we see that it is symmetric in a,e,c. Hence the first term vanishes. The argument for the second term is similar.

Finally, consider $\mathbb{F}=\mathbb{H}$. The variation in ϕ of the fermion kinetic term in (3.49) is

(3.72)

$$\begin{aligned}
\hat{\xi}_\phi\big(\frac{1}{4}\tilde h_{\alpha\beta}\nu^{abcd}\psi^\alpha_a\nabla_{cd}\psi^\beta_b\big)&=\hat{\xi}_\phi\big(\frac{1}{8}\nu^{abcd}\epsilon^{ij}\langle\psi_{ai},\nabla_{cd}\psi_{bj}\rangle\big)\\
&=\frac{1}{8}\nu^{abcd}\epsilon^{ij}\langle\psi_{ai},R(\hat\xi\phi,\partial_{cd}\phi)\psi_{bj}\rangle\\
&=\frac{1}{8}\nu^{abcd}\epsilon^{ij}R_{(\alpha i')(\beta j')(\gamma k)(\delta\ell)}\psi^{\alpha i'}_{ai}\psi^{\beta j'}_{bj}(\hat\xi\phi)^{\gamma k}(\partial_{cd}\phi)^{\delta\ell}\\
&=\frac{\kappa}{4}\nu^{abcd}\epsilon_{k\ell}\Omega_{\alpha\beta\gamma\delta}\eta^{ek}\psi^\alpha_a\psi^\beta_b\psi^\gamma_e(\partial_{cd}\phi)^{\delta\ell},
\end{aligned}$$

where $\kappa = \sqrt{2}$. This cancels against the variation in ψ of the curvature term in (3.49), which is

(3.73)

$$\begin{aligned}\hat{\xi}_\psi\big(-\frac{1}{24}\nu^{abcd}\Omega_{\alpha\beta\gamma\delta}\psi_a^\alpha\psi_b^\beta\psi_c^\gamma\psi_d^\delta\big) &= \frac{\kappa}{24}\nu^{abcd}\Omega_{\alpha\beta\gamma\delta}\epsilon_{ij}\eta^{ej}\Big[(\partial_{ae}\phi)^{\alpha i}\psi_b^\beta\psi_c^\gamma\psi_d^\delta - \psi_a^\alpha(\partial_{be}\phi)^{\beta i}\psi_c^\gamma\psi_d^\delta \\ &\qquad + \psi_a^\alpha\psi_b^\beta(\partial_{ce}\phi)^{\gamma i}\psi_d^\delta - \psi_a^\alpha\psi_b^\beta\psi_c^\gamma(\partial_{de}\phi)^{\delta i}\Big] \\ &= -\frac{\kappa}{6}\epsilon_{ij}\Omega_{\alpha\beta\gamma\delta}\nu^{abcd}\eta^{ej}\psi_b^\beta\psi_c^\gamma\psi_d^\delta(\partial_{ae}\phi)^{\alpha i}.\end{aligned}$$

Now we use the fact that $\dim S_0 = 4$, so that the cyclic sum in a, b, c, d, e of $\nu^{abcd}\eta^{ej}$ vanishes. After relabeling some indices we conclude that (3.73) simplifies to

$$\frac{\kappa}{4}\nu^{abcd}\epsilon_{k\ell}\Omega_{\alpha\beta\gamma\delta}\eta^{ek}\psi_a^\alpha\psi_b^\beta\psi_e^\gamma(\partial_{cd}\phi)^{\delta\ell}, \tag{3.74}$$

which cancels (3.72).

As in the previous cases, we complete the proof by showing that the variation in ϕ of the curvature term vanishes. That variation is a multiple of

$$\hat{\xi}_\phi\big(\nu^{abcd}\Omega(\psi_a,\psi_b,\psi_c,\psi_d)\big) = -\nu^{abcd}\eta^{ei}(\nabla_{\psi_{ei}}\Omega)(\psi_a,\psi_b,\psi_c,\psi_d). \tag{3.75}$$

Now the Bianchi identity and the symmetry of Ω imply that $(\nabla_{\psi_{ei}}\Omega)(\psi_a,\psi_b,\psi_c,\psi_d)$ is totally skew in a, b, c, d, e. Since the skew-symmetrization of $\nu^{abcd}\eta^{ei}$ vanishes ($\dim S_0 = 4$), the entire expression (3.75) vanishes.

In the nonlinear model the formulas (3.40) and (3.43) for the variational 1-form and the supercurrent are replaced by the covariant versions; that is, we use covariant derivatives in place of ordinary derivatives. We do not bother recording them here; see Theorem 6.33. The equations of motion (3.41) are affected in a more drastic way—the right hand sides are nonzero expressions in the curvature and its covariant derivative. They vanish if we set $\psi = 0$. (In fact, the terms are quadratic, cubic, and quartic in ψ.) The interested reader can work out the precise formulas, some of which appear in §4 and §5.

§3.4. Supersymmetric potential terms

To the bosonic σ-model lagrangian (3.15) with target X we can add a potential term. Let

$$V\colon X \longrightarrow \mathbb{R} \tag{3.76}$$

be a real-valued function, called the *potential energy*. We assume that V is bounded below. Then the bosonic lagrangian with potential is

$$L = \Big\{\frac{1}{2}|d\phi|^2 - \phi^*V\Big\}\,|d^nx|. \tag{3.77}$$

The moduli space of classical vacua for this theory—that is, the space of static field configurations of minimal energy—is

$$\mathcal{M}_{\text{vac}} = V^{-1}(0) \tag{3.78}$$

if we assume that the minimum value of V occurs at 0.

Now we consider supersymmetric extensions of (3.77). One basic principle, which follows from the supersymmetry algebra, is that in a supersymmetric theory the potential energy is nonnegative. That is manifest in our formulas below. Our starting point is the nonlinear σ-models developed in the previous section.

We will not carry out a detailed analysis in components, but rather content ourselves with pointing out a few elementary features. First, note that the variation of the potential term is, using (3.23),

$$\hat{\xi}(\phi^* V) = -\kappa \eta^a \phi^* \big(dV(\psi_a)\big). \tag{3.79}$$

Thus there must be an additional term in the lagrangian to cancel this. Power counting—that is, weighting the boson 0, the fermion 1/2, and a spacetime derivative 1—shows that in addition we must change the supersymmetry transformation law (3.23) for the fermion. The form of the new transformation is

$$\hat{\xi}\psi_a = \kappa \Gamma^\mu_{a\dot{b}}\, \bar{\eta}^{\dot{b}}\, \partial_\mu \phi + \kappa \epsilon_{ab} \eta^b \phi^* F_a \tag{3.80}$$

for some vector fields F_a on X, and the additional term in the lagrangian takes the form

$$\phi^* A^{ab}(\psi_a, \psi_b) \tag{3.81}$$

for some bilinear forms A^{ab} on X. Imposing the condition that the lagrangian be supersymmetric gives relations among V, A^{ab}, F_a. In fact, one can analyze those conditions in the dimensional reduction of the models to one spacetime dimension. Such an analysis leads to the form of the lagrangians in the following result.

Theorem 3.82. (*i*) ($\mathbb{F} = \mathbb{R}$) *Let X be a Riemannian manifold and $h\colon X \to \mathbb{R}$ a real function. Then the lagrangian*

$$\boxed{\begin{aligned} L = \Big\{ \frac{1}{2}|d\phi|^2 + \frac{1}{2}\langle \psi \not{D}_\phi \psi\rangle + \frac{1}{12}\epsilon^{ab}\epsilon^{cd}\langle \psi_a, \phi^* R(\psi_b, \psi_c)\psi_d\rangle \\ - \frac{1}{2}\phi^* |\operatorname{grad} h|^2 - \frac{1}{2}\epsilon^{ab}\phi^*(\operatorname{Hess} h)(\psi_a, \psi_b)\Big\}\, |d^3x|. \end{aligned}} \tag{3.83}$$

is supersymmetric. The supersymmetry transformation law is

$$\boxed{\begin{aligned} \hat{\xi}\phi &= -\eta^a\, \psi_a \\ \hat{\xi}\psi_a &= \Gamma^\mu_{ab}\, \eta^b\, \partial_\mu \phi + \epsilon_{ab}\eta^b \phi^* \operatorname{grad} h \end{aligned}} \tag{3.84}$$

(*ii*) ($\mathbb{F} = \mathbb{C}$) *Let X be a Kähler manifold and $W\colon X \to \mathbb{R}$ a holomorphic function. Then the lagrangian*

(3.85)

$$\boxed{\begin{aligned} L = \Big\{ \langle d\bar{\phi}, d\phi\rangle + \frac{1}{2}\langle \bar{\psi} \not{D}_\phi \psi\rangle - \frac{1}{2}\langle (\not{D}_\phi \bar{\psi})\psi\rangle - \frac{1}{4}\epsilon^{ac}\epsilon^{\dot{b}\dot{d}}\langle \psi_a, \phi^* R(\psi_c, \bar{\psi}_{\dot{d}})\bar{\psi}_{\dot{b}}\rangle \\ - \phi^* \|\operatorname{grad} W\|^2 - \operatorname{Re}\big[\epsilon^{ab}\phi^*(\operatorname{Hess} W)(\psi_a, \psi_b)\big]\Big\}\, |d^4x|. \end{aligned}}$$

is supersymmetric. The supersymmetry transformation law is

$$\boxed{\begin{aligned}\hat{\xi}\phi &= -\sqrt{2}\,\eta^a\,\psi_a \\ \hat{\xi}\psi_a &= \sqrt{2}\,\Gamma^\mu_{a\dot{b}}\,\bar{\eta}^{\dot{b}}\,\partial_\mu\phi + \sqrt{2}\,\epsilon_{ab}\eta^b\phi^*\,\operatorname{grad} W\end{aligned}} \tag{3.86}$$

Here $\operatorname{Hess} h$ is the covariant hessian

$$\operatorname{Hess} h = \nabla dh, \tag{3.87}$$

which is a symmetric bilinear form. Similarly,

$$\operatorname{Hess} W = \nabla\partial W. \tag{3.88}$$

The double bars denote the hermitian norm:

$$\|\operatorname{grad} W\|^2 = \langle\overline{\operatorname{grad} W}, \operatorname{grad} W\rangle. \tag{3.89}$$

For $\mathbb{F} = \mathbb{R}$ the potential energy is

$$V_{\mathbb{R}} = \frac{1}{2}|\operatorname{grad} h|^2; \tag{3.90}$$

for $\mathbb{F} = \mathbb{C}$ it is

$$V_{\mathbb{C}} = \|\operatorname{grad} W\|^2. \tag{3.91}$$

In both cases it is nonnegative. The function W is called the *superpotential*. The moduli space of classical vacua is

$$\mathcal{M}^{\mathbb{R}}_{\text{vac}} = \operatorname{Crit}(h) \tag{3.92}$$

for $\mathbb{F} = \mathbb{R}$ and

$$\mathcal{M}^{\mathbb{C}}_{\text{vac}} = \operatorname{Crit}(W) \tag{3.93}$$

for $\mathbb{F} = \mathbb{C}$. Here 'Crit' denotes the set of critical points.

We defer the proof of Theorem 3.82 to §4 and §5, where we give a manifestly supersymmetric version in superspace. There we also derive the supercurrent for the model with potential.

For the $\mathbb{F} = \mathbb{H}$ case there is no way to add a potential energy term and preserve supersymmetry. Presumably this follows in a straightforward way from the component analysis indicated above. On the other hand, the dimensional reduction of this 6-dimensional theory to 4 dimensions (and below) permits a supersymmetric potential energy term—the bosonic potential energy V is the norm square of a vector field on X which preserves the hyperkähler structure.

§3.5. Superspace construction

Consider a superspace $M^{n|s}$ in any dimension built out of any real spin representation, and let X be any Riemannian manifold. Introduce a scalar superfield

$$\Phi\colon M^{n|s} \longrightarrow X. \tag{3.94}$$

Let $i\colon \check{M}^n \hookrightarrow M^{n|s}$ be the inclusion. There are component fields

$$\begin{aligned} \phi &= i^*\Phi \\ \psi_a &= i^* D_a \Phi \end{aligned} \tag{3.95}$$

of the type we need, but in general higher derivatives of Φ lead to more components.

It is tempting to write a lagrangian

$$L = \check{L}\,|d^n x| \tag{3.96}$$

of the form $\check{L} = i^*S$ for some function S on superspace. We need 4 fermionic derivatives in superspace to have 2 spacetime derivatives on the scalar ϕ, and the natural invariant expression is

$$\check{L} = i^*\Big\{ k g_{\mu\nu} \tilde{\Gamma}^{\mu ac} \tilde{\Gamma}^{\nu bd} D_a D_b \langle D_c\Phi, D_d\Phi\rangle \Big\}, \tag{3.97}$$

where k is a constant. In fact, the term

$$\check{L} = i^*\Big\{ -k g_{\mu\nu} \tilde{\Gamma}^{\mu ac} \tilde{\Gamma}^{\nu bd} \langle D_a D_c\Phi, D_b D_d\Phi\rangle \Big\} \tag{3.98}$$

contains a multiple of the bosonic lagrangian (3.15), and this fixes the constant k. It is not hard to see that we also obtain the kinetic term for the fermion ψ. In addition—if there are at least 2 supersymmetries—we find terms with no derivatives built out of *auxiliary fields*, which are restrictions to $\check{M}^n$ of second derivatives of Φ. But in general this construction fails to give a supersymmetric lagrangian: there are too many component fields. We can eliminate some of them from the supersymmetry transformation laws derived from (1.29), but there is no guarantee that (3.97) is supersymmetric, nor that the supersymmetry algebra closes on-shell. In fact, this construction does work for $s = 1$ supersymmetry (1.52) and $s = 2$ supersymmetries (§4). For $s = 1$ there are no auxiliary fields; for $s = 2$ there is an auxiliary real scalar field. For $s = 4$ supersymmetries (§5) the superspace model presented here works if we impose a *constraint* on Φ which eliminates many of the component fields. There is then a single auxiliary complex scalar field. In all of these cases we express (3.97) in a form which is manifestly supersymmetric.

§3.6. Dimensional reduction

In a standard way we can reduce a theory on $\check{M}^n$ to a theory on $\check{M}^m$ for $m < n$ by considering fields on $\check{M}^n$ invariant under an $(n-m)$-dimensional space of spatial translations. The scalar field ϕ reduces to a scalar field on $\check{M}^m$, and the spinor field ψ reduces to (nonchiral) spinor fields on $\check{M}^m$. (A single spinor field on $\check{M}^n$ gives possibly many spinor fields on $\check{M}^m$. For example, the 8-component chiral spinor field in the 6-dimensional $\mathbb{F} = \mathbb{H}$ model reduces to 4 right handed and 4 left handed

spinor fields in $\check{M}^2$.) The reduced model has the same number of supersymmetries as the original.

In particular, we can relate the three supersymmetric σ-models considered here. For example, the dimensional reduction of the linear $\mathbb{F} = \mathbb{C}$ model to 3 dimensions gives two copies of the $\mathbb{F} = \mathbb{R}$ model. Similarly, the dimensional reduction of the nonlinear $\mathbb{F} = \mathbb{C}$ model with target a Kähler manifold X gives the nonlinear $\mathbb{F} = \mathbb{R}$ model with target the underlying real manifold. In this case the fermion $\hat{\psi}$ of the 3-dimensional model is defined in terms of the fermion ψ of the 4-dimensional model by

$$\boxed{\hat{\psi} = \frac{\psi + \bar{\psi}}{\sqrt{2}}.} \tag{3.99}$$

Then the lagrangian (3.48) reduces to twice the lagrangian (3.47). For example, if we plug (3.99) into the curvature term in (3.47) we have 6 nonzero terms with a factor of $1/48$ in front, and twice $6 \cdot 1/48$ is $1/4$. The sign also works. The dimensional reduction of the potential term (3.85) in the $\mathbb{F} = \mathbb{C}$ model gives the potential term (3.83) with

$$\boxed{h = \operatorname{Re} W.} \tag{3.100}$$

To dimensionally reduce the $\mathbb{F} = \mathbb{H}$ model to 4 dimensions, we must fix a Kähler structure on the hyperkähler target manifold X. In fact, there is a 2-sphere of such Kähler structures parametrized by the unit imaginary quaternions acting on the vector space W^* (cf. (3.51)). We choose the basis $\{e_1, e_2\}$ of W so that $V \otimes \mathbb{C}\, e^1$ is the bundle of type $(1,0)$ tangent vectors to X and $V \otimes \mathbb{C}\, e^2$ the bundle of type $(0,1)$ tangent vectors to X. As for the spinor fields, recall from §2.2 that under dimensional reduction we identify $S_0 \cong S' \oplus S''$, where $S' \oplus S''$ is the complexified spinor representation of $\mathrm{Spin}(1,3)$. Thus the spinor fields of the dimensionally reduced theory transform in $(S' \oplus S'') \otimes \phi^* V$. In indices, if $\hat{\xi}^{\alpha i}$ is a complex tangent vector to X, then we define

$$\boxed{\xi^\alpha = \hat{\xi}^{\alpha 1}, \qquad \bar{\xi}^{\bar{\alpha}} = \hat{\xi}^{\alpha 2}.} \tag{3.101}$$

Note ξ is real if $\bar{\xi}^{\bar{\alpha}}$ is the complex conjugate of ξ^α. If $\hat{\psi}_a^\alpha$ is a spinor field on $\check{M}^6$, we define a spinor field ψ on $\check{M}^4$ by

$$\boxed{\begin{aligned} \psi_1^\alpha &= \hat{\psi}_1^\alpha, & \bar{\psi}_{\dot{1}}^{\bar{\alpha}} &= J_\beta^{\bar{\alpha}} \hat{\psi}_3^\beta, \\ \psi_2^\alpha &= \hat{\psi}_2^\alpha, & \bar{\psi}_{\dot{2}}^{\bar{\alpha}} &= J_\beta^{\bar{\alpha}} \hat{\psi}_4^\beta, \end{aligned}} \tag{3.102}$$

where J is the pseudoreal structure on V. The reality condition on ψ is that $\bar{\psi}_{\dot{a}}^{\bar{\alpha}}$ is the complex conjugate of ψ_a^α. Armed with these formulas, together with (2.99) and (3.55), we dimensionally reduce (3.49) to $\check{M}^4$, i.e., we evaluate (3.49) on fields ϕ, ψ which satisfy $\partial_{12}\phi = \partial_{12}\psi = 0$. A bit of computation shows that we recover (3.48).

CHAPTER 4
The Supersymmetric σ-Model in Dimension 3

In this section we discuss the σ-model with 2 supersymmetries. We already described it in components in Chapter 3. Here we give a manifestly supersymmetric treatment in superspace. We begin in §4.1 by deriving the component fields and supersymmetry transformation laws, following the general principles laid out in §1.2. Then in §4.2 we state the superspace lagrangian and derive the component lagrangian. We recover the lagrangian of Theorem 3.46(i) except with the addition of an auxiliary field. One virtue of the auxiliary field is that now the bracketing relations among the supersymmetry transformations (4.14) are precisely those of the super Poincaré algebra; without the auxiliary field we need to impose the equations of motion to get the correct algebra. In §4.3 we prove Theorem 3.82(i), which describes the supersymmetric potential term. We carry out the analysis of the classical theory in §4.4—we compute the variational 1-form, equations of motion, symplectic structure, supercurrent—directly in superspace and from that rederive the component expressions obtained in Chapter 3. We conclude in §4.5 by briefly considering the dimensional reduction to 2 dimensions, where the Poisson brackets of the supercharges leads to a central extension of the super Poincaré algebra.

§4.1. Fields and supersymmetry transformations on $M^{3|2}$

Linear Case. We begin with a real *scalar superfield*

$$\Phi\colon M^{3|2} \longrightarrow \mathbb{R}. \tag{4.1}$$

Define the component fields as

$$\begin{aligned} \phi &= i^*\Phi \\ \psi_a &= i^* D_a \Phi \\ F &= i^*(-D^2)\Phi. \end{aligned} \tag{4.2}$$

So ϕ is a real function on Minkowski space $\check{M}^3$, the field $\psi = \psi_a f^a$ is a spinor field on $\check{M}^3$, and F is again a real function on $\check{M}^3$. Note that ϕ and F are *even* whereas ψ is *odd*. We will see that the field F only enters algebraically, and its equations of motion are algebraic. Such a field is termed *auxiliary* since we can solve for it algebraically in terms of the other fields. The collection of component fields is termed a *multiplet*, so here (ϕ, ψ, F) is a *real scalar multiplet*.

Let $\hat{\xi}$ be the vector field on the space $\mathcal{F}_{(\phi,\psi,F)}$ of component fields induced by the supersymmetry transformation $\eta^a Q_a$, where η^a are odd parameters. We compute the action of $\hat{\xi}$ on component fields using (1.29). For the lowest component ϕ this is straightforward:

$$\hat{\xi}\phi = -\eta^a i^* D_a \Phi = -\eta^a \psi_a. \tag{4.3}$$

For the action on ψ_a we use (2.42):

$$\begin{aligned} \hat{\xi}\psi_a &= -\eta^b i^* D_b D_a \Phi \\ &= \eta^b i^* (\partial_{ab} - \epsilon_{ba} D^2)\Phi \\ &= \eta^b \partial_{ab}\phi - \epsilon_{ab}\eta^b F. \end{aligned} \tag{4.4}$$

To compute the action on F we use (2.43):

$$\begin{aligned} \hat{\xi}F &= -\eta^a i^* D_a(-D^2)\Phi \\ &= -\eta^a \epsilon^{bc} i^* \partial_{ab} D_c \Phi \\ &= -\eta^a \epsilon^{bc} \partial_{ab}\psi_c \\ &= \eta^a (\not{D}\psi)_a. \end{aligned} \tag{4.5}$$

The reader should check the supersymmetry algebra (1.30) and should also compare (4.3) and (4.4) to (3.23).

Nonlinear Case. For the *nonlinear* σ-model into a Riemannian manifold X the superfield is a map

$$\Phi\colon M^{3|2} \longrightarrow X. \tag{4.6}$$

The components are defined by the (covariant) extensions of (4.2) using the Levi-Civita covariant derivative:

$$\boxed{\begin{aligned} \phi &= i^*\Phi \\ \psi_a &= i^* D_a \Phi \\ F &= i^*(-\mathcal{D}^2)\Phi. \end{aligned}} \tag{4.7}$$

Now $\phi\colon \check{M}^3 \to X$ is a map into the target manifold, i.e., the field of the underlying bosonic σ-model. Then $\psi = \psi_a f^a$ is a spinor field on $\check{M}^3$ with values in ϕ^*TX, i.e., $\psi\colon \check{M}^3 \to \Pi S \otimes \phi^*TX$. (Recall that 'Π' denotes parity reversal.) For pedagogical purposes we use the symbol '$\mathcal{D}_a$' to denote the covariant derivative of a section of ϕ^*TX in the direction D_a and '∇_{ab}' for the covariant derivative in the direction ∂_{ab}. Then $\mathcal{D}^2 = \frac{1}{2}\epsilon^{ab}\mathcal{D}_a\mathcal{D}_b$, and F is a scalar with values in ϕ^*TX, i.e., a section of ϕ^*TX on $\check{M}^3$.

To compute with the nonlinear superfield Φ it is useful to recall that

$$\mathcal{D}_a D_b \Phi = \mathcal{D}_b D_a \Phi, \tag{4.8}$$

since the Levi-Civita connection is torsionfree. Also,

$$R(D_a, D_b) = \mathcal{D}_a\mathcal{D}_b + \mathcal{D}_b\mathcal{D}_a + 2\nabla_{ab} \qquad \text{on } \Phi^*TX \tag{4.9}$$

by the definition of the curvature R of Φ^*TX and the sign rule. One must also be careful with the sign rule when applying the Bianchi identity.

Now the analogs of equations (2.43) for covariant derivatives are easily derived:

$$\mathcal{D}_a\mathcal{D}^2\Phi = -\epsilon^{bc}\big(\nabla_{ab} - \frac{1}{3}R(D_a\Phi, D_b\Phi)\big)D_c\Phi, \tag{4.10}$$

$$\mathcal{D}^2 D_a\Phi = \quad \epsilon^{bc}\big(\nabla_{ab} - \frac{1}{6}R(D_a\Phi, D_b\Phi)\big)D_c\Phi. \tag{4.11}$$

To illustrate the computations, we derive (4.11) for $a = 1$:

$$\begin{aligned} 2\mathcal{D}^2 D_1\Phi &= \mathcal{D}_1\mathcal{D}_2 D_1\Phi - \mathcal{D}_2\mathcal{D}_1 D_1\Phi \\ &= \big[R(D_1\Phi, D_2\Phi)D_1\Phi - \mathcal{D}_2\mathcal{D}_1 D_1\Phi - 2\nabla_{12}D_1\Phi\big] + \mathcal{D}_2\partial_{11}\Phi \\ &= 2D_2\partial_{11}\Phi - 2\nabla_{12}D_1\Phi + R(D_1\Phi, D_2\Phi)D_1\Phi \\ &= 2\nabla_{11}D_2\Phi - 2\nabla_{12}D_1\Phi + R(D_1\Phi, D_2\Phi)D_1\Phi. \end{aligned} \tag{4.12}$$

At the last stage, and also when we compare (4.12) with (4.11), we use the Bianchi identity $R(D_1\Phi, D_1\Phi)D_2\Phi = -2R(D_1\Phi, D_2\Phi)D_1\Phi$.

Using (4.10) and (4.11) it is a routine matter to derive the transformation laws for the fields in the nonlinear case. In fact, the only change comes in (4.5):

$$\begin{aligned} \hat{\xi}F &= -\eta^a i^*\mathcal{D}_a(-\mathcal{D}^2)\Phi \\ &= -\eta^a\epsilon^{bc}i^*\big(\nabla_{ab} - \frac{1}{3}R(D_a\Phi, D_b\Phi)\big)D_c\Phi \\ &= -\eta^a\epsilon^{bc}\big(\nabla_{ab} - \frac{1}{3}R(\psi_a, \psi_b)\big)\psi_c \\ &= \quad \eta^a\big[(\not{D}\psi)_a + \frac{1}{3}\epsilon^{bc}R(\psi_a, \psi_b)\psi_c\big]. \end{aligned} \tag{4.13}$$

For convenience we collect the transformation formulas for the component fields:

$$\boxed{\begin{aligned} \hat{\xi}\phi &= -\eta^a\psi_a \\ \hat{\xi}\psi_a &= \quad \eta^b(\partial_{ab}\phi - \epsilon_{ab}F) \\ \hat{\xi}F &= \quad \eta^a\big[(\not{D}\psi)_a + \frac{1}{3}\epsilon^{bc}R(\psi_a, \psi_b)\psi_c\big]. \end{aligned}} \tag{4.14}$$

These formulas have easy reductions to $M^{2|2}$. As mentioned after (2.106) we simply set $\partial_{12} = \nabla_{12} = 0$ and relabel indices as 1=+ and 2=−:

$$\boxed{\begin{aligned} \hat{\xi}\phi &= -\eta^+\psi_+ - \eta^-\psi_- \\ \hat{\xi}\psi_+ &= \eta^+\partial_+\phi - \eta^- F \\ \hat{\xi}\psi_- &= \eta^-\partial_-\phi + \eta^+ F \\ \hat{\xi}F &= -\eta^+\nabla_+\psi_- + \eta^-\nabla_-\psi_+ - \eta^+R(\psi_+, \psi_-)\psi_+ + \eta^-R(\psi_-, \psi_+)\psi_-. \end{aligned}} \tag{4.15}$$

§4.2. The σ-model action on $M^{3|2}$

Since the computations in the nonlinear case are no more difficult than in the linear case, we proceed directly to it. So our superfield (4.6) is a map $\Phi\colon M^{3|2} \to X$ into a Riemannian manifold X. The lagrangian density on $M^{3|2}$ is

$$\boxed{L_0 = |d^3x|\, d^2\theta\, \frac{1}{4}\epsilon^{ab}\langle D_a\Phi, D_b\Phi\rangle,} \tag{4.16}$$

where $\langle\cdot,\cdot\rangle$ is the metric on X pulled back to a metric on Φ^*TX. Our goal is to derive the component lagrangian as defined in (1.33). Recall that in this case we use integration over the θ variables, expressed as differentiation in (2.40). The resulting lagrangian *function* on Minkowski space is

$$\check{L}_0 = \int d^2\theta\, \frac{1}{4}\epsilon^{ab}\langle D_a\Phi, D_b\Phi\rangle. \tag{4.17}$$

We carry out the integration using (2.40), at intermediate stages using (4.11) and the covariant analog of (2.42), and finally using the definition of the component fields (4.7) and formulas (2.26), (2.28) to restrict to Minkowski space:

(4.18)

$$\begin{aligned}
4\check{L}_0 &= -\epsilon^{ab} i^* D^2\langle D_a\Phi, D_b\Phi\rangle \\
&= i^*\left\{-2\epsilon^{ab}\langle D_a\Phi, \mathcal{D}^2 D_b\Phi\rangle + \epsilon^{ab}\epsilon^{cd}\langle \mathcal{D}_c D_a\Phi, \mathcal{D}_d D_b\Phi\rangle\right\} \\
&= i^*\Big\{-2\epsilon^{ab}\epsilon^{cd}\langle D_a\Phi, \nabla_{bc} D_d\Phi\rangle + \frac{1}{3}\epsilon^{ab}\epsilon^{cd}\langle D_a\Phi, R(D_b\Phi, D_c\Phi)D_d\Phi\rangle \\
&\qquad\qquad + \epsilon^{ab}\epsilon^{cd}\langle \partial_{ca}\Phi - \epsilon_{ca}\mathcal{D}^2\Phi\ ,\ \partial_{db}\Phi - \epsilon_{db}\mathcal{D}^2\Phi\rangle\Big\} \\
&= 2\langle\psi \not{D}\psi\rangle + \frac{1}{3}\epsilon^{ab}\epsilon^{cd}\langle\psi_a, R(\psi_b,\psi_c)\psi_d\rangle + 2|d\phi|^2 + 2|F|^2.
\end{aligned}$$

Of course, here the Dirac form (2.28) is defined with a covariant derivative. So the final formula is[30]

$$\boxed{\check{L}_0 = \frac{1}{2}|d\phi|^2 + \frac{1}{2}\langle\psi \not{D}_\phi\psi\rangle + \frac{1}{12}\epsilon^{ab}\epsilon^{cd}\langle\psi_a, \phi^*R(\psi_b,\psi_c)\psi_d\rangle + \frac{1}{2}|F|^2.} \tag{4.19}$$

The resulting equations of motion imply that $F = 0$. Putting $F = 0$ in (4.19) we recover (3.47).

§4.3. The potential term on $M^{3|2}$

Let $h\colon X \to \mathbb{R}$ be a real-valued function on X, and set

$$\boxed{L_1 = |d^3x|\, d^2\theta\ \Phi^*(h).} \tag{4.20}$$

[30] We put 'ϕ' and 'ϕ^*' into the notation to remind ourselves that these terms contribute to the equation of motion for ϕ.

The expansion in components is

$$\begin{aligned}\check{L}_1 &= -\frac{1}{2}i^*\epsilon^{ab}D_aD_b\Phi^*(h)\\ &= -\frac{1}{2}i^*\epsilon^{ab}D_a\big(\iota(D_b\Phi)dh\big)\\ &= -\frac{1}{2}i^*\epsilon^{ab}\big(\iota(\mathcal{D}_aD_b\Phi)dh - \iota(D_b\Phi)D_a\Phi^*dh\big)\\ &= \iota(F)\phi^*dh - \frac{1}{2}\epsilon^{ab}\phi^*(\operatorname{Hess} h)(\psi_a,\psi_b).\end{aligned} \tag{4.21}$$

In this expression $\operatorname{Hess} h = \nabla dh$ is the covariant hessian of h, a symmetric tensor on X. We include 'ϕ^*' in our equations to make clear the dependence of ϕ. In the total lagrangian function $\check{L}_0 + \check{L}_1$ the field F still enters algebraically, and its equation of motion is

$$\boxed{F = -\phi^*\operatorname{grad} h.} \tag{4.22}$$

Substituting into the lagrangian we obtain

$$\begin{aligned}\check{L}_0 + \check{L}_1 \sim \frac{1}{2}|d\phi|^2 + \frac{1}{2}\langle\psi\not{D}_\phi\psi\rangle + \frac{1}{12}\epsilon^{ab}\epsilon^{cd}\langle\psi_a,\phi^*R(\psi_b,\psi_c)\psi_d\rangle\\ - \frac{1}{2}\phi^*|\operatorname{grad} h|^2 - \frac{1}{2}\epsilon^{ab}\phi^*(\operatorname{Hess} h)(\psi_a,\psi_b).\end{aligned} \tag{4.23}$$

The '$\sim$' indicates that we have eliminated auxiliary fields. This is precisely the lagrangian (3.83), and since the superspace model is manifestly supersymmetric we have proved Theorem 3.82(i). The potential energy for the bosons is

$$V = \frac{1}{2}|\operatorname{grad} h|^2. \tag{4.24}$$

It is nonnegative, as we expect in general for a supersymmetric lagrangian. The bilinear form in ψ is a mass term for the fermions, which we abbreviate as

$$\phi^*(\operatorname{Hess} h)(\psi,\psi). \tag{4.25}$$

A *vacuum solution* of a classical field theory on Minkowski space is a field configuration with all fermions set to zero and all scalar fields set to constants which minimize the energy. (If there are gauge fields, then they are chosen to be trivial.) For $\check{L}_0 + \check{L}_1$ this means that $\psi \equiv 0$ and ϕ is a constant which is a critical point of (4.24). Now

$$dV = \langle\operatorname{grad} h, \nabla\operatorname{grad} h\rangle. \tag{4.26}$$

This certainly vanishes at critical points of h. At such points the potential energy vanishes and supersymmetry is *unbroken*. This means that the solution is annihilated by the supersymmetry transformation (4.14). For any vacuum solution (4.14) simplifies to (using (4.22))

$$\begin{aligned}\hat{\xi}\phi &= 0\\ \hat{\xi}\psi_a &= \eta^b\epsilon_{ab}\phi^*\operatorname{grad} h,\end{aligned} \tag{4.27}$$

and clearly $\hat{\xi}\psi = 0$ only at a critical point of h. Thus at critical points of V which are *not* critical points of h, supersymmetry is broken. At such a point $\operatorname{grad} h$ is in the kernel of the mass form (4.25) for the fermions. In other words, at such a point there is a massless fermion, the so-called *Goldstone fermion*.

§4.4. Analysis of the classical theory

We first compute the variation of the lagrangian density L_0+L_1 under an arbitrary *even* variation $\delta\Phi$ of the superfield Φ. One should interpret L_0+L_1 as living on $\mathcal{F}_\Phi \times M^{3|2}$, where $\mathcal{F}_\Phi$ is the supermanifold of superfields Φ. Then 'δ' is the component of the differential along $\mathcal{F}_\Phi$. We have

$$\delta(L_0+L_1) = |d^3x|\,d^2\theta\,\Big\{\frac{1}{2}\epsilon^{ab}\langle\delta_\nabla D_a\Phi, D_b\Phi\rangle + \iota(\delta\Phi)dh\Big\}, \tag{4.28}$$

where δ_∇ is the extension of the differential δ using the covariant derivative. The first term in braces in (4.28) is

$$\begin{aligned} &|d^3x|\,d^2\theta\,\frac{1}{2}\epsilon^{ab}\langle\delta_\nabla D_a\Phi, D_b\Phi\rangle \\ &\qquad = |d^3x|\,d^2\theta\,\frac{1}{2}\epsilon^{ab}\langle\mathcal{D}_a\delta\Phi, D_b\Phi\rangle \\ &\qquad = |d^3x|\,d^2\theta\,\frac{1}{2}\epsilon^{ab}\Big\{D_a\langle\delta\Phi, D_b\Phi\rangle - \langle\delta\Phi, \mathcal{D}_aD_b\Phi\rangle\Big\} \\ &\qquad = d\Big\{\iota(D_a)\big(|d^3x|\,d^2\theta\,\frac{1}{2}\epsilon^{ab}\langle D_b\Phi, \delta\Phi\rangle\big)\Big\} - |d^3x|\,d^2\theta\,\langle\delta\Phi, \mathcal{D}^2\Phi\rangle. \end{aligned} \tag{4.29}$$

At the last stage we use the fact that the canonical density $|d^3x|\,d^2\theta$ is invariant under D_a and we use the Cartan formula for the Lie derivative. So altogether

$$\begin{aligned} \delta(L_0+L_1) = d\Big\{\iota(D_a)\big(|d^3x|\,d^2\theta\,\frac{1}{2}\epsilon^{ab}\langle D_b\Phi, \delta\Phi\rangle\big)\Big\} \\ - \ |d^3x|\,d^2\theta\,\langle\delta\Phi\,,\,\mathcal{D}^2\Phi - \Phi^*\operatorname{grad} h\rangle. \end{aligned} \tag{4.30}$$

From this we read off the equation of motion in superspace

$$\boxed{\mathcal{D}^2\Phi = \Phi^*\operatorname{grad} h} \tag{4.31}$$

and the variational 1-form

$$\boxed{\gamma = -\iota(D_a)|d^3x|\,d^2\theta\,\frac{1}{2}\epsilon^{ab}\langle D_b\Phi, \delta\Phi\rangle.} \tag{4.32}$$

The appellation '1-form' refers to the fact that γ is a 1-form on the space $\mathcal{F}_\Phi$ of superfields. With respect to $M^{3|2}$ it is an integral density of degree -1, written here as a Berezinian contracted with a vector field, i.e., $\gamma \in \Omega^{1,|-1|}(\mathcal{F}_\Phi \times M^{3|2})$. The differential $\delta\gamma$ is $\omega \in \Omega^{2,|-1|}(\mathcal{F}_\Phi \times M^{3|2})$:

$$\boxed{\omega = \iota(D_a)|d^3x|\,d^2\theta\,\frac{1}{2}\epsilon^{ab}\langle\mathcal{D}_b\delta\Phi\wedge\delta\Phi\rangle.} \tag{4.33}$$

The 2-form ω is a local version of the symplectic form on the space of classical solutions $\mathcal{M} \subset \mathcal{F}_\Phi$; the global symplectic 2-form on $\mathcal{M}$ is obtained by integrating ω over a spacelike hypersurface of codimension $1|0$ in $M^{3|2}$.

This completes the analysis in superspace. Now we expand in components. We begin with the equation of motion (4.31). Restricting to Minkowski space, we

simply recover the equation of motion (4.22) for F. The equation of motion for ψ is found by applying $i^*\mathcal{D}_a$ to (4.31) and using (4.10):

$$\boxed{(\not{D}_\phi\psi)_a = -\epsilon^{bc}\,\phi^*\nabla_{ab}\psi_c = -\frac{1}{3}\epsilon^{bc}\phi^* R(\psi_a,\psi_b)\psi_c + \phi^*(\nabla\,\mathrm{grad}\,h)(\psi_a).} \tag{4.34}$$

The equation of motion for ϕ is a complicated expression which schematically is

$$\boxed{\Box\phi = -\frac{1}{2}\phi^*\,\mathrm{grad}\,|dh|^2 + Q^{(2)}_\phi(\psi) + R^{(2)}_\phi(\psi)d\phi + Q^{(4)}_\phi(\psi),} \tag{4.35}$$

where '$\Box$' is the covariant wave operator

$$\Box = \nabla_{11}\partial_{22} - \nabla_{12}\partial_{12} \tag{4.36}$$

and $Q^{(2)}_\phi$, $R^{(2)}_\phi$, and $Q^{(4)}_\phi$ are forms in ψ whose coefficients are made from the pullback of derivatives of h and derivatives of the curvature R. The precise formula is not of interest to us here. The *bosonic* classical equation for ϕ, obtained by setting the fermion $\psi = 0$, is

$$\Box\phi = -\phi^*\,\mathrm{grad}\,V, \tag{4.37}$$

where V is the potential energy (4.24). Newton's law at last!

Next we compute $\check{\gamma}, \check{\omega}$ which are obtained from γ, ω by integrating out the θ's. Recall from (2.16) that $D_a = \partial_a - \theta^c\partial_{ac}$. Now in (4.32) the term with $\iota(\partial_a)$ drops out after $\int d^2\theta$, so

(4.38)

$$\begin{aligned}
\check{\gamma} &= \iota(\partial_{ac})\,|d^3x|\ \int d^2\theta\ \frac{1}{2}\epsilon^{ab}\theta^c\langle D_b\Phi, \delta\Phi\rangle \\
&= -\iota(\partial_{ac})\,|d^3x|\,\Big\{\frac{1}{4}\epsilon^{ab}\epsilon^{de}i^* D_d D_e\theta^c\langle D_b\Phi,\delta\Phi\rangle\Big\} \\
&= -\iota(\partial_{ac})\,|d^3x|\,\Big\{-\frac{1}{2}\epsilon^{ab}\epsilon^{cd}i^*\big[\langle\mathcal{D}_d D_b\Phi,\delta\Phi\rangle - \langle D_d\Phi,\delta D_b\Phi\rangle\big]\Big\} \\
&= -\iota(\partial_{ac})\,|d^3x|\,\Big\{\frac{1}{2}\epsilon^{ab}\epsilon^{cd}\big[\langle\partial_{db}\phi,\delta\phi\rangle + \epsilon_{db}\langle F,\delta\phi\rangle + \langle\psi_b,\delta\psi_d\rangle\big]\Big\}.
\end{aligned}$$

Note that '$\delta\psi_d$' is a covariant derivative of ψ_d. Simplifying further we obtain

$$\boxed{\begin{aligned}
\check{\gamma} &= -\iota(\partial_{cd})\,|d^3x|\,\Big\{\frac{1}{2}\epsilon^{bd}\epsilon^{ac}\big[\langle\partial_{ab}\phi,\delta\phi\rangle + \langle\psi_a,\delta\psi_b\rangle\big]\Big\} \\
&= -\iota(\partial_\mu)\ \ |d^3x|\,\Big\{g^{\mu\nu}\langle\partial_\nu\phi,\delta\phi\rangle + \frac{1}{2}\langle\psi\,\tilde{\Gamma}^\mu\,\delta\psi\rangle\Big\}.
\end{aligned}} \tag{4.39}$$

This last expression is written in terms of the basic bilinear forms: the inverse metric $g^{\mu\nu}$ and the paring $\tilde{\Gamma}^\mu$ of (1.2). It agrees with (3.40). As expected, no derivatives of fermions and only one derivative of a boson enter into the formula for $\check{\gamma}$, and the

potential terms do not contribute. We compute $\check{\omega}$ analogously from (4.33), or by simply differentiating (4.39). The results agree:

(4.40)
$$\boxed{\begin{aligned}\check{\omega} &= \iota(\partial_{cd})\,|d^3x|\,\Big\{\frac{1}{2}\epsilon^{bd}\epsilon^{ac}\big[\langle\partial_{ab}\delta\phi\wedge\delta\phi\rangle + \langle\delta\psi_a\wedge\delta\psi_b\rangle + \langle\psi_a\wedge R_\nabla(\delta\phi\wedge\delta\phi)\psi_b\rangle\big]\Big\}\\ &= \iota(\partial_\mu)\ |d^3x|\,\Big\{g^{\mu\nu}\langle\partial_\nu\delta\phi\wedge\delta\phi\rangle + \frac{1}{2}\langle\delta\psi\tilde{\Gamma}^\mu\,\delta\psi\rangle + \frac{1}{2}\langle\psi\tilde{\Gamma}^\mu R_\nabla(\delta\phi\wedge\delta\phi)\psi\rangle\Big\}.\end{aligned}}$$

We compute the global symplectic form Ω on the space of solutions $\mathcal{M}$ by integrating $\check{\omega}$ over a spacelike hypersurface. Using coordinates x^0, x^1, x^2 as in (5.8), with x^0 representing time, and integrating over a constant time, we obtain

(4.41)
$$\boxed{\Omega = \int_{x^0=\text{const}} dx^1dx^2\,\Big\{\langle\delta\dot{\phi}\wedge\delta\phi\rangle - \frac{1}{2}\langle\delta\psi\tilde{\Gamma}^0\delta\psi\rangle + \frac{1}{2}\langle\psi\tilde{\Gamma}^0R_\nabla(\delta\phi\wedge\delta\phi)\psi\rangle\Big\}.}$$

Here '$\dot{\phi}$' denotes the time derivative of ϕ and $\tilde{\Gamma}^0$ is the symmetric form (1.2) on spinors evaluated in the time direction.

We compute the Noether current of the supersymmetry transformation η^aQ_a in two ways. First, we work with superfields—so on $\mathcal{F}_\Phi\times M^{3|2}$—where supersymmetry is manifest. The supersymmetry transformation η^aQ_a induces the vector field

$$\xi = \eta^a\tau_{Q_a} + \widehat{\Xi} \tag{4.42}$$

on $\mathcal{F}_\Phi\times M^{3|2}$, where

$$\widehat{\Xi}\Phi = -\eta^c\tau_{Q_c}\Phi. \tag{4.43}$$

Then

$$J = \iota(\xi)(L_0 + L_1 + \gamma). \tag{4.44}$$

Alternatively, we can work in components—so on $\mathcal{F}_{(\phi,\psi,F)}\times\check{M}^3$—with nonmanifest supersymmetry and compute

$$\check{J} = \iota(\hat{\xi})\check{\gamma} - \alpha, \tag{4.45}$$

where α satisfies

$$\text{Lie}(\hat{\xi})\big(|d^3x|\ (\check{L}_0 + \check{L}_1)\big) = d\alpha \tag{4.46}$$

and may be computed directly from (1.42). Of course, we pass from J to $\check{J}$ by integration (1.32) over the odd variables.

As a first step in computing (4.44) we have

(4.47)
$$\begin{aligned}\iota(\xi)L_0 &= \iota(\eta^c\tau_{Q_c})L_0\\ &= \iota(\eta^c\partial_c + \eta^c\theta^d\partial_{cd})\,|d^3x|\,d^2\theta\ \frac{1}{4}\epsilon^{ab}\langle D_a\Phi, D_b\Phi\rangle\\ &= \iota(\partial_c)\,|d^3x|\,d^2\theta\ \frac{1}{4}\epsilon^{ab}\eta^c\langle D_a\Phi, D_b\Phi\rangle + \iota(\partial_{cd})\,|d^3x|\,d^2\theta\ \frac{1}{4}\epsilon^{ab}\eta^c\theta^d\langle D_a\Phi, D_b\Phi\rangle.\end{aligned}$$

Similarly,

$$\iota(\xi)L_1 = \iota(\partial_c)\,|d^3x|\,d^2\theta\;\eta^c\Phi^*(h) + \iota(\partial_{cd})\,|d^3x|\,d^2\theta\;\eta^c\theta^d\Phi^*(h). \tag{4.48}$$

From (4.32) we compute the $(0,|-1|)$ component of $\iota(\xi)\gamma$; the other components play no role.

$$\begin{aligned}\big(\iota(\xi)\gamma\big)^{0,|-1|} &= \iota(\widehat{\Xi})\,\iota(D_a)\,|d^3x|\,d^2\theta\;\frac{1}{2}\epsilon^{ab}\langle D_b\Phi,\delta\Phi\rangle\\ &= \iota(D_a)\,|d^3x|\,d^2\theta\;\Big\{\frac{1}{2}\epsilon^{ab}\langle D_b\Phi,-\eta^c\tau_{Q_c}\Phi\rangle\Big\}\\ &= \iota(\partial_a)\,|d^3x|\,d^2\theta\;\Big\{\frac{1}{2}\epsilon^{ab}\eta^c\langle D_b\Phi,\tau_{Q_c}\Phi\rangle\Big\}\\ &\qquad + \iota(\partial_{ad})\,|d^3x|\,d^2\theta\;\Big\{\frac{1}{2}\epsilon^{ab}\eta^c\theta^d\langle D_b\Phi,\tau_{Q_c}\Phi\rangle\Big\}.\end{aligned} \tag{4.49}$$

The $(0,|-1|)$ component of J is the sum of (4.47), (4.48), and (4.49).

Next, we integrate J over the odd variables using (2.40). The terms with $\iota(\partial_a)$ drop out; only the terms with $\iota(\partial_{ab})$ survive. From (4.47) we have

$$\begin{aligned}\iota(\partial_{cd})\,|d^3x|\;&\int d^2\theta\,\frac{1}{4}\epsilon^{ab}\eta^c\theta^d\langle D_a\Phi,D_b\Phi\rangle\\ &= \iota(\partial_{cd})\,|d^3x|\,\Big\{-\frac{1}{8}\epsilon^{ab}\epsilon^{ef}\eta^c i^* D_eD_f\theta^d\langle D_a\Phi,D_b\Phi\rangle\Big\}\\ &= \iota(\partial_{cd})\,|d^3x|\,\Big\{\frac{1}{2}\epsilon^{ab}\epsilon^{de}\eta^c i^*\langle D_eD_a\Phi,D_b\Phi\rangle\Big\}\\ &= \iota(\partial_{cd})\,|d^3x|\,\Big\{\epsilon^{ab}\epsilon^{de}\eta^c\langle-(\partial_{ae}\phi+\epsilon_{ea}F),\psi_b\rangle\Big\}\\ &= \iota(\partial_{cd})\,|d^3x|\,\Big\{-\frac{1}{2}\epsilon^{ab}\epsilon^{de}\eta^c\langle\partial_{ae}\phi,\psi_b\rangle+\frac{1}{2}\epsilon^{db}\eta^c\langle F,\psi_b\rangle\Big\}.\end{aligned} \tag{4.50}$$

From (4.48) we have

$$\begin{aligned}\iota(\partial_{cd})\,|d^3x|\;&\int d^2\theta\,\eta^c\theta^d\Phi^*(h)\\ &= \iota(\partial_{cd})\,|d^3x|\,\Big\{-\frac{1}{2}\epsilon^{ab}\eta^c i^* D_aD_b\theta^d\Phi^*(h)\Big\}\\ &= \iota(\partial_{cd})\,|d^3x|\,\Big\{-\epsilon^{ad}\eta^c i^*\langle\Phi^*\operatorname{grad}h,D_a\Phi\rangle\Big\}\\ &= \iota(\partial_{cd})\,|d^3x|\,\Big\{-\epsilon^{ad}\eta^c\langle\phi^*\operatorname{grad}h,\psi_a\rangle.\Big\}\end{aligned} \tag{4.51}$$

Finally, from (4.49) we compute

(4.52)
$$\begin{aligned}
&\iota(\partial_{cd})\,|d^3x|\int d^2\theta\,\frac{1}{2}\epsilon^{cb}\eta^a\theta^d\langle D_b\Phi,\tau_{Q_a}\Phi\rangle\\
&\qquad=\iota(\partial_{cd})\,|d^3x|\,\Big\{-\frac{1}{4}\epsilon^{cb}\epsilon^{ef}\eta^a i^*D_eD_f\theta^d\langle D_b\Phi,\tau_{Q_a}\Phi\rangle\Big\}\\
&\qquad=\iota(\partial_{cd})\,|d^3x|\,\Big\{-\frac{1}{2}\epsilon^{cb}\epsilon^{ef}\eta^a i^*\big[\langle D_eD_b\Phi,\tau_{Q_a}\Phi\rangle-\langle D_b\Phi,D_e\tau_{Q_a}\Phi\rangle\big]\Big\}\\
&\qquad=\iota(\partial_{cd})\,|d^3x|\,\Big\{-\frac{1}{2}\epsilon^{cb}\epsilon^{ed}\eta^a i^*\big[\langle D_eD_b\Phi,D_a\Phi\rangle+\langle D_b\Phi,D_aD_e\Phi\rangle\big]\Big\}\\
&\qquad=\iota(\partial_{cd})\,|d^3x|\,\Big\{-\frac{1}{2}\epsilon^{ed}(\epsilon^{cb}\eta^a+\epsilon^{ac}\eta^b)i^*\langle D_eD_b\Phi,D_a\Phi\rangle+\epsilon^{cb}\epsilon^{ed}\eta^a i^*\langle\partial_{ea}\Phi,D_b\Phi\rangle\Big\}\\
&\qquad=\iota(\partial_{cd})\,|d^3x|\,\Big\{\frac{1}{2}\epsilon^{ed}\epsilon^{ab}\eta^c\langle\partial_{eb}\phi+\epsilon_{ab}F,\psi_a\rangle+\epsilon^{cb}\epsilon^{ed}\eta^a i^*\langle\partial_{ea}\phi,\psi_b\rangle\Big\}.
\end{aligned}$$

Summing (4.50), (4.51), and (4.52) we find after using (4.22) that

$$\check{J}=\iota(\partial_{cd})\,|d^3x|\,\Big\{\epsilon^{bc}\epsilon^{de}\eta^a\langle\partial_{ae}\phi,\psi_b\rangle+\epsilon^{db}\eta^c\langle\phi^*\operatorname{grad}h,\psi_b\rangle\Big\}. \tag{4.53}$$

The supercurrent j_a, which is *minus* the Noether current of the supersymmetry transformation Q_a, is

$$\boxed{j_a=\iota(\partial_{cd})\,|d^3x|\,\Big\{\epsilon^{cb}\epsilon^{de}\langle\partial_{ae}\phi,\psi_b\rangle+\epsilon^{bd}\delta^c_a\langle\phi^*\operatorname{grad}h,\psi_b\rangle\Big\}.} \tag{4.54}$$

As a check, we compute $\check{J}$ directly using (4.45). Using (4.39) we compute the first term of (4.45):

(4.55)
$$\begin{aligned}
\iota(\hat{\xi})\check{\gamma}&=\frac{1}{2}\epsilon^{ac}\epsilon^{bd}\iota(\partial_{cd})\,|d^3x|\,\Big\{\langle\partial_{ab}\phi,-\eta^e\psi_e\rangle+\langle\psi_b,\eta^e(\partial_{ae}\phi-\epsilon_{ae}F)\rangle\Big\}\\
&=\iota(\partial_{cd})\,|d^3x|\,\Big\{\frac{1}{2}\epsilon^{ac}(\epsilon^{db}\eta^e+\epsilon^{de}\eta^b)\langle\partial_{ab}\phi,\psi_e\rangle-\frac{1}{2}\epsilon^{ec}\eta^d\langle\phi^*\operatorname{grad}h,\psi_e\rangle\Big\}.
\end{aligned}$$

Note we use (4.22). Next, we compute α directly from (1.42), setting $\Delta=0$:

(4.56)
$$\begin{aligned}
\alpha&=\iota(\partial_\mu)\,|d^3x|\,\Gamma^\mu_{ab}(-\frac{1}{2}\epsilon^{ef})i^*D_eD_f\eta^a\theta^b\big[\frac{1}{4}\epsilon^{gh}\langle D_g\Phi,D_h\Phi\rangle+\Phi^*(h)\big]\\
&=\iota(\partial_{cd})\,|d^3x|\,\Big\{\frac{1}{2}\epsilon^{ef}\eta^c i^*D_eD_f\theta^d\big[\frac{1}{4}\epsilon^{ab}\langle D_a\Phi,D_b\Phi\rangle+\Phi^*(h)\big]\Big\}\\
&=\iota(\partial_{cd})\,|d^3x|\,\Big\{\frac{1}{2}\epsilon^{ed}\eta^c i^*\big[\epsilon^{ab}\langle D_eD_a\Phi,D_b\Phi\rangle+2\langle\Phi^*\operatorname{grad}h,D_e\Phi\rangle\big]\Big\}\\
&=-\iota(\partial_{cd})\,|d^3x|\,\Big\{\frac{1}{2}\epsilon^{ab}\epsilon^{ed}\eta^c\langle\partial_{ae}\phi,\psi_b\rangle-\frac{1}{2}\epsilon^{ec}\eta^d\langle\phi^*\operatorname{grad}h,\psi_e\rangle\big]\Big\}.
\end{aligned}$$

The first term agrees with (3.29). Combining (4.55) and (4.56) we recover (4.53).

§4.5. Reduction to $M^{2|(1,1)}$

We can reduce the model to $n = 2$ dimensions either in superspace or in components. In both cases we consider the $n = 3$ dimensional theory with fields constrained to be invariant under ∂_{12}. All the formulas worked out in $n = 3$ dimensions hold, but there are simplifications. We already wrote in (4.15) the supersymmetry transformations. We record here the supercharges $\tilde{Q}_+, \tilde{Q}_-$ obtained by integrating the Noether current (4.54) over space:

$$\begin{aligned} \tilde{Q}_+ &= \int_{x^0=\text{const}} \Big\{ \langle \partial_+\phi, \psi_+ \rangle - \langle \phi^* \operatorname{grad} h, \psi_- \rangle \Big\} \, |dx^1| \\ \tilde{Q}_- &= \int_{x^0=\text{const}} \Big\{ \langle \partial_-\phi, \psi_- \rangle + \langle \phi^* \operatorname{grad} h, \psi_+ \rangle \Big\} \, |dx^1|. \end{aligned} \tag{4.57}$$

In the abstract supersymmetry algebra (1.15) the Lie bracket $[Q_+, Q_-]$ of the supersymmetry generators vanishes. However, the Poisson bracket $\{\tilde{Q}_+, \tilde{Q}_-\}$ of the Noether charges is nonzero in this case; it is a locally constant function on the space of classical solutions. We compute it by acting $\widehat{Q}_+$ on $\tilde{Q}_-$, where $\widehat{Q}_+$ is the Hamiltonian vector field which corresponds to the function $\tilde{Q}_+$; its action on fields is given in (4.15):

$$\begin{aligned} \widehat{Q}_+\phi &= -\psi_+ \\ \widehat{Q}_+\psi_+ &= \partial_+\phi \\ \widehat{Q}_+\psi_- &= -\phi^* \operatorname{grad} h \\ \widehat{Q}_+(\phi^* \operatorname{grad} h) &= \nabla_+\psi_- + R(\psi_+, \psi_-)\psi_+. \end{aligned} \tag{4.58}$$

In the following computation most terms drop out after we use the equation of motion (4.34) and the Bianchi identity:

(4.59)

$$\begin{aligned} \frac{1}{2}\{\tilde{Q}_+, \tilde{Q}_-\} &= \frac{1}{2} \int_{x^0=\text{const}} \Big\{ -\langle \partial_-\phi, \phi^* \operatorname{grad} h \rangle + \langle \partial_+\phi, \phi^* \operatorname{grad} h \rangle \Big\} \, |dx^1| \\ &= \int_{x^0=\text{const}} \langle \partial_1\phi, \phi^* \operatorname{grad} h \rangle \, |dx^1| \\ &= \int_{x^0=\text{const}} \partial_1(\phi^* h) \, |dx^1| \\ &= h\big(\phi(\infty)\big) - h\big(\phi(-\infty)\big). \end{aligned}$$

A scalar field $\phi = \phi(x^1)$ depending on the spatial variable has finite energy only if it has limits as $x^1 \to \pm\infty$ which are critical points of h. The Poisson bracket (4.59) measures the difference of the critical values. It induces a central extension of the abstract supersymmetry algebra.

CHAPTER 5
The Supersymmetric σ-Model in Dimension 4

In this chapter we consider the σ-model with 4 supersymmetries, which was described in components in Chapter 3. Here we give a manifestly supersymmetric treatment in superspace. In many respects the treatment parallels that of Chapter 4, so we are briefer here. (One new feature over theories with 2 supersymmetries is the presence of an R-symmetry.) Although the model is well-defined and supersymmetric for any Kähler target manifold X, the superspace model is nicest when X is Hodge. In that case we give a global superspace lagrangian (see (5.22)). In these theories the scalar potential is the norm square of the gradient of a holomorphic function called the *superpotential*. We describe it in §5.3, and in particular prove Theorem 3.82(ii). We do not give a detailed analysis of the classical theory, but simply summarize the most important equations in §5.4.

§5.1. Fields and supersymmetry transformations on $M^{4|4}$

Linear Case. The simplest superfield for which we can write sensible lagrangians is the (linear) *chiral superfield*

$$\Phi\colon M^{4|4} \longrightarrow \mathbb{C}, \tag{5.1}$$

which is required to satisfy

$$\overline{D}_{\dot{a}}\Phi = 0. \tag{5.2}$$

The complex conjugate of a chiral superfield is called an *antichiral multiplet*. Define the component fields as

$$\boxed{\begin{aligned} \phi &= i^*\Phi \\ \psi_a &= \frac{1}{\sqrt{2}} i^* D_a \Phi \\ F &= \frac{1}{2} i^*(-D^2)\Phi. \end{aligned}} \tag{5.3}$$

Component fields for the antichiral superfield $\overline{\Phi}$ are defined by the conjugate equations to (5.3). Note that F is a *complex* scalar field.

Let $\hat{\xi}$ be the vector field on the space $\mathcal{F}_{(\phi,\psi,F)}$ of component fields induced by the supersymmetry transformation $\eta^a Q_a + \bar{\eta}^{\dot{a}}\overline{Q}_{\dot{a}}$. Here $\eta^a, \bar{\eta}^{\dot{a}}$ are independent odd parameters. The action on the components of Φ is computed by similar manipulations as in (4.3)–(4.5). For example, using (2.73) and (2.75) we compute

$$\begin{aligned}\hat{\xi}F &= -i^*(\eta^a D_a + \bar{\eta}^{\dot{a}}\overline{D}_{\dot{a}})(\frac{1}{2}D^2)\Phi \\ &= -\bar{\eta}^{\dot{a}}\epsilon^{bc}i^*\partial_{b\dot{a}}D_c\Phi \\ &= -\sqrt{2}\bar{\eta}^{\dot{a}}\epsilon^{bc}\partial_{b\dot{a}}\psi_c \\ &= \sqrt{2}\bar{\eta}^{\dot{a}}(\not{D}\psi)_{\dot{a}}.\end{aligned} \tag{5.4}$$

Collecting the results of similar computations, we display the supersymmetry transformation laws:

$$\boxed{\begin{aligned}\hat{\xi}\phi &= -\sqrt{2}\eta^a\psi_a \\ \hat{\xi}\psi_a &= \sqrt{2}(\bar{\eta}^{\dot{b}}\partial_{a\dot{b}}\phi - \eta^b\epsilon_{ab}F) \\ \hat{\xi}F &= \sqrt{2}\bar{\eta}^{\dot{a}}(\not{D}\psi)_{\dot{a}}.\end{aligned}} \tag{5.5}$$

The transformation laws for the components of $\overline{\Phi}$ are obtained by conjugating the equations in (5.5).

Nonlinear Case. The nonlinear chiral superfield is a map

$$\Phi\colon M^{4|4} \longrightarrow X, \tag{5.6}$$

where X is a complex manifold and Φ is required to pullback holomorphic functions on X to linear chiral superfields on $M^{4|4}$. Another way to state this is that $D_a\Phi$ is a vector field of type $(1,0)$, and so $\overline{D}_{\dot{a}}\overline{\Phi}$ is a vector field of type $(0,1)$. Again the composition of a chiral superfield and a holomorphic function is a chiral superfield. The conjugate conditions define a nonlinear antichiral superfield. To define the component fields we assume that X is Kähler[31] and use the Levi-Civita covariant derivative as in (4.7):

$$\boxed{\begin{aligned}\phi &= i^*\Phi \\ \psi_a &= \frac{1}{\sqrt{2}}i^*D_a\Phi \\ F &= \frac{1}{2}i^*(-\mathcal{D}^2)\Phi.\end{aligned}} \tag{5.7}$$

Note that ψ_a and F have type $(1,0)$. The conjugate fields $\bar{\psi}_{\dot{a}}$ and $\overline{F}$ have type $(0,1)$.

Now the analogs of equations (2.74) and (2.75) for covariant derivatives of nonlinear chiral superfields are easily derived:

$$\mathcal{D}_a\overline{\mathcal{D}}^2\Phi = \overline{\mathcal{D}}^2 D_a\Phi = 0, \tag{5.8}$$

$$\overline{\mathcal{D}}_{\dot{a}}\mathcal{D}^2\Phi = -2\epsilon^{bc}\Big\{\nabla_{b\dot{a}} - \frac{1}{4}R(D_b\Phi, \overline{D}_{\dot{a}}\overline{\Phi})\Big\}D_c\Phi. \tag{5.9}$$

[31] In any case X must be Kähler to define the σ-model action.

The conjugate equations to (5.8) and (5.9) give information about covariant derivatives of nonlinear antichiral superfields. There is only one curvature term which alters (5.5) in the nonlinear case:

$$\boxed{\begin{aligned}\hat{\xi}\phi &= -\sqrt{2}\eta^a\psi_a \\ \hat{\xi}\psi_a &= \sqrt{2}(\bar{\eta}^{\dot{b}}\partial_{a\dot{b}}\phi - \eta^b\epsilon_{ab}F) \\ \hat{\xi}F &= \sqrt{2}\bar{\eta}^{\dot{a}}\big[(\not{D}\psi)_{\dot{a}} + \frac{1}{4}\epsilon^{bc}R(\psi_b,\bar{\psi}_{\dot{a}})\psi_c\big].\end{aligned}} \tag{5.10}$$

§5.2. The σ-model action on $M^{4|4}$

Linear Case. Here it is worth computing separately for the linear superfield (5.1). The lagrangian density on $M^{4|4}$ is

$$\mathcal{L}_0 = |d^4x|\, d^4\theta\, \frac{1}{4}\overline{\Phi}\Phi. \tag{5.11}$$

Recall from the discussion following (2.72) that we compute component lagrangians by integrating over the θ's and adding the wave operator. Hence by (2.72)

(5.12)

$$\begin{aligned}4\check{\mathcal{L}}_0 &= \left\{\int d^4\theta + \Box i^*\right\}\overline{\Phi}\Phi \\ &= \frac{1}{2}i^*(D^2\overline{D}^2 + \overline{D}^2D^2 + 4\Box)\overline{\Phi}\Phi \\ &= \frac{1}{2}i^*\big\{D^2(\overline{D}^2\overline{\Phi}\cdot\Phi) + \overline{D}^2(\overline{\Phi}\cdot D^2\Phi) + 4\Box(\overline{\Phi}\Phi)\big\} \\ &= \frac{1}{2}i^*\big\{(D^2\overline{D}^2 + 4\Box)\overline{\Phi}\cdot\Phi + \overline{\Phi}\cdot(\overline{D}^2D^2 + 4\Box)\Phi + 8\langle d\overline{\Phi}, d\Phi\rangle \\ &\qquad + 2\,\overline{D}^2\overline{\Phi}\cdot D^2\Phi + \epsilon^{ab}D_a\overline{D}^2\overline{\Phi}\cdot D_b\Phi - \epsilon^{\dot{a}\dot{b}}\overline{D}_{\dot{b}}\overline{\Phi}\cdot\overline{D}_{\dot{a}}D^2\Phi\big\} \\ &= \frac{1}{2}\big\{8\langle d\bar{\phi}, d\phi\rangle + 8\overline{F}F + 4\epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\psi_b\partial_{a\dot{c}}\bar{\psi}_{\dot{d}} + 4\epsilon^{\dot{a}\dot{b}}\epsilon^{cd}\bar{\psi}_{\dot{b}}\partial_{c\dot{a}}\psi_d\big\},\end{aligned}$$

where we use (2.74)–(2.77) and the definition (5.7) of component fields in the last step. Finally, by (2.56)

$$\boxed{\check{\mathcal{L}}_0 = \langle d\bar{\phi}, d\phi\rangle + \frac{1}{2}(\bar{\psi}\not{D}\psi + \psi\not{D}\bar{\psi}) + \overline{F}F.} \tag{5.13}$$

The Dirac form in (5.13) is real, but up to an exact term we can write it as the complex form $\bar{\psi}\not{D}\psi$:

$$\frac{1}{2}(\bar{\psi}\not{D}\psi + \psi\not{D}\bar{\psi}) = \bar{\psi}\not{D}\psi - \frac{1}{2}\partial_\mu(\bar{\psi}\Gamma^\mu\psi). \tag{5.14}$$

We drop this exact term in subsequent formulas. As in 3 dimensions, F enters the lagrangian (5.13) as an auxiliary field and can be eliminated through its equation of motion $F = 0$.

Nonlinear Case. Now we carry out the same computation for the nonlinear chiral superfield (5.6) with values in a Kähler manifold X. The superfield expression for the lagrangian depends on the choice of a local Kähler potential K, which is a locally defined real-valued function on X such that the Kähler form is

$$\omega = \sqrt{-1}\,\partial\bar{\partial}K \tag{5.15}$$

and the metric is

$$\langle \cdot, \cdot \rangle = 2\sqrt{-1}\,\omega = 2\bar{\partial}\partial K, \tag{5.16}$$

viewed as a *bilinear* form on $\overline{TX} \otimes TX$. For example, on $\mathbb{C}$ we have a global Kähler potential $K(\bar{z}, z) = \frac{1}{2}\|z\|^2$. We use double bar notation $\|\cdot\|$ throughout for the norm on a hermitian vector space.

The lagrangian density on $M^{4|4}$ is

$$\boxed{\mathcal{L}_0 = |d^4x|\, d^4\theta\, \frac{1}{2} K(\overline{\Phi}, \Phi).} \tag{5.17}$$

Although it depends on a choice of Kähler potential, the component lagrangian makes sense globally by the following argument. Consider a change $K \to K + f + \bar{f}$ for f a holomorphic function on X. Of course, it does not affect the Kähler form (5.15) and is exactly the ambiguity in the choice of Kähler potential. Now since we compute the component lagrangian using the first term of (2.71)—the integral over θ plus the wave operator—and since that operation annihilates chiral and antichiral superfields, such a change in K does not affect the component lagrangian.

There is a global formulation of the lagrangian density $\mathcal{L}_0$ on superspace when X is a *Hodge* manifold. A Hodge manifold is a Kähler manifold X together with a holomorphic hermitian line bundle $L_\omega \to X$ whose curvature is related to the Kähler form by

$$\operatorname{curv}(L_\omega) = \sqrt{-1}\,\omega. \tag{5.18}$$

Recall that if s is a local holomorphic section of $L_\omega \to X$, then

$$\operatorname{curv}(L_\omega) = \bar{\partial}\partial \log \|s\|^2, \tag{5.19}$$

and so $\log \|s\|^2$ is a local Kähler potential. Consider a chiral superfield $\Phi\colon M^{4|4} \to X$. Since Minkowski space is contractible, there is a *global* lift to a chiral superfield[32]

$$\tilde{\Phi}\colon M^{4|4} \longrightarrow \Phi^* L_\omega \tag{5.21}$$

which is a section of $\Phi^* L_\omega \to M^{4|4}$. For any such lift we write the lagrangian (5.17) globally as

$$\boxed{\mathcal{L}_0 = |d^4x|\, d^4\theta\, \frac{1}{2} \log \|\tilde{\Phi}\|^2.} \tag{5.22}$$

[32]The existence of a global lift depends on the fact that

$$H^1(M^{4|4}, \mathcal{C}^*) = 0, \tag{5.20}$$

where $\mathcal{C}^*$ is the sheaf of nonzero chiral superfields. The vanishing (5.20) is obvious in components.

By the argument in the previous paragraph the corresponding component lagrangian is independent of the choice of lift $\tilde{\Phi}$. We remark that it is not unfamiliar to have an action whose lagrangian density depends on an auxiliary map: consider, for example, the usual formulation of the Wess-Zumino-Witten term in two or four dimensions.

As a preliminary to computing the component lagrangian from (5.17), note that the Cartan formula $\mathrm{Lie}(\eta) = d\iota(\eta) + \iota(\eta)d$ for the Lie derivative, valid for all vector fields η, implies the following equations when acting on 1-forms:

$$\iota(\zeta)\iota(\zeta')d = -\mathrm{Lie}(\zeta)\iota(\zeta') - \mathrm{Lie}(\zeta')\iota(\zeta) + \iota([\zeta,\zeta']), \qquad \zeta,\zeta' \text{ odd}; \tag{5.23}$$

$$\iota(\zeta)\iota(\xi)d = \mathrm{Lie}(\xi)\iota(\zeta) - \mathrm{Lie}(\zeta)\iota(\xi) + \iota([\zeta,\xi]), \qquad \zeta \text{ odd}, \ \xi \text{ even}. \tag{5.24}$$

We use these formulas several times in the next computation.

We choose (2.72) to compute the component lagrangian, though (2.71) also leads to a straightforward computation. Thus we expand (5.17):

(5.25)

$$\begin{aligned}
\overline{\mathcal{D}}^2\mathcal{D}^2\Phi^*K &= \frac{1}{2}\epsilon^{ab}\overline{\mathcal{D}}^2 D_a\iota(D_b\Phi)\partial K \\
&= \frac{1}{4}\epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\overline{D}_{\dot{c}}(-\mathcal{D}_a\overline{D}_{\dot{d}} - 2\partial_{a\dot{d}})\iota(D_b\Phi)\partial K \\
&= \frac{1}{4}\epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\overline{D}_{\dot{c}}\Big\{D_a\iota(\overline{D}_{\dot{d}}\overline{\Phi})\iota(D_b\Phi)\bar{\partial}\partial K + 2D_a\iota(\partial_{b\dot{d}}\Phi)\partial K \\
&\qquad\qquad - 2\iota(D_b\Phi)\iota(\partial_{a\dot{d}}\Phi)\bar{\partial}\partial K - 2D_b\iota(\partial_{a\dot{d}}\Phi)\partial K\Big\} \\
&= \frac{1}{4}\epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\overline{D}_{\dot{c}}\Big\{-\frac{1}{2}D_a\langle\overline{D}_{\dot{d}}\overline{\Phi}, D_b\Phi\rangle - \langle\partial_{a\dot{d}}\overline{\Phi}, D_b\Phi\rangle + 4D_a\iota(\partial_{b\dot{d}}\Phi)\partial K\Big\} \\
&= \frac{1}{4}\epsilon^{\dot{c}\dot{d}}\overline{D}_{\dot{c}}\langle\overline{D}_{\dot{d}}\overline{\Phi}, \mathcal{D}^2\Phi\rangle + \epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\overline{D}_{\dot{c}}D_a\iota(\partial_{b\dot{d}}\Phi)\partial K.
\end{aligned}$$

The first term is

(5.26)

$$\begin{aligned}
\frac{1}{4}\epsilon^{\dot{c}\dot{d}}\overline{D}_{\dot{c}}\langle\overline{D}_{\dot{d}}\overline{\Phi}, \mathcal{D}^2\Phi\rangle &= \frac{1}{2}\langle\overline{\mathcal{D}}^2\overline{\Phi}, \mathcal{D}^2\Phi\rangle - \frac{1}{4}\epsilon^{\dot{c}\dot{d}}\langle\overline{D}_{\dot{d}}\overline{\Phi}, \overline{D}_{\dot{c}}\mathcal{D}^2\Phi\rangle \\
&= \frac{1}{2}\epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\Big\{\langle\overline{D}_{\dot{d}}\overline{\Phi}, \nabla_{a\dot{c}}D_b\Phi\rangle \\
&\qquad - \frac{1}{4}\langle\overline{D}_{\dot{d}}\overline{\Phi}, R(D_a\Phi, \overline{D}_{\dot{c}}\overline{\Phi})D_b\Phi\rangle\Big\} + \frac{1}{2}\langle\overline{\mathcal{D}}^2\overline{\Phi}, \mathcal{D}^2\Phi\rangle.
\end{aligned}$$

The second term in (5.25) is

(5.27)

$$\begin{aligned}
\epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\overline{D}_{\dot{c}}D_a\iota(\partial_{b\dot{d}}\Phi)\partial K &= \epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\Big\{-\mathcal{D}_a\overline{D}_{\dot{c}}\iota(\partial_{b\dot{d}}\Phi)\partial K - 2\partial_{a\dot{c}}\iota(\partial_{b\dot{d}}\Phi)\partial K\Big\} \\
&= \epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\Big\{-\frac{1}{2}D_a\langle\overline{D}_{\dot{c}}\overline{\Phi}, \partial_{b\dot{d}}\Phi\rangle - 2\partial_{a\dot{c}}\iota(\partial_{b\dot{d}}\Phi)\partial K\Big\} \\
&= \epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\Big\{\frac{1}{2}\langle\overline{D}_{\dot{c}}\overline{\Phi}, \nabla_{b\dot{d}}D_a\Phi\rangle + \langle\partial_{a\dot{c}}\overline{\Phi}, \partial_{b\dot{d}}\Phi\rangle \\
&\qquad\qquad - 2\partial_{a\dot{c}}\iota(\partial_{b\dot{d}}\Phi)\partial K\Big\}.
\end{aligned}$$

Combining (5.25)–(5.27) we have

$$\begin{aligned}(5.28)\quad &\frac{1}{2}\big(\overline{\mathcal{D}}^2\mathcal{D}^2+2\epsilon^{ab}\epsilon^{\dot c\dot d}\partial_{a\dot c}\iota(\partial_{b\dot d}\Phi)\partial K\big)(\frac{1}{2}\Phi^*K)\\ &\qquad=\epsilon^{ab}\epsilon^{\dot c\dot d}\Big\{\frac{1}{4}\langle\partial_{b\dot d}\overline{\Phi},\partial_{a\dot c}\Phi\rangle+\frac{1}{4}\langle\overline{D}_{\dot d}\overline{\Phi},\nabla_{a\dot c}D_b\Phi\rangle\\ &\qquad\qquad-\frac{1}{32}\langle\overline{D}_{\dot d}\overline{\Phi},R(D_a\Phi,\overline{D}_{\dot c}\overline{\Phi})D_b\Phi\rangle\Big\}+\frac{1}{8}\langle\overline{\mathcal{D}}^2\overline{\Phi},\mathcal{D}^2\Phi\rangle.\end{aligned}$$

Now we add (5.28) to its complex conjugate. Since

$$(5.29)\qquad 2\epsilon^{ab}\epsilon^{\dot c\dot d}\partial_{a\dot c}\iota(\partial_{b\dot d}\Phi)(\partial K+\bar\partial K)=2\epsilon^{ab}\epsilon^{\dot c\dot d}\partial_{a\dot c}\partial_{b\dot d}(\Phi^*K)=4\Box(\Phi^*K),$$

the left hand side restricted to $\check M^4$ is the component lagrangian (as in the second line of (5.12)), and so finally

$$(5.30)\qquad \boxed{\check{\mathcal{L}}_0=\langle\overline{d\phi},d\phi\rangle+\langle\bar\psi\not{D}_\phi\psi\rangle-\frac{1}{4}\epsilon^{ab}\epsilon^{\dot c\dot d}\langle\bar\psi_{\dot d},\phi^*R(\psi_a,\bar\psi_{\dot c})\psi_b\rangle+\langle\overline{F},F\rangle.}$$

Recall that we have absorbed an exact term (5.14) in writing the Dirac form. With this understood, the lagrangian (5.30) agrees with (3.48) after eliminating the auxiliary field.

We check in Problem FP10(c) of [I-**Homework**] that the lagrangian (5.17) on $M^{4|4}$ reduces to (4.16) in $M^{3|2}$.

A new aspect in four dimensions is the $U(1)$ R-symmetry (2.81). We claim that the σ-model lagrangian is invariant under R. First, in the superspace expression (5.17) (or (5.22)) we note that $d^4\theta=d^2\theta\,d^2\bar\theta$ is invariant. So if R acts trivially on Φ, then $\mathcal{L}_0$ is obviously invariant. When R acts nontrivially it is easier to argue in components. Suppose for simplicity that Φ takes values in $\mathbb{C}$, and let R acts on Φ with weight k. Then

$$(5.31)\qquad \boxed{\begin{array}{ll}R\phi=k\phi, & R\bar\phi=-k\bar\phi\\ R\psi=(k+1)\psi, & R\bar\psi=-(k+1)\bar\psi\\ RF=(k+1)F, & R\overline{F}=-(k+2)\overline{F}.\end{array}}$$

Direct inspection of (5.30) shows that $\check{\mathcal{L}}_0$ is R-invariant.

§5.3. The superpotential term on $M^{4|4}$

As a preliminary, note that if Λ is any chiral superfield, then $|d^4x|\int d^2\theta\,\Lambda$ makes sense as a density on Minkowski space. Similarly, for $\overline{\Lambda}$ antichiral we can define $|d^4x|\int d^2\bar\theta\,\overline{\Lambda}$. One view is that $M^{4|4}$ is a split cs-manifold with a complex two dimensional odd tangent bundle. In fact, there are two such cs-manifolds, one for each of the complex structures on the real four dimensional spin representation (see (2.2)). These cs-manifolds have canonical densities—denoted $|d^4x|\,d^2\theta$ (resp. $|d^4x|\,d^2\bar\theta$)—and global functions on these cs-manifolds are chiral (resp. antichiral) superfields. To compute these chiral integrals we use (2.40).

The distinction between terms in the action of the form $\int|d^4x|\,d^4\theta$ and $\int|d^4x|\,d^2\theta$ is important in $M^{4|4}$ and also in $M^{2|(2,2)}$.

We introduce a superpotential term, and we may as well work in the nonlinear σ-model. Let $W\colon X \to \mathbb{C}$ be a *holomorphic* function, and set

$$\boxed{\mathcal{L}_1 = |d^4x|\ \mathrm{Re}\Big\{d^2\theta\ \Phi^*(W)\Big\}.} \tag{5.32}$$

It lives in the direct sum of the spaces of chiral and antichiral densities. The superpotential term in the component action is the integral of $\mathcal{L}_1$:

$$\int\mathcal{L}_1 = \int|d^4x|\ \frac{1}{2}\left\{\int d^2\theta\ \Phi^*(W)\ +\ \int d^2\bar\theta\ \Phi^*(\overline{W})\right\} = \int|d^4x|\ \check{\mathcal{L}}_1. \tag{5.33}$$

We compute $\check{\mathcal{L}}_1$ exactly as in (4.21); recalling our normalizations (5.7) for the component fields we obtain

$$\check{\mathcal{L}}_1 = \iota(F)\phi^*\partial W + \iota(\overline{F})\phi^*\overline{\partial W} - \frac{1}{2}\epsilon^{ab}\phi^*\nabla\partial W(\psi_a,\psi_b) - \frac{1}{2}\epsilon^{\dot a\dot b}\phi^*\nabla\overline{\partial W}(\bar\psi_{\dot a},\bar\psi_{\dot b}). \tag{5.34}$$

We eliminate the auxiliary fields $F, \overline{F}$ from $\check{\mathcal{L}}_0 + \check{\mathcal{L}}_1$ using their equations of motion

$$\boxed{\begin{aligned} F &= -\phi^*\operatorname{grad} W \\ \overline{F} &= -\phi^*\overline{\operatorname{grad} W}, \end{aligned}} \tag{5.35}$$

where now $\overline{\operatorname{grad} W}$ is the vector field of type $(0,1)$ corresponding to the $(1,0)$-form ∂W using the hermitian metric on X. The total lagrangian with auxiliary fields eliminated is then

$$\boxed{\begin{aligned} \check{\mathcal{L}}_0 + \check{\mathcal{L}}_1 \sim\ & \langle\overline{d\phi}, d\phi\rangle + \langle\bar\psi \not{D}_\phi\psi\rangle - \frac{1}{4}\epsilon^{ab}\epsilon^{\dot c\dot d}\langle\bar\psi_{\dot d}, \phi^*R(\psi_a,\bar\psi_{\dot c})\psi_b\rangle \\ & - \phi^*\|\operatorname{grad} W\|^2 - \Big(\frac{1}{2}\epsilon^{ab}\phi^*\nabla\partial W(\psi_a,\psi_b) - \frac{1}{2}\epsilon^{\dot a\dot b}\phi^*\nabla\overline{\partial W}(\bar\psi_{\dot a},\bar\psi_{\dot b})\Big). \end{aligned}} \tag{5.36}$$

This is the lagrangian (3.85), and since the superspace model is manifestly supersymmetric we have proved Theorem 3.82(ii). Of course, the potential energy in (5.36) is

$$V = \|\operatorname{grad} W\|^2. \tag{5.37}$$

Other aspects of the discussion following (4.23) carry over directly.

Finally, we consider the R-symmetry (2.81). We already noted that it leaves $\mathcal{L}_0$ and $\check{\mathcal{L}}_0$ invariant, even if Φ is assigned a nonzero weight. On the other hand, $d^2\theta$ has weight 2 and so if $\mathcal{L}_1$ is to be invariant we must be able to assign Φ a weight so that $\Phi^*(W)$ has weight -2. (In general, there are several Φ_i with different weights.) Such a W is termed *quasi-homogeneous* in the physics literature.

§5.4. Analysis of the classical theory

We do not give details, having done so exhaustively in §4 for the σ-model on $M^{3|2}$. Rather, we simply record the formulas

$$\boxed{\check\gamma = \iota(\partial_{c\dot d})\,|d^4x|\ \frac{1}{\sqrt{2}}\epsilon^{ac}\epsilon^{\dot b\dot d}\big[\langle\partial_{a\dot b}\phi, \delta\bar\phi\rangle + \langle\partial_{a\dot b}\bar\phi, \delta\phi\rangle + \langle\psi_a, \delta\bar\psi_{\dot b}\rangle + \langle\bar\psi_{\dot b}, \delta\psi_a\rangle\big]} \tag{5.38}$$

for the variational 1-form,

$$\boxed{\check{\omega} = \iota(\partial_{c\dot{d}})\,|d^4x|\;\frac{1}{\sqrt{2}}\epsilon^{ac}\epsilon^{\dot{b}\dot{d}}\big[\langle\partial_{a\dot{b}}\delta\phi\wedge\delta\bar{\phi}\rangle + \langle\partial_{a\dot{b}}\delta\bar{\phi}\wedge\delta\phi\rangle + \langle\delta\psi_a\wedge\delta\bar{\psi}_{\dot{b}}\rangle + \langle\delta\bar{\psi}_{\dot{b}}\wedge\delta\psi_a\rangle\big]} \tag{5.39}$$

for the local symplectic form, and

$$\boxed{j_a = \iota(\partial_{c\dot{d}})\,|d^4x|\;\Big\{\sqrt{2}\epsilon^{cb}\epsilon^{\dot{d}\dot{e}}\langle\partial_{a\dot{e}}\phi,\psi_b\rangle + \sqrt{2}\,\epsilon^{\dot{b}\dot{d}}\delta^c_a\langle\phi^*\,\mathrm{grad}\,W,\bar{\psi}_{\dot{b}}\rangle\Big\}} \tag{5.40}$$

for the supercurrent corresponding to the supersymmetry transformation Q_a.

CHAPTER 6
Supersymmetric Yang-Mills Theories

Now we consider theories with gauge fields (connections). In the bosonic case there is a Yang-Mills theory in any dimension. The supersymmetric extension involves a *dual* spinor field as well, and it exists in dimensions $n = 3, 4, 6, 10$ with minimal supersymmetry. Our treatment in §6.1 is based on a special property of spinors which holds only in these cases: the quadratic form $s^* \mapsto \Gamma(s^*, s^*)$ on S^* takes values on the light cone. In §6.2 we describe the supersymmetric extensions of the most general bosonic theory, which is variously called a *gauge theory with matter* or a *gauged σ-model*. We consider theories in dimensions $n = 3, 4, 6$ with minimal supersymmetry. (The only theory in $n = 10$ is pure Yang-Mills.) The basic lagrangians, supersymmetry transformations, and supercurrents are summarized in Theorem 6.33. There are new terms which appear neither in the pure σ-model nor in the pure Yang-Mills theory. There is a superspace formulation of supersymmetric gauge theories in dimensions $n = 3, 4, 6$ (with minimal supersymmetry); we describe some common features in §6.3. The condition on the quadratic form has a geometric significance here as well. Details of the superspace formulation occupy the remaining chapters.

§6.1. The minimal theory in components

The bosonic Yang-Mills theory is determined by the data:

(6.1)
$$\begin{array}{ll} G & \text{compact Lie group} \\ \langle \cdot, \cdot \rangle & \text{bi-invariant scalar product on } \mathfrak{g} \end{array}$$

The lagrangian in n-dimensional Minkowski space is

(6.2)
$$L = -\frac{1}{2}|F_A|^2 \, |d^n x|,$$

where A is a connection on a principal G bundle P. The minimal supersymmetric extension has a single dual spinor field λ with values in the adjoint bundle and lagrangian

(6.3)
$$\boxed{L = \{-\frac{1}{2}|F_A|^2 + \frac{1}{2}\langle \lambda \not{D}_A \lambda \rangle\} \, |d^n x|.}$$

We ask: When is (6.3) supersymmetric? That is, in which dimensions and for what type of spinor field λ is (6.3) supersymmetric? Consider first the abelian theory, which is free. Suppose the spinor field has s components. Upon quantization,[33] the gauge field A has $n-2$ physical degrees of freedom and the spinor field has $s/2$ physical degrees of freedom. With one exception (explained presently) the number of bosonic and fermionic degrees of freedom in a supersymmetric theory must be equal, from which $n = 2 + s/2$. This is satisfied only for $n = 3, 4, 6, 10$ with minimal supersymmetry $s = 2, 4, 8, 16$.

Theorem 6.4. *The super Yang-Mills lagrangian* (6.3) *is supersymmetric in* $n = 3, 4, 6, 10$ *dimensions with minimal supersymmetry.*

There is an additional supersymmetric theory—the exception referred to above—in $n = 2$ dimensions with $s = (1, 0)$ supersymmetry; we will not consider it. The quantum argument above shows that (6.3) cannot be supersymmetric in other dimensions and with other amounts of supersymmetry.

Proof. We work in n-dimensional Minkowski space and fix a real spin representation S. Thus we have pairings (1.1), (1.2) which satisfy the Clifford relation (1.3). The spinor field λ takes values in $S^* \otimes \operatorname{ad} P$. The lagrangian (6.3) is

$$\boxed{L = \left(-\frac{1}{4} g^{\mu\sigma} g^{\nu\rho} \langle F_{\mu\nu}, F_{\sigma\rho}\rangle + \frac{1}{2}\Gamma^{\sigma}_{ab}\langle \lambda^a, \nabla_\sigma \lambda^b\rangle\right) |d^n x|,} \tag{6.5}$$

where $\langle \cdot, \cdot \rangle$ is the invariant inner product on the gauge algebra, the covariant derivative is $\nabla_\sigma = \partial_\sigma + A_\sigma$, and the curvature is

$$F_A = \frac{1}{2} F_{\mu\nu}\, dx^\mu \wedge dx^\nu, \tag{6.6}$$

where

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]. \tag{6.7}$$

We work with *real* bases for spinors and vectors. We postulate the induced action of the supersymmetry transformation $\eta^a Q_a$ on the fields to be

$$\boxed{\begin{aligned} (\hat{\xi} A)_\mu &= \eta^a g_{\mu\nu} \Gamma^\nu_{ab} \lambda^b \\ \hat{\xi}\lambda^a &= \frac{1}{2}\eta^b \tilde{\Gamma}^{\mu ac} \Gamma^\nu_{cb} F_{\mu\nu}. \end{aligned}} \tag{6.8}$$

Each side of the first equation is a component of a 1-form with values in the adjoint bundle. These equations are motivated by Lorentz invariance, gauge invariance, parity, and power counting.

There are two things we need to check: (i) $\hat{\xi}$ is a (nonmanifest) symmetry of L; and (ii) the Lie bracket of transformations (6.8) is consistent with (1.30). We will find the conditions under which (i) and (ii) hold.

[33] The free quantum theory is only used here to rule out some cases; otherwise the argument is entirely classical.

First, we investigate $\mathrm{Lie}(\hat{\xi})L$. The variation of the curvature is

$$\boxed{\hat{\xi}F_{\mu\nu} = \eta^a g_{\nu\sigma}\Gamma^\sigma_{ab}\nabla_\mu\lambda^b - \eta^a g_{\mu\sigma}\Gamma^\sigma_{ab}\nabla_\nu\lambda^b,} \tag{6.9}$$

from which we easily compute

$$\hat{\xi}\big(-\frac{1}{4}g^{\mu\sigma}g^{\nu\rho}\langle F_{\mu\nu}, F_{\sigma\rho}\rangle\big) = -\eta^a g^{\mu\sigma}\Gamma^\nu_{ab}\langle F_{\mu\nu}, \nabla_\sigma\lambda^b\rangle. \tag{6.10}$$

In the variation of the second term in (6.5) we must remember to include the variation of the covariant derivative $\nabla_\sigma = \partial_\sigma + A_\sigma$:

(6.11)

$$\begin{aligned}\hat{\xi}\big(\frac{1}{2}\Gamma^\sigma_{ab}\langle\lambda^a, \nabla_\sigma\lambda^b\rangle\big) &= \frac{1}{2}\Gamma^\sigma_{ab}\langle\eta^c\tilde{\Gamma}^{\mu ad}\Gamma^\nu_{dc}F_{\mu\nu}, \nabla_\sigma\lambda^b\rangle + \frac{1}{2}\Gamma^\sigma_{ab}\langle\lambda^a, \eta^c g_{\sigma\nu}\Gamma^\nu_{cd}[\lambda^d, \lambda^b]\rangle \\ &\qquad + EXACT \\ &= \frac{1}{2}\eta^a\Gamma^\sigma_{bd}\tilde{\Gamma}^{\mu cd}\Gamma^\nu_{ac}\langle F_{\mu\nu}, \nabla_\sigma\lambda^b\rangle - \frac{1}{2}\eta^c g_{\sigma\nu}\Gamma^\sigma_{ab}\Gamma^\nu_{cd}\langle\lambda^a, [\lambda^d, \lambda^b]\rangle \\ &\qquad + EXACT \\ &= \eta^a g^{\mu\sigma}\Gamma^\nu_{ab}\langle F_{\mu\nu}, \nabla_\sigma\lambda^b\rangle - \frac{1}{2}\eta^c g_{\sigma\nu}\Gamma^\sigma_{ab}\Gamma^\nu_{cd}\langle\lambda^a, [\lambda^d, \lambda^b]\rangle \\ &\qquad + EXACT,\end{aligned}$$

where

$$EXACT = \partial_\sigma\big(\frac{1}{4}\eta^a\tilde{\Gamma}^\sigma_{bd}\Gamma^{\mu cd}\tilde{\Gamma}^\nu_{ac}\langle F_{\mu\nu}, \lambda^b\rangle\big). \tag{6.12}$$

In the last step we use the Clifford relation $\Gamma^\mu_{bd}\tilde{\Gamma}^{\sigma cd} + \Gamma^\sigma_{bd}\tilde{\Gamma}^{\mu cd} = 2g^{\mu\sigma}\delta^c_b$ and the Bianchi identity $\nabla_\sigma F_{\mu\nu} + \nabla_\nu F_{\sigma\mu} + \nabla_\mu F_{\nu\sigma} = 0$. Now the first term in the last line cancels against (6.10), and the second term vanishes if and only if the quartic form

$$Q_{abcd} = g_{\sigma\nu}\big(\Gamma^\sigma_{ab}\Gamma^\nu_{cd} + \Gamma^\sigma_{ad}\Gamma^\nu_{bc} + \Gamma^\sigma_{ac}\Gamma^\nu_{db}\big) \tag{6.13}$$

vanishes. There are two conclusions from this computation. First, if the gauge algebra is abelian, then $\hat{\xi}$ is a symmetry of L in any dimension and with any spin representation S. Second, if the gauge algebra is nonabelian, then $\hat{\xi}$ is a symmetry of L if and only if the quartic form

$$Q\colon s^* \longmapsto |\,\Gamma(s^*, s^*)\,|^2, \qquad s^* \in S^*, \tag{6.14}$$

vanishes identically. This holds for $n = 3, 4, 6, 10$ as in those dimensions the spin group acts transitively on the nonzero spinors. For $n = 3, 4, 6$ the spin representation is $SL(2, \mathbb{F})$ acting on $\mathbb{F}^2$ for $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$. For $n = 10$ octonions can be similarly used [I-**Spinors**, §6.5–§6.7].

Next, we investigate the Lie algebra generated by the symmetry transformation (6.8). Let η^a_1, η^b_2 be odd parameters and $\hat{\xi}_i$ the action (6.8) of $\eta^a_i Q_a$. Recall from (1.30) that for any field f we need to check

$$[\hat{\xi}_1, \hat{\xi}_2]f = 2\eta^a_1\eta^b_2\Gamma^\sigma_{ab}\partial_\sigma f \tag{6.15}$$

When acting on the spinor field $f = \lambda$, we replace ∂_σ with the covariant derivative ∇_σ. When acting on the connection $f = A$, we use Lie derivative by the horizontal lift, and it is easy to compute that the action of ∂_σ on A has μ^{th} component equal to $F_{\sigma\mu}$.

We first verify (6.15) for the connection field. Using (6.8) we find:

$$\begin{aligned}\big([\hat{\xi}_1, \hat{\xi}_2]A\big)_\mu &= -\frac{1}{2}\eta_1^a\eta_2^b g_{\mu\nu}(\tilde{\Gamma}^{\rho cd}\Gamma^\nu_{bc}\Gamma^\sigma_{ad} + \tilde{\Gamma}^{\rho cd}\Gamma^\nu_{ac}\Gamma^\sigma_{bd})F_{\rho\sigma} \\ &= -\frac{1}{2}\eta_1^a\eta_2^b g_{\mu\nu}(4g^{\rho\nu}\Gamma^\sigma_{ab})F_{\rho\sigma} \\ &= 2\eta_1^a\eta_2^b\Gamma^\sigma_{ab}F_{\sigma\mu}.\end{aligned} \tag{6.16}$$

To pass to the second line we added to the first line terms in parentheses with ρ and ν exchanged; then we used the Clifford identity (1.3). This is the desired result. To check the commutation relation (6.15) on the spinor field we need to also impose the equation of motion

$$\Gamma^\sigma_{ab}\nabla_\sigma\lambda^a = 0, \qquad \text{for all } b; \tag{6.17}$$

that is, we only verify the supersymmetry algebra *on-shell*. Then we compute from (6.8)

$$\begin{aligned}[\hat{\xi}_a, \hat{\xi}_2]\lambda^c &= -\frac{1}{2}\eta_1^a\eta_2^b g_{\mu\nu}(\tilde{\Gamma}^{\sigma cd}\Gamma^\nu_{db}\Gamma^\mu_{ae} + \tilde{\Gamma}^{\sigma cd}\Gamma^\nu_{da}\Gamma^\mu_{be} \\ &\qquad - \tilde{\Gamma}^{\nu cd}\Gamma^\sigma_{db}\Gamma^\mu_{ae} - \tilde{\Gamma}^{\nu cd}\Gamma^\sigma_{da}\Gamma^\mu_{be})\nabla_\sigma\lambda^e \\ &= \frac{1}{2}\eta_1^a\eta_2^b g_{\mu\nu}\tilde{\Gamma}^{\sigma cd}\Gamma^\nu_{de}\Gamma^\mu_{ab}\nabla_\sigma\lambda^e - \frac{1}{2}\eta_1^a\eta_2^b\tilde{\Gamma}^{\sigma cd}Q_{abde}\nabla_\sigma\lambda^e \\ &\qquad + \frac{1}{2}\eta_1^a\eta_2^b g_{\mu\nu}(\tilde{\Gamma}^{\nu cd}\Gamma^\sigma_{db}\Gamma^\mu_{ae} + \tilde{\Gamma}^{\nu cd}\Gamma^\sigma_{da}\Gamma^\mu_{be})\nabla_\sigma\lambda^e.\end{aligned} \tag{6.18}$$

Let 'LT' denote the last term. Applying the Clifford identity we see that the symmetric part in ν, σ vanishes by the equation of motion, whence

$$\begin{aligned}LT &= \frac{1}{2}LT - \frac{1}{4}\eta_1^a\eta_2^b g_{\mu\nu}(\tilde{\Gamma}^{\sigma cd}\Gamma^\nu_{db}\Gamma^\mu_{ae} + \tilde{\Gamma}^{\sigma cd}\Gamma^\nu_{da}\Gamma^\mu_{be})\nabla_\sigma\lambda^e \\ &= \frac{1}{2}LT + \frac{1}{4}\eta_1^a\eta_2^b g_{\mu\nu}\tilde{\Gamma}^{\sigma cd}\Gamma^\nu_{de}\Gamma^\mu_{ab}\nabla_\sigma\lambda^e - \frac{1}{4}\eta_1^a\eta_2^b\tilde{\Gamma}^{\sigma cd}Q_{abde}\nabla_\sigma\lambda^e.\end{aligned} \tag{6.19}$$

Solving for LT and plugging into (6.18) we have

$$\begin{aligned}[\hat{\xi}_1, \hat{\xi}_2]\lambda^c &= \eta_1^a\eta_2^b g_{\mu\nu}\tilde{\Gamma}^{\sigma cd}\Gamma^\nu_{de}\Gamma^\mu_{ab}\nabla_\sigma\lambda^e - \eta_1^a\eta_2^b\tilde{\Gamma}^{\sigma cd}Q_{abde}\nabla_\sigma\lambda^e \\ &= 2\eta_1^a\eta_2^b\Gamma^\sigma_{ab}\nabla_\sigma\lambda^c - \eta_1^a\eta_2^b\tilde{\Gamma}^{\sigma cd}Q_{abde}\nabla_\sigma\lambda^e.\end{aligned} \tag{6.20}$$

Here we use the equation of motion (6.17) to see that the last term in the first line vanishes when σ and ν are exchanged. If the second term vanishes, then this is the desired bracket. Hence the supersymmetry algebra is satisfied on-shell if and only if the quartic form (6.14) vanishes. This applies both to the abelian and nonabelian theories.

This completes the proof that (6.3) is supersymmetric.

We give another interpretation of the quartic form below (6.69).

We now compute the classical equations of motion and the variational 1-form. First, the variation of the curvature (6.7) is

$$\delta F_{\mu\nu} = \nabla_\mu \delta A_\nu - \nabla_\nu \delta A_\mu. \tag{6.21}$$

Now write the lagrangian (6.5) as $L = \check{L}\,|d^n x|$. Then a short computation yields

$$\begin{aligned}\delta\check{L} = \langle \delta A_\rho\,,\, g^{\mu\sigma}g^{\nu\rho}\nabla_\sigma F_{\mu\nu} - \frac{1}{2}\Gamma^\rho_{ab}[\lambda^a,\lambda^b]\rangle + \langle \delta\lambda^a\,,\,\Gamma^\sigma_{ab}\nabla_\sigma\lambda^b\rangle \\ + \partial_\sigma\Big\{-g^{\mu\sigma}g^{\nu\rho}\langle F_{\mu\nu},\delta A_\rho\rangle + \frac{1}{2}\Gamma^\sigma_{ab}\langle\lambda^a,\delta\lambda^b\rangle\Big\}.\end{aligned} \tag{6.22}$$

This leads to the equations of motion

$$\boxed{\begin{aligned} g^{\mu\sigma}g^{\nu\rho}\nabla_\sigma F_{\mu\nu} &= \frac{1}{2}\Gamma^\rho_{ab}[\lambda^a,\lambda^b] \\ \Gamma^\sigma_{ab}\nabla_\sigma\lambda^b &= 0\end{aligned}} \tag{6.23}$$

and the variational 1-form

$$\boxed{\gamma = \Big(-g^{\mu\sigma}g^{\nu\rho}\langle F_{\mu\nu},\delta A_\rho\rangle + \frac{1}{2}\Gamma^\sigma_{ab}\langle\lambda^a,\delta\lambda^b\rangle\Big)\,\frac{\partial}{\partial x^\sigma}\otimes|d^n x|.} \tag{6.24}$$

In coordinate-free notation we write the equations of motion (6.23) as

$$\boxed{\begin{aligned} -d_A * F_A &= \frac{1}{2}[\lambda\,\Gamma\,\lambda]\;|d^n x| \\ \not{D}_A\lambda &= 0.\end{aligned}} \tag{6.25}$$

The Noether current associated to $\hat{\xi}$ is computed from (6.12) and (6.24) as

$$\iota(\hat{\xi})\gamma - \Big(\frac{1}{4}\eta^a\Gamma^\sigma_{bd}\tilde{\Gamma}^{\mu cd}\Gamma^\nu_{ac}\langle F_{\mu\nu},\lambda^b\rangle\,\frac{\partial}{\partial x^\sigma}\otimes|d^n x|\Big) = -\eta^a g^{\mu\sigma}\Gamma^\nu_{ab}\langle F_{\mu\nu},\lambda^b\rangle\,\frac{\partial}{\partial x^\sigma}\otimes|d^n x|. \tag{6.26}$$

The *supercurrent* j_a is *minus* the Noether current for Q_a:

$$\boxed{j_a = g^{\mu\sigma}\Gamma^\nu_{ab}\langle F_{\mu\nu},\lambda^b\rangle\,\frac{\partial}{\partial x^\sigma}\otimes|d^n x|.} \tag{6.27}$$

§6.2. Gauge theories with matter

The most general bosonic model without gravity[34] on Minkowski space $\check{M}^n$ is specified by the following data:

(6.28)	G	compact Lie group
	$\langle\cdot,\cdot\rangle$	bi-invariant scalar product on $\mathfrak{g}$
	X	Riemannian manifold on which G acts by isometries
	$V\colon X \longrightarrow \mathbb{R}$	potential function invariant under G

[34]We include only scalar fields and gauge fields. There are also models with p-form fields for $p \geq 2$, for example. The models discussed here cover most *fundamental* (vs. *effective*) lagrangians without gravity.

The fields in the model are

$$\begin{array}{ll} A & \text{connection on some principal } G \text{ bundle } P \longrightarrow \check{M}^n \\ \phi & \text{section of the associated bundle } X^P = P \times_G X \longrightarrow \check{M}^n \end{array} \tag{6.29}$$

The standard fundamental lagrangian is then the "minimally coupled" σ-model lagrangian (3.77) and Yang-Mills lagrangian (6.2):

$$L = \left\{ -\frac{1}{2}|F_A|^2 + \frac{1}{2}|d_A\phi|^2 - \phi^* V \right\} |d^n x|. \tag{6.30}$$

The moduli space of classical vacua on $\check{M}^n$ is the quotient of the set of minima of V—which we assume occurs at $V = 0$—by the action of constant gauge transformations:

$$\mathcal{M}_{\text{vac}} = V^{-1}(0)/G. \tag{6.31}$$

Now we consider supersymmetric extensions of (6.30). As in §3 we work with models in 3, 4, and 6 dimensions with minimal supersymmetry. As in §3 we refer to the model in $3, 4, 6$ dimensions as the $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ model. In these dimensions we have already constructed separately the supersymmetric σ-model and the supersymmetric gauge theory. The next theorem summarizes what happens when we combine them. In these models the bosons are given in (6.29) and the fermions are

$$\begin{array}{ll} \psi & \text{section of } \phi^*\big(P \times_G (TX \otimes_{\mathbb{F}} \Pi\mathbb{F}^2)\big) \\ \lambda & \text{section of } P \times_G (\mathfrak{g} \otimes \Pi\mathbb{F}^{2^*}) \end{array} \tag{6.32}$$

In the $\mathbb{F} = \mathbb{H}$ case we use complex notation for ψ as in (3.49).

A discussion—including an explanation of notation—follows the statement of the theorem.

Theorem 6.33. *Fix*

$$\begin{array}{ll} G & \textit{compact Lie group} \\ \langle \cdot, \cdot \rangle & \textit{bi-invariant scalar product on } \mathfrak{g} \end{array} \tag{6.34}$$

(*i*) ($\mathbb{F} = \mathbb{R}$) *Suppose we are given*

$$\begin{array}{ll} X & \textit{Riemannian manifold with } G \textit{ action by isometries} \\ h\colon X \longrightarrow \mathbb{R} & G\textit{-invariant real-valued function} \end{array} \tag{6.35}$$

Then the lagrangian

$$\boxed{\begin{aligned} L = \Big\{ &-\frac{1}{2}|F_A|^2 + \frac{1}{2}|d_A\phi|^2 + \frac{1}{2}\langle \lambda \not{D}_A \lambda \rangle + \frac{1}{2}\langle \psi (\not{D}_A)_\phi \, \psi \rangle - \frac{1}{2}\phi^* |\operatorname{grad} h|^2 \\ &- \frac{1}{2}\epsilon^{ab}\phi^*(\operatorname{Hess} h)(\psi_a, \psi_b) + \langle \hat{\lambda}^a_\phi, \psi_a \rangle \\ &+ \frac{1}{12}\epsilon^{ab}\epsilon^{cd}\langle \psi_a, \phi^* R(\psi_b, \psi_c)\psi_d \rangle \Big\} \, |d^3 x|. \end{aligned}} \tag{6.36}$$

on $\check{M}^3$ is supersymmetric with the supersymmetry transformation laws

$$\boxed{\begin{aligned}\hat{\xi}\phi &= -\eta^a\,\psi_a\\ (\hat{\xi}A)_{ab} &= \eta^c(\epsilon_{ac}\epsilon_{bd}+\epsilon_{ad}\epsilon_{bc})\lambda^d\\ \hat{\xi}\psi_a &= \quad \eta^b\big((d_A)_{ab}\phi+\epsilon_{ab}\phi^*\operatorname{grad}h\big)\\ \hat{\xi}\lambda^a &= -\eta^b\epsilon^{ac}f_{bc}\end{aligned}} \tag{6.37}$$

The potential energy is

$$\boxed{V=\frac{1}{2}|\operatorname{grad}h|^2} \tag{6.38}$$

and the moduli space of vacua is

$$\boxed{\mathcal{M}_{\text{vac}}=\operatorname{Crit}(h)/G.} \tag{6.39}$$

The supercurrent j_a, which is minus the Noether current for Q_a, is

$$\boxed{\begin{aligned}j_a=\Big\{\epsilon^{cb}\epsilon^{de}\langle\partial_{ae}\phi,\psi_b\rangle+\epsilon^{bd}\delta^c_a\langle\phi^*\operatorname{grad}h,\psi_b\rangle+\frac{1}{2}\epsilon^{bd}\langle f_{ba},\lambda^c\rangle\Big\}\frac{\partial}{\partial y^{cd}}\otimes|d^3x|\\ +\frac{1}{2}\epsilon^{cb}\langle f_{cd},\lambda^d\rangle\,\frac{\partial}{\partial y^{ab}}\otimes|d^3x|\end{aligned}} \tag{6.40}$$

(ii) ($\mathbb{F}=\mathbb{C}$) *Suppose we are given*

$$\begin{array}{ll} X & \textit{Kähler manifold with } G \textit{ action}\\ \mu\colon X\longrightarrow\mathfrak{g}^* & \textit{moment map for } G \textit{ action}\\ W\colon X\longrightarrow\mathbb{C} & G\textit{-invariant holomorphic function}\end{array} \tag{6.41}$$

Then the lagrangian

$$\boxed{\begin{aligned}L=\Big\{-\frac{1}{2}|F_A|^2+\langle d_A\bar{\phi},d_A\phi\rangle+\langle\bar{\lambda}\not{D}_A\lambda\rangle+\langle\bar{\psi}(\not{D}_A)_\phi\,\psi\rangle\\ -\phi^*\|\operatorname{grad}W\|^2-2\phi^*|\mu|^2-\operatorname{Re}\big[\epsilon^{ab}\phi^*(\operatorname{Hess}W)(\psi_a,\psi_b)\big]\\ +\sqrt{2}\left(\langle\hat{\lambda}^a_\phi,\psi_a\rangle-\langle\bar{\psi}_{\dot{a}},\hat{\bar{\lambda}}^{\dot{a}}_\phi\rangle\right)-\frac{1}{4}\epsilon^{ac}\epsilon^{\dot{b}\dot{d}}\langle\psi_a,\phi^*R(\psi_c,\bar{\psi}_{\dot{d}})\bar{\psi}_{\dot{b}}\rangle\Big\}\,|d^4x|.\end{aligned}}$$

on $\check{M}^4$ is supersymmetric with the supersymmetry transformation laws

$$\boxed{\begin{aligned}\hat{\xi}\phi &= -\sqrt{2}\,\eta^a\psi_a\\ (\hat{\xi}A)_{a\dot{b}} &= 2\eta^c\,\epsilon_{ac}\epsilon_{\dot{b}\dot{d}}\,\bar{\lambda}^{\dot{d}}+2\bar{\eta}^{\dot{c}}\,\epsilon_{ad}\epsilon_{\dot{b}\dot{c}}\,\lambda^d\\ \hat{\xi}\psi_a &= \sqrt{2}\left[\bar{\eta}^{\dot{b}}(d_A)_{a\dot{b}}\phi+\eta^b\epsilon_{ab}\phi^*\operatorname{grad}W\right]\\ \hat{\xi}\lambda^a &= -\eta^b\epsilon^{ac}F_{bc}-2\eta^a\sqrt{-1}\phi^*\check{\mu}\end{aligned}} \tag{6.43}$$

The potential energy is

$$\boxed{V = \|\operatorname{grad} W\|^2 + 2|\mu|^2} \tag{6.44}$$

and the moduli space of vacua is

$$\boxed{\mathcal{M}_{\text{vac}} = \mu^{-1}(0) \cap \operatorname{Crit}(W) \,/\, G.} \tag{6.45}$$

The supercurrent j_a, which is minus the Noether current for Q_a, is

(6.46)

$$\boxed{\begin{aligned} j_a = \Big\{ \sqrt{2}\epsilon^{cb}\epsilon^{\dot d\dot e}\langle \partial_{a\dot e}\phi, \psi_b\rangle + \sqrt{2}\,\epsilon^{\dot b\dot d}\delta_a^c\langle \phi^* \operatorname{grad} W, \bar\psi_{\dot b}\rangle + \frac{1}{2}\epsilon^{bc}\langle F_{ba}, \bar\lambda^{\dot d}\rangle \Big\} \frac{\partial}{\partial y^{c\dot d}} \otimes |d^4x| \\ + \Big\{ \frac{1}{2}\epsilon^{\dot c\dot b}\langle \overline{F}_{\dot c\dot d}, \bar\lambda^{\dot d}\rangle - 2\sqrt{-1}\,\phi^*\mu(\bar\lambda^{\dot b}) \Big\} \frac{\partial}{\partial y^{a\dot b}} \otimes |d^4x| \end{aligned}}$$

(*iii*) ($\mathbb{F} = \mathbb{H}$) *Suppose we are given*

$$\begin{array}{ll} X & \textit{hyperkähler manifold with } G \textit{ action} \\ \mu_{\mathbb{H}} \colon X \longrightarrow \mathfrak{g}^* \otimes \operatorname{Im}\mathbb{H} & \textit{hyperkähler moment map} \end{array} \tag{6.47}$$

Then the lagrangian

$$\boxed{\begin{aligned} L = \Big\{ -\frac{1}{2}|F_A|^2 + \frac{1}{16}\tilde h_{\alpha\beta}\epsilon_{ij}\nu^{abcd}(\partial_{ab}\phi)^{\alpha i}(\partial_{cd}\phi)^{\beta j} + \frac{1}{2}\langle \lambda \not{D}_A \lambda\rangle \\ + \frac{1}{4}h_{\alpha\beta}\nu^{abcd}\psi_a^\alpha(\phi^*\nabla_{cd})\psi_b^\beta - 2\phi^*|\mu_{\mathbb{H}}|^2 + \sqrt{2}\,\tilde h_{\alpha\beta}\epsilon_{ij}(\hat\lambda_\phi^{aj})^{\alpha i}\psi_a^\beta \\ - \frac{1}{24}\nu^{abcd}\phi^*\Omega_{\alpha\beta\gamma\delta}\psi_a^\alpha\psi_b^\beta\psi_c^\gamma\psi_d^\delta \Big\}\, |d^6x|. \end{aligned}} \tag{6.48}$$

on $\check M^6$ is supersymmetric with the supersymmetry transformation laws

$$\boxed{\begin{aligned} (\hat\xi\phi)^{\alpha i} &= -\sqrt{2}\,\eta^{ai}\psi_a^\alpha \\ (\hat\xi A)_{pq} &= 2\eta^{ai}\nu_{pqab}\epsilon_{ij}\lambda^{bj} \\ \hat\xi\psi_a^\alpha &= \sqrt{2}\,\epsilon_{ij}\eta^{bj}\big((d_A)_{ab}\phi\big)^{\alpha i} \\ \hat\xi\lambda^{ai} &= -\eta^{bi} f_b^a - 2\eta^{aj}\sqrt{-1}\,\phi^*(\check\mu_{\mathbb{H}})_j^i. \end{aligned}} \tag{6.49}$$

The potential energy is

$$\boxed{V = 2|\mu_{\mathbb{H}}|^2} \tag{6.50}$$

and the moduli space of vacua is the hyperkähler quotient

$$\boxed{\mathcal{M}_{\text{vac}} = \mu_{\mathbb{H}}^{-1}(0) \,/\, G.} \tag{6.51}$$

The supercurrent j_{ai}, which is minus the Noether current for Q_{ai}, is

(6.52)
$$\boxed{\begin{aligned} j_{ai} = &\left\{\frac{1}{\sqrt{2}}\nu^{bcde}\langle\partial_{ac}\phi,\psi_{bi}\rangle + \frac{1}{2}\epsilon_{ij}\langle f_a^e,\lambda^{dj}\rangle\right\}\frac{\partial}{\partial y^{de}}\otimes|d^6x| \\ &-\left\{\frac{1}{2}\epsilon_{ij}\langle f_b^c,\lambda^{bj}\rangle + 2\sqrt{-1}\,\epsilon_{jk}\langle\phi^*(\check{\mu}_{\mathbb{H}})_i^j,\lambda^{ck}\rangle\right\}\frac{\partial}{\partial y^{ac}}\otimes|d^6x|. \end{aligned}}$$

The proof for $\mathbb{F}=\mathbb{R}$ is given in §7.3, and the proof[35] for $\mathbb{F}=\mathbb{C}$ (at least when X is Hodge) in §8.3. In both cases we give a manifestly supersymmetric formulation is superspace. We do not give a proof for $\mathbb{F}=\mathbb{H}$, a case in which there is no known superspace formulation for an arbitrary hyperkähler manifold X, though we have done enough of the computations in components that the reader should have no problem completing the proof directly in components. The reduction of (6.48) to 4 dimensions is discussed in §10.3.

In (6.37) the Hodge dual of the curvature F_A is

$$*F_A = f_{ab}dy^{ab}. \tag{6.53}$$

In (6.43) the curvature F_A has been decomposed into its self-dual and anti-self-dual parts:

$$F_A = \frac{1}{2}(F_{ab}\epsilon_{\dot{c}\dot{d}} + \overline{F}_{\dot{c}\dot{d}}\epsilon_{ab})\,dy^{a\dot{c}}\wedge dy^{b\dot{d}}. \tag{6.54}$$

In (6.49) the curvature 2-form

$$F_A = \frac{1}{8}\,F_{(ab)(cd)}\,dy^{ab}\wedge dy^{cd} \tag{6.55}$$

has been rewritten in terms of a traceless 4×4 matrix f:

$$f_b^a = \frac{1}{4}\nu^{acde}F_{(bc)(de)}. \tag{6.56}$$

We explain this further in §10.1.

For $\mathbb{F}=\mathbb{C}$ the scalar product on $\mathfrak{g}$ identifies $\mathfrak{g}^*\cong\mathfrak{g}$. We let

$$\check{\mu}\colon X\to\mathfrak{g} \tag{6.57}$$

denote the composition of the moment map with this isomorphism; for $\mathbb{F}=\mathbb{H}$ we have $\check{\mu}_{\mathbb{H}}\colon X\to\mathfrak{g}\otimes\operatorname{Im}\mathbb{H}$. In the last equation in (6.49) we identify $\operatorname{Im}\mathbb{H}$ with traceless hermitian 2×2 matrices:

$$aI+bJ+cK\leftrightarrow\begin{pmatrix} a\sqrt{-1} & -b+c\sqrt{-1}\\ b+c\sqrt{-1} & -a\sqrt{-1}\end{pmatrix}. \tag{6.58}$$

In all three cases there are new terms of the form $\langle\hat{\lambda}_\phi,\psi\rangle$ which appear in the lagrangian. Here $\hat{\lambda}_\phi$ is the odd vector field on $P\times_G X$ corresponding to λ, and the

[35]See Problem ASH4 of [I-**Homework**] for an example, which in particular serves as a check of the constants in (6.42) and (6.43).

inner product is the Riemannian metric on X. For $\mathbb{F} = \mathbb{R}$, in the linear case X is an orthogonal vector space and G acts linearly by an orthogonal representation $\rho\colon G \to O(X)$. Then

$$\hat{\lambda}_\phi = \dot{\rho}(\lambda)\phi, \tag{6.59}$$

where $\dot{\rho}\colon \mathfrak{g} \to \mathfrak{o}(X)$ is the infinitesimal action. This term can be predicted by supersymmetry: the variation in A of $\langle \psi \not{D}_A \psi \rangle$ is canceled by the variation in ϕ of this term. For $\mathbb{F} = \mathbb{C}$ note that ψ_a is a vector field of type $(1,0)$ on X, so only the $(0,1)$-component of $\hat{\lambda}^a_\phi$ and its conjugate enter (6.42). In the linear case G acts by a unitary representation $\rho\colon G \to U(X)$ on a hermitian vector space X. Then the $(1,0)$-component of $\hat{\lambda}_\phi$ is

$$(\hat{\lambda}_\phi)^{1,0} = \dot{\rho}_{\mathbb{C}}(\lambda)\phi, \tag{6.60}$$

where $\dot{\rho}_{\mathbb{C}}\colon \mathfrak{g}_{\mathbb{C}} \to \mathfrak{gl}_{\mathbb{C}}(X)$ is the complexification of the differential to ρ. To write a formula for the $(0,1)$-component we identify $\overline{X} \cong X^*$ via the hermitian metric; then

$$(\hat{\lambda}_\phi)^{0,1} = -\dot{\rho}_{\mathbb{C}}(\lambda)^*\phi^* \in X^* \cong \overline{X} \tag{6.61}$$

at the point $\phi^* \in X^*$. (As a real manifold, $X^* \cong \overline{X} \cong X$ so we can identify points of X with points of X^*.) Note that when λ is real, then $\dot{\rho}_{\mathbb{C}}(\lambda)$ is skew-hermitian and (6.61) is the complex conjugate of (6.60), i.e., $\hat{\lambda}_\phi$ is real. Also,[36]

$$\hat{\bar{\lambda}} = \overline{\hat{\lambda}}, \tag{6.62}$$

which also holds in the nonlinear case. This ensures that (6.42) is real.

The quotient in (6.45) is the symplectic (Kähler) quotient, which can often be identified with the complex quotient of $\mathrm{Crit}(W)$ by the complexification of G. The quotient in (6.51) is the hyperkähler quotient. (It was in this context that the hyperkähler quotient was discovered.) For $\mathbb{F} = \mathbb{C}$ the abelian group $\big(\mathfrak{g}/[\mathfrak{g},\mathfrak{g}]\big)^*$ of infinitesimal characters acts by translations on the space of moment maps. Theories with different choices of moment map are different. Notice that when X is a point we must still choose a moment map, which is now simply an infinitesimal character. The resulting term $-\big(2|\mu|^2\big)$ is usually called a *Fayet-Iliopoulos term.* For $\mathbb{F} = \mathbb{H}$ the moment map is also ambiguous, this time up to $\big(\mathfrak{g}/[\mathfrak{g},\mathfrak{g}]\big)^* \otimes \operatorname{Im}\mathbb{H}$. For X a point this leads to a Fayet-Iliopoulos term.

The action of R-symmetries for $\mathbb{F} = \mathbb{C}$ and $\mathbb{F} = \mathbb{H}$ is summarized in (5.31), (8.24), and (10.21).

We have not written the equations of motion for these general lagrangians; it seems better to work out the detailed form on a case-by-case basis. However, we indicate here some general features. For the scalar field ϕ we find an equation of the form

$$\Box_A \phi = -\phi^* \operatorname{grad} V + Q^{(2)}_\phi(\psi,\lambda) + R^{(2)}_\phi(\psi)d\phi + Q^{(4)}(\psi). \tag{6.63}$$

[36] We caution the reader that '$\bar{\lambda}$' denotes the conjugate in the Lie algebra. For example, if $G = \mathbb{T}$ is the circle group, then $\mathfrak{g} \cong \sqrt{-1}\,\mathbb{R}$ and $\mathfrak{g}_{\mathbb{C}} \cong \mathbb{C}$ with conjugation *minus* the usual complex conjugation in $\mathbb{C}$. See Problem ASH4 of [I-**Homework**] for a sample lagrangian in this case.

Here $\Box_A = d_A^* d_A$ is the wave operator on sections of $X^P \to \check{M}^n$. The coefficients of the quadratic form $Q_\phi^{(2)}$ depend on the pullback of the hessian of h or W (for $\mathbb{F} = \mathbb{R}$ or $\mathbb{F} = \mathbb{C}$) and also on the action of $\mathfrak{g}$ on X; the coefficients of the quadratic form $R_\phi^{(2)}$ depend on the pullback of the Riemann curvature tensor; and the coefficients of the quartic form $Q_\phi^{(4)}$ depend on the pullback of derivatives of the Riemann curvature tensor. Now the action of $\mathfrak{g}$ on X induces a map from $\mathfrak{g}$ to vertical vector fields on X^P, or equivalently a $\mathfrak{g}^*$-valued vector field $\hat{\zeta}$ on X^P. This vector field enters into some equations of motion, for example in the equation

$$\not{D}_A \lambda = -\langle \hat{\zeta}_\phi, \psi \rangle. \tag{6.64}$$

Here $\hat{\zeta}_\phi$ means the pullback of $\hat{\zeta}$ by the section ϕ.

§6.3. Superspace construction

We consider only pure gauge theory. As usual we fix data (6.1). Consider a superspace $M^{n|s}$ in any dimension built out of any real spin representation. Suppose $\mathcal{P} \to M^{n|s}$ is a principal G bundle. Let $\mathcal{A}$ be a connection on $\mathcal{P}$. If $\mathcal{A}$ is unconstrained, then there are too many component fields to obtain a sensible theory, even with n, s small. Hence we impose a constraint. Namely, we require that the curvature $F_\mathcal{A}$ be constrained to vanish along the left-invariant odd distribution τ on $M^{n|s}$:

$$\iota(D_b)\iota(D_a)F_\mathcal{A} = 0. \tag{6.65}$$

In that case the "lowest" piece of the curvature is in the even-odd direction. Let $\mathcal{D}_a$ be the horizontal lift of D_a and ∇_μ the horizontal lift of ∂_μ. Define

$$\boxed{\Lambda^a = \frac{1}{n}\tilde{\Gamma}^{\mu ab}\iota(D_b)\iota(\partial_\mu)F_\mathcal{A} = \frac{1}{n}\tilde{\Gamma}^{\mu ab}[\mathcal{D}_b, \nabla_\mu].} \tag{6.66}$$

(The constant is chosen with hindsight.) Also, we identify G-invariant vertical vector fields with sections of the adjoint bundle; see [I-**Classical Fields**, §4.2].) This is a "projection" of the lowest nonvanishing part of the curvature; Λ^a is an odd section of the adjoint bundle $\operatorname{ad} P = \mathcal{P} \times_G \mathfrak{g} \to M^{n|s}$. There are component fields

$$\begin{aligned} A = i^*\mathcal{A} \qquad & \text{connection on } P = i^*\mathcal{P} \longrightarrow \check{M}^n \\ \lambda^a = i^*\Lambda^a \qquad & \text{dual spinor field on } \check{M}^n \text{ with values in } \operatorname{ad} P \end{aligned} \tag{6.67}$$

and possibly additional component fields as well. To see what other component fields there are, we examine consequences of the Bianchi identity.

First, we apply Bianchi to D_a, D_b, D_c to learn that

$$\text{The symmetrization of } \Gamma^\mu_{ab}[\mathcal{D}_c, \nabla_\mu] \text{ in } a, b, c \text{ vanishes.} \tag{6.68}$$

To interpret this, consider the sequence of maps

$$\begin{array}{ccccc} \operatorname{Sym}^3(S^*) & \xrightarrow{\operatorname{Sym}^3(\Gamma\otimes\mathrm{id})} & V \otimes S^* & \xrightarrow{g\Gamma} & S \\ f_a f_b f_c & \longrightarrow & \operatorname{sym}(\Gamma^\mu_{ab} e_\mu \otimes f_c) & & \\ & & e_\mu \otimes f_c & \longrightarrow & g_{\mu\nu}\Gamma^\nu_{cd} f^d \end{array} \tag{6.69}$$

The composition vanishes if and only if for all s^* in S^*, $\Gamma(s^*, s^*) \otimes s^*$ maps to 0 in S, i.e., if and only if for all $t^* \in S^*$,

$$\langle \text{image of } \Gamma(s^*, s^*) \otimes s^*, t^* \rangle = \langle \Gamma(s^*, s^*), \Gamma(s^*, t^*) \rangle \tag{6.70}$$

vanishes. The last expression is obtained by polarization from$\langle \Gamma(s^*, s^*), \Gamma(s^*, s^*) \rangle$, and the composition vanishes if and only if the quadratic form (6.14) does. As we have seen, this occurs for $n = \dim V = 3, 4, 6, 10$ with minimal supersymmetry $s = \dim S = 2, 4, 8, 16$. Furthermore, in these cases the sequence (6.69) is exact. Assume from now on that we are in one of these situations. Now the dual to (6.69) is

$$S^* \longrightarrow V^* \otimes S \longrightarrow \mathrm{Sym}^3(S). \tag{6.71}$$

The even-odd piece of the curvature can be identified with an element of $\Omega^0(M^{n|s}, \mathrm{ad}\,\mathcal{P}) \otimes V^* \otimes S$;then (6.68) asserts that its image in $\Omega^0(M^{n|s}, \mathrm{ad}\,\mathcal{P}) \otimes \mathrm{Sym}^3(S)$ vanishes. By the exactness of (6.71) we can then identify the even-odd piece of the curvature with an element $\Lambda \in \Omega^0(M^{n|s}, \mathrm{ad}\,\mathcal{P}) \otimes S^*$, as defined in (6.66). In fact, the Clifford identity implies that

$$\begin{aligned} V^* \otimes S &\longrightarrow S^* \\ e^\mu \otimes f^a &\longmapsto \frac{1}{n} \tilde{\Gamma}^{\mu ab} f_b \end{aligned} \tag{6.72}$$

is a splitting of (6.71), whence the formula (6.66) for Λ. To summarize, in the four cases we are considering the even-odd curvature can be expressed in terms of Λ:

$$\boxed{[\mathcal{D}_a, \nabla_\mu] = g_{\mu\nu} \Gamma^\nu_{ab} \Lambda^b.} \tag{6.73}$$

Next, we apply Bianchi to D_a, D_b, ∂_μ. After some manipulation with (6.73) and the Clifford identity we find

$$-2\tilde{\Gamma}^{\mu ac} \Gamma^\nu_{cb} \mathcal{F}_{\mu\nu} = g_{\mu\nu} \tilde{\Gamma}^{\mu ac} \Gamma^\nu_{bd} \mathcal{D}_c \Lambda^d + n \mathcal{D}_b \Lambda^a, \tag{6.74}$$

where the even-even part of the curvature is

$$\mathcal{F}_{\mu\nu} = \iota(\partial_\nu)\iota(\partial_\mu) F_{\mathcal{A}} = -[\nabla_\mu, \nabla_\nu]. \tag{6.75}$$

(The funny sign is due to the fact that we take the bracket as vector fields on P, not as operators on an associated bundle; cf. [I-**Classical Fields**, §4.2].) The first term on the right hand side of (6.74) depends on the particular case, and its precise form determines the structure of the auxiliary fields. In $n = 3, 4, 6$ dimensions we will find a formula of the form

$$\boxed{\mathcal{D}_b \Lambda^a = -\frac{1}{2} \tilde{\Gamma}^{\mu ac} \Gamma^\nu_{cb} \mathcal{F}_{\mu\nu} + \mathcal{E},} \tag{6.76}$$

where $i^*\mathcal{E}$ consists of auxiliary fields. (Compare with the second equation in (6.8).)

We do not learn anything new from the remaining cases of the Bianchi identity. Applied to $D_a, \partial_\mu, \partial_\nu$ we obtain the Jacobi identity for three vector fields $\Lambda^a, \partial_\mu, \partial_\nu$

on $\mathcal{P}$. (Here we identify Λ^a with a G-invariant vertical vector field. Upon restriction to $\check{M}^n$ this equation gives the supersymmetry transformation law (6.9) for the curvature.) Bianchi applied to $\partial_\mu, \partial_\nu, \partial_\rho$ restricts on $\check{M}^n$ to the usual Bianchi identity for A.

Set

$$F_{\mu\nu} = i^*\mathcal{F}_{\mu\nu}; \tag{6.77}$$

then

$$F_A = \frac{1}{2}F_{\mu\nu}dx^\mu \wedge dx^\nu \tag{6.78}$$

is the curvature of A.

From the general formula (1.29) and equations (6.73) and (6.76) we recover the supersymmetry transformation law (6.8), at least if we set $\mathcal{E} = 0$. For example, to compute the action of $\hat{\xi}$ on the connection A, we lift the vector field $\eta^a D_a$ on $M^{n|s}$ to the horizontal vector field $-\eta^a\mathcal{D}_a$ on $\mathcal{P}$ and apply the covariant derivative. From (6.73) the μ^{th} component of the answer is

$$\begin{aligned}(\hat{\xi}A)_\mu &= \iota(\partial_\mu)i^*\operatorname{Lie}(-\eta^a\mathcal{D}_a)A \\ &= i^*\iota(\nabla_\mu)\operatorname{Lie}(-\eta^a\mathcal{D}_a)A \\ &= i^*\Big[\operatorname{Lie}(-\eta^a\mathcal{D}_a)\iota(\nabla_\mu)A - \iota\big([-\eta^a\mathcal{D}_a, \nabla_\mu]\big)A\Big] \\ &= i^*\eta^a[\mathcal{D}_a, \nabla_\mu] \\ &= \eta^a g_{\mu\nu}\Gamma^\nu_{ab}\lambda^b,\end{aligned} \tag{6.79}$$

which agrees with (6.8). The precise formula for the variation of the fermion λ, which includes the auxiliary fields, must be analyzed on a case-by-case basis.

In §7, §8, and §10 we construct the superspace models for the cases $n = 3, 4$, and 6 (previously referred to as $\mathbb{F} = \mathbb{R}, \mathbb{C}$, and $\mathbb{H}$). Supersymmetry is manifest in the superspace formulation, and we determine the auxiliary fields by computing $\mathcal{E}$ in (6.76). In addition, for $n = 3, 4$ we give a superspace formulation of the gauge theory coupled to matter, and so prove Theorem 6.33(i) and Theorem 6.33(ii). There is no known superspace formulation for the $n = 10$ dimensional theory.

There are also superspace models for dimensional reductions of the $n = 3, 4, 6$ theories; particular cases are treated in §9 and §10. We remark here that (6.65) is no longer the correct curvature constraint in superspace; some components of the odd-odd curvature are allowed to be nonzero.

CHAPTER 7
$N = 1$ Yang-Mills Theory in Dimension 3

For the scalar superfield of Chapter 4 and the chiral superfield of Chapter 5 it is easy to see that the supermanifold of superfields is diffeomorphic to the supermanifold of component fields. But for constrained connections the argument is more complicated due to gauge symmetry. The precise statement is the main task of §7.1. In §7.2 and §7.3 we describe the superspace formulation of the Yang-Mills lagrangian and of the lagrangian including matter. There are no surprises, though we do explain where the new term in the component lagrangian (6.36) (coupling the two sorts of spinors) arises from the superspace point of view.

§7.1. Constrained connections on $M^{3|2}$

We specialize the discussion in §6.3 to $M^{3|2}$. Our first task is to evaluate (6.74). To that end we compute using the formulas of §2.3:

$$\begin{aligned} g_{\mu\nu}\tilde{\Gamma}^{\mu ac}\Gamma^{\nu}_{bd} &= g_{\mu\nu}\epsilon^{aa'}\epsilon^{cc'}\Gamma^{\mu}_{a'c'}\Gamma^{\nu}_{bd} \\ &= \frac{1}{2}(\epsilon_{a'b}\epsilon_{c'd} + \epsilon_{a'd}\epsilon_{c'b})(\epsilon^{aa'}\epsilon^{cc'} + \epsilon^{ac'}\epsilon^{ca'}) \\ &= \delta^a_b\delta^c_d + \delta^a_d\delta^c_b. \end{aligned} \tag{7.1}$$

Plugging into (6.74) we obtain

$$-2\tilde{\Gamma}^{\mu ac}\Gamma^{\nu}_{cb}\mathcal{F}_{\mu\nu} = 4\mathcal{D}_b\Lambda^a + \delta^a_b\mathcal{D}_c\Lambda^c. \tag{7.2}$$

Setting $a = b$ and summing we find

$$6\mathcal{D}_c\Lambda^c = -2\epsilon^{aa'}\epsilon^{cc'}\Gamma^{\mu}_{a'c'}\Gamma^{\nu}_{ac}\mathcal{F}_{\mu\nu} = 0, \tag{7.3}$$

since ϵ and $\mathcal{F}$ are skew-symmetric. Thus (7.2) becomes

$$\mathcal{D}_b\Lambda^a = -\frac{1}{2}\tilde{\Gamma}^{\mu ac}\Gamma^{\nu}_{cb}\mathcal{F}_{\mu\nu}. \tag{7.4}$$

This implies $\mathcal{E} = 0$ in (6.76), which means that there are no auxiliary fields.

The multiplet $\{A, \lambda\}$ of component fields is called the *vector multiplet*, and from the vector multiplet we can reconstruct the constrained connection $\mathcal{A}$. The precise statement is the following.

Proposition 7.5. *The category of connections $\mathcal{A}$ on $M^{3|2}$ whose curvature vanishes on the odd distribution τ is equivalent to the category of pairs $\{A, \lambda\}$ consisting of a connection A on $\check{M}^3$ and a dual spinor field λ with values in the adjoint bundle.*

We first comment on the meaning of Proposition 7.5. As odd fields are involved, it is crucial to work over a base S: for any supermanifold S, what Proposition 7.5 claims is that the functor which attaches the component fields to a constrained connection is an equivalence between the following categories: (a) G-torsors on $M^{3|2} \times S$ provided with a relative connection $\mathcal{A}$ satisfying the curvature constraint; (b) G-torsors on $\check{M}^3 \times S$ provided with a relative connection A and an adjoint bundle valued dual spinor field λ. Note that in (b) we have $\check{M}^3 \times S$, not the reduced space of $M^{3|2} \times S$. As usual, in the proof we will keep S silent.

If $\mathcal{P}$ is a G-torsor with a connection $\mathcal{A}$ on $M^{3|2}$ (i.e. on $M^{3|2} \times S$), any automorphism of $(\mathcal{P}, \mathcal{A})$ which induces the identity on the restriction P of $\mathcal{P}$ to $\check{M}^3$(i.e. $\check{M}^3 \times S$) is the identity. Because of this, $(\mathcal{P}, \mathcal{A})$ can be viewed as a structure on P. Concretely, once some choices have been made on $\check{M}^3$ (see the proof of Proposition 7.5 for an example), if $(\mathcal{P}, \mathcal{A})$ is a G-torsor with connection on $M^{3|2}$ with restriction P to $\check{M}^3$, and if s is a trivialization of P, the connection gives rise to a canonical extension of s to a trivialization of $\mathcal{P}$. Given s, then $\mathcal{A}$ becomes a 1-form α with values in the Lie algebra $\mathfrak{g}$ obeying suitable constraints. What Proposition 7.5 claims is that the map from the space of those α's to the space of (A, λ) (A a connection on P, and λ a $\mathfrak{g}$-valued dual spinor field) is bijective. In other words, for P a G-torsor on $\check{M}^3$ the map from (a) the set of isomorphism classes of G-torsors $\mathcal{P}$ on $M^{3|2}$, given with a constrained connection $\mathcal{A}$ and an isomorphism $P \xrightarrow{\sim} \mathcal{P} \mid \check{M}^3$, to (b) the set of (A, λ), is bijective. In (a), the objects considered have no non-trivial isomorphisms. Because of this, the question is local, and one may assume that P is the trivial G-torsor.

Proof. The data on $\mathcal{P}$ of a constrained connection amounts to that of a partial connection in the direction of the odd distribution τ: a G-invariant lift to $\mathcal{P}$ not of all vector fields, but only of those in τ. If $\mathcal{D}_a$ (resp. ∇_{ab}) is the horizontal lift of D_a (resp. ∂_{ab}), the vanishing along τ of F means that

$$\nabla_{ab} = -\frac{1}{2}[\mathcal{D}_a, \mathcal{D}_b]. \tag{7.6}$$

If only the $\mathcal{D}_a$ are given, the corresponding constrained connection is obtained by defining ∇_{ab} by this formula.

The functor implicit in Proposition 7.5 attaches to $(\mathcal{P}, \nabla)$: the restriction to $\check{M}^3$ of $\mathcal{P}$ (denoted P), that of $[\mathcal{D}_a, \mathcal{D}_b]$, and that of $[[\mathcal{D}_a, \mathcal{D}_b], \mathcal{D}_c]$; respectively: a G-torsor on $\check{M}^3$, a connection on it, and a section of the adjoint bundle with vanishing symmetrization in a, b, c. (cf. (6.68)).

We now choose a coordinate system as in §2.3, giving rise to the basis D_1, D_2 of τ. We will use D_1, D_2 to define a retraction q: $M^{3|2} \to \check{M}^3$, and, for $\mathcal{P}$ a torsor with connection on $M^{3|2}$, with restriction P to $\check{M}^3$, an isomorphism of torsors $q^*P \to \mathcal{P}$ (partial gauge fixing).

We first explain an analog, in classical differential geometry of what we will do. Suppose N is of codimension 2 in a variety M, and that D_1, D_2 are vector fields which, on N, span the normal bundle T_M/T_N. In a neighborhood of N, the flow $\exp(tD_1)$ generates from N a codimension 1 subvariety M_1, with a retraction

$q_1\colon M \to N$ constant on the flow lines. Repeating the process with M_1 and D_2, we obtain in a neighborhood of N a retraction $q_2\colon M \to M_1$, constant on the flow lines of D_2:



If $\mathcal{P}$ is a torsor on M with connection, with restriction P to N, integrating the connection on the flow lines of D_1, we obtain an isomorphism $\alpha_1\colon q_1^*P \to \mathcal{P}$ on M_1. Doing the same with D_2, we obtain $\alpha_2\colon q_2^*(\mathcal{P}|M_1)\tilde{\to}\mathcal{P}$.

In our case of codimension $0|2$, the situation is different in that (a) the vector field D_1 does not integrate to a foliation with leaves $\mathbb{R}^{1|0}$: the obstruction is D_1^2 mod D_1. (b) On $\mathbb{R}^{0|1}$, a connection ∇ does not need to be flat: curvature obstruction ∇_θ^2.

It however remains true that for $N \subset M^{3|2}$ a subvariety and D an odd vector field transversal to N, the vector field D, in fact just its restriction to N, generates from N an embedding $N \times \mathbb{R}^{0|1} \hookrightarrow M^{3|2}$. A function f on $M^{3|2}$ vanishes on the image M_1 if and only if f and Df vanish on N, and the retraction $q_1\colon M_1 \to N$ is such that for any function u on N, $Dq_1^*(u) = 0$. If $\mathcal{P}$ is a torsor with connection on $M^{3|2}$, with restriction P to N, we have on M_1 a unique isomorphism of torsors $q_1^*P \to \mathcal{P}|M_1$ which is the identity on N and such that on N the liftings of D agree. One should beware that if D is given as a vector field on $M^{3|2}$, it does not need to be tangent to M_1. This tangency to M_1 holds only on N.

We apply this twice, to $\check{M}^3 \subset M^{3|2}$ and D_1, then to M_1 and D_2, obtaining $q\colon M^{3|2} \to M_1 \to \check{M}^3$ and $\alpha\colon q^*P \to \mathcal{P}$. The construction of α uses only the connection in the directions of D_1 and D_2: for $\mathcal{P}$ a torsor with a connection along τ, we obtain $\alpha\colon q^*P \to \mathcal{P}$, and the data of α is equivalent to that of part of the connection, viz. $\mathcal{D}_1$ on $\check{M}^3$ and $\mathcal{D}_2$ on M_1. It remains to check that to complete this partial data to that of a connection ∇ along τ amounts to giving $[\mathcal{D}_a, \mathcal{D}_b]$ on $\check{M}^3$ (a connection on P) and $[[\mathcal{D}_a, \mathcal{D}_b], \mathcal{D}_c]$ on $\check{M}^3$ (a section on $\check{M}^3$ of $\mathrm{ad}\, P$ with vanishing symmetrization in a, b, c, i.e., the data of just $[[\mathcal{D}_1, \mathcal{D}_1], \mathcal{D}_2]$ and $[[\mathcal{D}_2, \mathcal{D}_2], \mathcal{D}_1]$).

Let us choose a local trivialization of P, hence of $\mathcal{P}$. The connection along τ is given by a section a of τ^* with values in the Lie algebra, i.e., by $a_1 = \langle D_1, a\rangle$ and $a_2 = \langle D_2, a\rangle$. That the connection is partially given by $\alpha\colon q^*P \to \mathcal{P}$ amounts to $a_1 = 0$ on $\check{M}^3$ and $a_2 = 0$ on M_1, i.e. to $a_1 = a_2 = D_1a_2 = 0$ on $\check{M}^3$.

For the simplicity of notations, let us assume that G is $GL(n)$, i.e., that we are considering a vector bundle. We have then

$$\begin{aligned} [\mathcal{D}_1, \mathcal{D}_1] &= -2\partial_{11} + 2D_1a_1 \quad \text{on} \quad \check{M}^3 \\ [\mathcal{D}_1, \mathcal{D}_2] &= -2\partial_{12} + D_2a_1 \quad \text{on} \quad \check{M}^3 \\ [\mathcal{D}_2, \mathcal{D}_2] &= -2\partial_{22} + 2D_2a_1 \quad \text{on} \quad \check{M}^3\ , \end{aligned} \tag{7.7}$$

so the data of the first order jet of a_1 and a_2 along $\check{M}^3$ amounts to the data of the connection on P. The remaining freedom: the addition of $\theta_1\theta_2b_1$ to a_1 and $\theta_1\theta_2b_2$ to a_2 amounts to the data of $[[\mathcal{D}_i, \mathcal{D}_j], \mathcal{D}_k]$ on $\check{M}^3$. Indeed, by $a_i \mapsto a_i + \theta_1\theta_2b_i$, $[[\mathcal{D}_1, \mathcal{D}_1], \mathcal{D}_2]$ on $\check{M}^3$ is increased by $-2b_1$ and $[[\mathcal{D}_2, \mathcal{D}_2], \mathcal{D}_1]$ by $-2b_2$.

It is easy in this case to reformulate a constrained connection in terms of an *unconstrained* field. Namely, take the unconstrained field to be a partial connection on $\mathcal{P}$ along the odd distribution τ. Then if $\mathcal{D}_a$ is the horizontal lift of D_a, define the horizontal lift ∇_{ab} of ∂_{ab} to be

$$\nabla_{ab} = -\frac{1}{2}[\mathcal{D}_a, \mathcal{D}_b]. \tag{7.6}$$

In other words, the constraint that the odd-odd components of the curvature vanish uniquely determines the connection $\mathcal{A}$ from the restriction of $\mathcal{A}$ to τ.

It is convenient to lower an index on the spinor field and define[37]

$$\boxed{\Lambda_a = \Lambda^b \epsilon_{ba} = \frac{1}{3}\epsilon^{bc}[\nabla_{ab}, \mathcal{D}_c].} \tag{7.8}$$

Equation (6.73) specializes to

$$[\mathcal{D}_a, \Lambda_{bc}] = \epsilon_{ab}\Lambda_c + \epsilon_{ac}\Lambda_b. \tag{7.9}$$

Note that (7.3) is now

$$\epsilon^{ab}\mathcal{D}_a\Lambda_b = 0. \tag{7.10}$$

We note a useful consequence of (7.10):

$$\mathcal{D}^2\Lambda_a = -\epsilon^{bc}\nabla_{ab}\Lambda_c. \tag{7.11}$$

It is convenient to set

$$f_{ac} = \frac{1}{2}\epsilon^{bd}F_{(ab)(cd)}, \tag{7.12}$$

which is symmetric in a, c. Then

$$*F_A = f_{ab}dy^{ab} \tag{7.13}$$

is the Hodge dual of the curvature (c.f. (2.38)).

In these variables the component fields are

$$\boxed{\begin{array}{c} A = i^*\mathcal{A} \\ \lambda_a = i^*\Lambda_a \end{array}} \tag{7.14}$$

and from (7.3) and (7.10) we have

$$\boxed{f_{ab} = i^*\mathcal{D}_a\Lambda_b = i^*\mathcal{D}_b\Lambda_a.} \tag{7.15}$$

The supersymmetry transformation laws (6.8) specialize to

$$\boxed{\begin{array}{l} (\hat{\xi}A)_{ab} = -\eta^c(\epsilon_{ac}\,\lambda_b + \epsilon_{bc}\,\lambda_a) \\ \quad \hat{\xi}\lambda_a = -\eta^b f_{ab} \end{array}} \tag{7.16}$$

and (6.9) specializes to

$$\boxed{\hat{\xi}f_{ab} = \quad \eta^c\big[\nabla_{ca}\lambda_b + \epsilon_{ac}(\not{D}\lambda)_b\big].} \tag{7.17}$$

[37] This goes against our general wish not to raise or lower a single index using a skew-symmetric tensor, since there is a sign ambiguity. However, defining Λ_a facilitates comparison with the physics literature.

§7.2. The Yang-Mills action on $M^{3|2}$

Assume that G is compact. Fix a bi-invariant inner product $\langle\cdot,\cdot\rangle$ on $\mathfrak{g}$. For a constrained connection $\mathcal{A}$ on $M^{3|2}$ the basic lagrangian density is

$$\boxed{L_2 = |d^3x|\, d^2\theta \, \Big\{-\frac{1}{4}\epsilon^{ab}\langle\Lambda_a, \Lambda_b\rangle\Big\}.} \tag{7.18}$$

We compute the component lagrangian function $\check{L}_2$ using (2.40):

$$\begin{aligned} -4\check{L}_2 &= -\epsilon^{ab} i^* D^2\langle\Lambda_a, \Lambda_b\rangle \\ &= -\epsilon^{ab} i^*\Big\{2\langle\mathcal{D}^2\Lambda_a, \Lambda_b\rangle + \epsilon^{cd}\langle\mathcal{D}_d\Lambda_a, \mathcal{D}_c\Lambda_b\rangle\Big\} \\ &= 2\epsilon^{ab}\epsilon^{cd}\langle\nabla_{ac}\lambda_d, \lambda_b\rangle - \epsilon^{ab}\epsilon^{cd}\langle f_{da}, f_{cb}\rangle. \end{aligned} \tag{7.19}$$

In the last step we use (7.15), (7.11), and (7.14). So from (2.27), (2.28), and the fact that $|{*F_A}|^2 = |F_A|^2$ we conclude

$$\boxed{\check{L}_2 = -\frac{1}{2}|F_A|^2 + \frac{1}{2}\langle\lambda \not{D}_A \lambda\rangle.} \tag{7.20}$$

This is the super Yang-Mills lagrangian (6.3). Thus we do have a superspace formulation of this lagrangian in 3 dimensions.

Next, we begin the analysis of the superspace lagrangian directly in superspace. This is a redundant exercise, since we already have the component expressions in (6.22) and the equations which follow. But it is instructive to see how the computation looks in superspace.

We compute the superspace equation of motion by varying the *unconstrained* partial connection $\mathcal{D}_a$. Notice that a variation $\delta\mathcal{D}_a$ is a G-invariant vertical vector field, so corresponds to a section $\delta\mathcal{A}_a$ of the adjoint bundle. From (7.8), (6.66), and (7.6) we compute

$$\delta\Lambda_a = -\frac{1}{6}\epsilon^{cd}\Big\{[[\delta\mathcal{D}_a, \mathcal{D}_c], \mathcal{D}_d] + [[\mathcal{D}_a, \delta\mathcal{D}_c], \mathcal{D}_d] - 2[\nabla_{ac}, \delta\mathcal{D}_d]\Big\}. \tag{7.21}$$

We use the invariance of $\langle\cdot,\cdot\rangle$, (7.10), and (7.11) to find

(7.22)

$$\begin{aligned} \delta L_2 &= |d^3x|\, d^2\theta \, \frac{-1}{2}\epsilon^{ab}\langle\delta\Lambda_a, \Lambda_b\rangle \\ &= |d^3x|\, d^2\theta \, \frac{1}{12}\epsilon^{ab}\epsilon^{cd}\Big\{\langle\delta\mathcal{D}_a, [\mathcal{D}_c, [\mathcal{D}_d, \Lambda_b]]\rangle + \langle\delta\mathcal{D}_c, [\mathcal{D}_a, [\mathcal{D}_d, \Lambda_b]]\rangle \\ &\qquad\qquad + 2\langle\delta\mathcal{D}_d, [\nabla_{ac}, \Lambda_b]\rangle\Big\} + EXACT \\ &= |d^3x|\, d^2\theta \, \frac{-1}{2}\epsilon^{ab}\epsilon^{cd}\langle\delta\mathcal{A}_a, \nabla_{bc}\Lambda_d\rangle + EXACT. \end{aligned}$$

So the superspace equation of motion is

$$\boxed{\epsilon^{cd}\nabla_{bc}\Lambda_d = 0 \qquad \text{for all } b.} \tag{7.23}$$

The exact term is $-d\gamma$ for the variational 1-form γ. We do not compute the precise expression, since it is unwieldy and not useful. One interesting feature emerges quickly, though. Namely, γ contains a term proportional to

$$\epsilon^{ab}\epsilon^{cd}\iota(D_d)\ |d^3x|\,d^2\theta\ D_c\langle\delta\mathcal{A}_a,\Lambda_b\rangle, \tag{7.24}$$

and so as a differential 1-form on the space of fields γ is *not* linear over functions. (In other words, $\delta\mathcal{A}_a$ is differentiated.)

We extract from (7.23) the equations of motion in components. Restricting (7.23) to $\check{M}^3$ we find

$$\boxed{\not{D}_A\lambda = 0.} \tag{7.25}$$

This agrees with (6.25). To obtain the equation of motion for A, we either apply $i^*\mathcal{D}_e$ to (7.23) or specialize (6.23). In either case we obtain

$$\boxed{\epsilon^{cd}\nabla_{ad}f_{bc} = [\lambda_a,\lambda_b] \qquad \text{for all } a,b.} \tag{7.26}$$

This agrees with (6.23). Finally, the supercurrent (6.27) specializes to

$$\boxed{j_a = \frac{1}{2}\epsilon^{cb}\epsilon^{de}\langle f_{cd},\lambda_e\rangle\ \frac{\partial}{\partial y^{ab}}\otimes|d^3y| + \frac{1}{2}\epsilon^{cd}\epsilon^{be}\langle f_{ca},\lambda_e\rangle\ \frac{\partial}{\partial y^{db}}\otimes|d^3y|.} \tag{7.27}$$

Since we didn't compute the exact term in (7.22), we do not give a superspace computation of the supercurrent.

§7.3. Gauge theory with matter on $M^{3|2}$

In this subsection we prove Theorem 6.33(i). Suppose we are given the data in (6.35): a Riemannian manifold X on which G acts by isometries and a G-invariant function $h\colon X\to\mathbb{R}$. Fix a G bundle $\mathcal{P}\to M^{3|2}$. Then we can form the associated bundle $X^{\mathcal{P}}\to M^{3|2}$ whose typical fiber is X. The superfields in the theory are a constrained connection $\mathcal{A}$ on $\mathcal{P}$ and a section Φ of $X^{\mathcal{P}}$. We view Φ as an equivariant map

$$\Phi\colon \mathcal{P}\longrightarrow X. \tag{7.28}$$

Then the formulas in §4 make sense provided we interpret the vector fields D_a,∂_{ab} as the horizontal lifts to $\mathcal{P}$ of the indicated vector fields on $M^{3|2}$. For example, the component fields are still defined by (4.7), but now $\psi_a = i^*\mathcal{D}_a\Phi$ where $\mathcal{D}_a$ acts by the covariant derivative associated to $\mathcal{A}$. Also, in the definition of the auxiliary field $F = \frac{1}{2}i^*(\mathcal{D}_2\mathcal{D}_1 - \mathcal{D}_1\mathcal{D}_2)\Phi$ the outer covariant derivatives are constructed from both $\mathcal{A}$ and the Levi-Civita connection on X.

The superspace lagrangian for the theory is $L_0+L_1+L_2$, where the individual terms are defined in (4.16), (4.20), and (7.18). The only possible changes to the computation of the component lagrangian are additional curvature terms from the connection $\mathcal{A}$. Recall that $\mathcal{A}$ is constrained to be flat on the odd distribution τ, and this constraint eliminates many such possible terms. With this in mind we examine the computation (4.18). There is a change in the formula (4.11) for $\mathcal{D}^2\mathcal{D}_b\Phi$, which

is used to pass from the second line of (4.18) to the third line. Examining (4.12) we see that there is a new term from the even-odd curvature of $\mathcal{A}$ in passing to the last line. Hence (4.11) is replaced by[38]

$$\mathcal{D}^2\mathcal{D}_a\Phi = \epsilon^{bc}\big(\nabla_{ab} - \frac{1}{6}R(\mathcal{D}_a\Phi, \mathcal{D}_b\Phi)\big)\mathcal{D}_c\Phi + 2(\Lambda_a)_\Phi, \tag{7.29}$$

where $(\Lambda_a)_\Phi$ is the vector field on $X^{\mathcal{P}}$ induced by the action of Λ_a on X. The new term in (7.29) contributes an extra term to (4.18), and the new contribution to the component lagrangian is

$$-\epsilon^{ab}\langle\psi_a, (\hat{\lambda}_b)_\phi\rangle = -\langle\psi_a, \hat{\lambda}^a_\phi\rangle = \langle\hat{\lambda}^a_\phi, \psi_a\rangle. \tag{7.30}$$

The computation (4.21) with the potential term remains unchanged. Altogether, we find the total component lagrangian to be (6.36), and this completes the proof of Theorem 6.33(i).

[38] Recall a tricky sign in [I-**Classical Fields**, §4.2]: An equivariant vertical vector field is identified with a section of the adjoint bundle, and the action on a section of an associated bundle is by *minus* the directional derivative.

CHAPTER 8
$N = 1$ Yang-Mills Theory in Dimension 4

Here we complete the discussion initiated in §6.3 of the supersymmetric formulation of super Yang-Mills theory in dimension 4. This chapter parallels the previous one. One new feature in §8.1 is the appearance of an auxiliary field as well as the existence of an R-symmetry. Another new feature appears in §8.2, where the coupling constant of the Yang-Mills lagrangian is complex. Its imaginary part is proportional to the usual coupling constant, whereas its real part is the coefficient of a Chern-Weil form. (If this term is present, then the theory is not invariant under orientation-reversing symmetries.) In §8.3 we describe the superspace approach to gauge theories with matter in case the target manifold of the scalar field is Hodge. Then there is a geometric explanation for the appearance of the moment map in the component lagrangian (6.42).

§8.1. Constrained connections on $M^{4|4}$

We specialize the discussion in §6.3 to $M^{4|4}$. As explained in §2.4 we work in a complex basis. Our first task is to evaluate (6.74). To that end we compute

$$\begin{aligned} g_{\mu\nu}\tilde{\Gamma}^{\mu a\dot{c}}\Gamma^{\nu}_{b\dot{d}} &= g_{\mu\nu}\epsilon^{aa'}\epsilon^{\dot{c}\dot{c}'}\Gamma^{\mu}_{a'\dot{c}'}\Gamma^{\nu}_{b\dot{d}} \\ &= \frac{1}{2}\epsilon_{a'b}\epsilon_{\dot{c}'\dot{d}}\epsilon^{aa'}\epsilon^{\dot{c}\dot{c}'} \\ &= 2\delta^a_b\delta^{\dot{c}}_{\dot{d}} \end{aligned} \tag{8.1}$$

Plugging into (6.74) we obtain

$$\mathcal{D}_b\Lambda^a = -\frac{1}{2}\tilde{\Gamma}^{\mu ac}\Gamma^{\nu}_{cb}\mathcal{F}_{\mu\nu} - \frac{1}{2}\delta^a_b\overline{\mathcal{D}}_{\dot{c}}\overline{\Lambda}^{\dot{c}}. \tag{8.2}$$

Thus there is an auxiliary field in this theory. Setting $a = b$ and summing, the first term on the right hand side of (8.2) vanishes, and so

$$\mathcal{D}_c\Lambda^c = -\overline{\mathcal{D}}_{\dot{c}}\overline{\Lambda}^{\dot{c}}. \tag{8.3}$$

Define the (auxiliary) field

$$E = -\frac{\sqrt{-1}}{2}\mathcal{D}_c\Lambda^c. \tag{8.4}$$

Equation (8.3) asserts that E is real:

$$E = \overline{E}. \tag{8.5}$$

From (6.74) we also learn that

$$\overline{\mathcal{D}}_{\dot{b}}\Lambda^a = 0; \tag{8.6}$$

in other words, Λ^a is a chiral superfield.

Let A, λ, E be the restrictions of $\mathcal{A}, \Lambda, E$ to $\check{M}^4$. (It should cause no confusion that the restriction of E to $\check{M}^4$ is also denoted 'E'.) The multiplet $\{A, \lambda, E\}$ of component fields is called the *vector multiplet*, and from the vector multiplet we can reconstruct the constrained connection $\mathcal{A}$. The precise statement is the following.

Proposition 8.7. *The category of connections $\mathcal{A}$ on $M^{4|4}$ whose curvature vanishes on the odd distribution τ is equivalent to the category of triples $\{A, \lambda, E\}$ consisting of a connection A on $\check{M}^4$, a dual real spinor field λ with values in the adjoint bundle, and a section E of the adjoint bundle.*

For an elucidation of what Proposition 8.7 claims, we refer to the comments following Proposition 7.5. The following proof contains a description of an unconstrained field equivalent to a constrained connection.

Proof. We first consider the complex analog of Proposition 8.7. For the complex super Minkowski space $M_{\mathbb{C}} = M_{\mathbb{C}}^{4|4}$, instead of a complex structure on the distribution τ, we get a decomposition $\tau = \tau' \oplus \tau''$, and the distributions τ' and τ'' are integrable. Dividing by the corresponding foliations, we obtain

$$M'' \overset{\mathrm{pr}''}{\leftarrow} M_{\mathbb{C}} \overset{\mathrm{pr}'}{\rightarrow} M', \tag{8.8}$$

with τ' (resp τ'') being the relative tangent bundle of $M_{\mathbb{C}}/M''$ (resp $M_{\mathbb{C}}/M'$). Let $\check{M}_{\mathbb{C}} \subset M_{\mathbb{C}}$ be the ordinary Minkowski space, with images $\check{M}'$ in M' and $\check{M}''$ in M''.

Put $M_1 = (\mathrm{pr}'')^{-1}(\check{M}'')$ and $M_2 := (\mathrm{pr}')^{-1}(\check{M}')$. The projections pr' and pr'' induce isomorphisms

$$M_1 \overset{\sim}{\rightarrow} M' \tag{8.9}$$

and

$$M_2 \overset{\sim}{\rightarrow} M''. \tag{8.10}$$



If P is a G-torsor on $M_{\mathbb{C}}$, a constrained connection on $\mathcal{P}$ is determined by its restriction to τ and, arguing as in the proof of Proposition 7.5, one sees that constrained connections correspond one to one to connections in the direction of τ, which are integrable in the directions of τ' and τ''. An integrable connection in the direction of τ'' (resp τ') amounts to a way to descend $\mathcal{P}$ from $M_{\mathbb{C}}$ to M' (resp M''). A constrained connection hence amounts to the data of $\mathcal{P}'$ on M' and $\mathcal{P}''$ on M'' descending $\mathcal{P}$:

$$\mathcal{P} \xrightarrow{\sim} (\mathrm{pr}')^{-1}(\mathcal{P}') \tag{8.11}$$

and

$$\mathcal{P} \xrightarrow{\sim} (\mathrm{pr}'')^{-1}(\mathcal{P}''). \tag{8.12}$$

In the τ' (resp. τ'') direction, the connection is given by (8.12) (resp. (8.11)).

Let P be the restriction of $\mathcal{P}$ to $\check{M}_{\mathbb{C}}$, and let us choose a trivialization of P. From (8.12) (resp (8.11)) we get a trivialization of $\mathcal{P}$ on M_1 (resp M_2): the only one extending the given trivialization and compatible with the connection along τ' (resp τ''). From (8.9) (resp (8.10)) and (8.11) (resp (8.12)), we then get trivializations s' and s'' of $\mathcal{P}'$ and $\mathcal{P}''$.

We keep noting s' and s'' their inverse images in $\mathcal{P}$ and let $g\colon M_{\mathbb{C}} \to G$ be the map such that $s'g = s''$. The map g to G, trivial on M_1 and M_2, is an unconstrained field equivalent to the constrained connection. It can be written $g = \exp(U)$, with U a map to $\mathfrak{g} = \mathrm{Lie}(G)$ vanishing on M_1 and M_2. In terms of g, the data $(\mathcal{P}, \mathcal{A})$ is the trivial torsor, with the trivial connection in the direction of τ'' and, in the direction of τ', the connection for which g is horizontal (given by the 1-form $-g^{-1}dg$).

Let us use coordinates as in §2.4. In those coordinates, M_1 is defined by $\bar{\theta}^{\dot{b}} = 0$, and M_2 by $\theta^a = 0$. Let J be the ideal of functions vanishing on M_1 and M_2, and I be the ideal generated by all θ and $\bar{\theta}$. Then, $(J \cap I^n)/J \cap I^{n+1}$ vanishes for $n \neq 2, 3, 4$ and is spanned respectively by

$$\begin{aligned} n = 2 &: \text{ the } \theta^a \bar{\theta}^{\dot{b}} \\ n = 3 &: \text{ the } \theta^1\theta^2\bar{\theta}^{\dot{b}} \text{ and the } \theta^a\bar{\theta}^{\dot{1}}\bar{\theta}^{\dot{2}} \\ n = 4 &: \ \theta^1\theta^2\bar{\theta}^{\dot{1}}\bar{\theta}^{\dot{2}}, \end{aligned}$$

the coefficients being functions on $\check{M}_{\mathbb{C}}$. We have

$$\begin{aligned} \mathcal{D}_a &= D_a, \\ \mathcal{D}_{\dot{a}} &= D_a - g^{-1}D_a g. \end{aligned} \tag{8.13}$$

It is then easy to check that $U \pmod{I}^3$ gives the connection on P; that changing U by a section of $(J \cap I^3) \otimes \mathfrak{g}$, taken modulo $J \cap I^4$, gives λ; and changing U by a section of $(J \cap I^4) \otimes \mathfrak{g}$ gives E.

We now consider the real case. The arguments used in the complex case work in just the same way for the cs case, where super Minkowski space is complexified just in the odd direction. As explained in the comments after Proposition 7.5, we have to prove that some map is a bijection. This map is obtained from the

similar map in the cs case by taking the fixed points of complex conjugation, and Proposition 8.7 follows.

In the physics literature the field U is referred to as the vector superfield in "Wess-Zumino gauge".

It is convenient to lower an index on the spinor field and define the chiral superfield[39]

$$\boxed{W_a = \Lambda^b \epsilon_{ba} = \frac{1}{4}\epsilon^{\dot{b}\dot{c}}[\nabla_{a\dot{b}}, \overline{\mathcal{D}}_{\dot{c}}]} \tag{8.14}$$

together with the conjugate antichiral superfield $\overline{W}_{\dot{a}}$. Equation (6.73) specializes to

$$[\mathcal{D}_a, \nabla_{b\dot{c}}] = 2\epsilon_{ab}\overline{W}_{\dot{c}}. \tag{8.15}$$

The auxiliary field E is

$$E = -\frac{\sqrt{-1}}{2}\epsilon^{ab} i^* \mathcal{D}_a W_b. \tag{8.16}$$

The reality condition (8.5) (equivalently (8.3)) is

$$\epsilon^{ab}\mathcal{D}_a W_b + \epsilon^{\dot{a}\dot{b}}\overline{\mathcal{D}}_{\dot{a}}\overline{W}_{\dot{b}} = 0 \tag{8.17}$$

This has the useful consequence

$$\mathcal{D}^2 W_a = -2\epsilon^{\dot{b}\dot{c}}\nabla_{a\dot{b}}\overline{W}_{\dot{c}}. \tag{8.18}$$

It is convenient to write the curvature as

$$F_A := \frac{1}{2}(F_{ab}\epsilon_{\dot{c}\dot{d}} + \overline{F}_{\dot{c}\dot{d}}\epsilon_{ab})\, dy^{a\dot{c}} \wedge dy^{b\dot{d}}, \tag{8.19}$$

where F_{ab} is symmetric in a, b. Our convention is that the curvature is real—$\overline{F}_{\dot{a}\dot{b}} = \overline{F_{ab}}$—for a compact gauge group G. *Warning*: The involution on the complexified Lie algebra $\mathfrak{g}_{\mathbb{C}}$ is such that the real subalgebra is the Lie algebra of the compact group. Thus for $G = \mathbb{T}$ the circle group the real subalgebra in $\mathfrak{g}_{\mathbb{C}} = \mathbb{C}$ is $i\mathbb{R} \subset \mathbb{C}$.

In these variables the component fields are

$$\boxed{\begin{array}{c} A = i^*\mathcal{A} \\ \lambda_a = i^* W_a, \end{array}} \tag{8.20}$$

and from (8.2) and (8.16) we have

$$\boxed{F_{ab} + \sqrt{-1}\,\epsilon_{ab} E = i^* \mathcal{D}_a W_b.} \tag{8.21}$$

[39]The symbol 'W' is standard in the physics literature.

Let $\hat{\xi}$ be the vector field on the space of component fields corresponding to the supersymmetry transformation $\eta^a Q_a + \bar{\eta}^{\dot{a}} \overline{Q}_{\dot{a}}$. We compute it from the general formula (1.29) and the other formulas in this section. We also can check the result against (6.8) and (6.9) if we set the auxiliary field E to zero. The result is:

$$\boxed{\begin{aligned} (\hat{\xi}A)_{a\dot{b}} &= -2\eta^c\,\epsilon_{ac}\,\bar{\lambda}_{\dot{b}} - 2\bar{\eta}^{\dot{c}}\,\epsilon_{\dot{b}\dot{c}}\,\lambda_a \\ \hat{\xi}\lambda_a &= -\eta^b F_{ab} + \eta^b \epsilon_{ab}\sqrt{-1}E \\ \hat{\xi}E &= \quad \eta^a\sqrt{-1}(\not{D}\bar{\lambda})_a - \bar{\eta}^{\dot{a}}\sqrt{-1}\,(\not{D}\lambda)_{\dot{a}} \end{aligned}} \tag{8.22}$$

and

$$\boxed{\hat{\xi}F_{ab} = \bar{\eta}^{\dot{c}}\big\{\nabla_{a\dot{c}}\lambda_b + \nabla_{b\dot{c}}\lambda_a\big\} + \eta^c\big\{\epsilon_{ac}(\not{D}\bar{\lambda})_b + \epsilon_{bc}(\not{D}\bar{\lambda})_a\big\}.} \tag{8.23}$$

Under the R-symmetry (2.81) the component fields transform by (see (8.14) and (8.16))

$$\boxed{\begin{aligned} [R, A_{a\dot{b}}] &= 0 \\ [R, \lambda_a] &= -\lambda_a \\ [R, \bar{\lambda}_{\dot{a}}] &= \bar{\lambda}_{\dot{a}} \\ [R, E] &= 0. \end{aligned}} \tag{8.24}$$

§8.2. The Yang-Mills action on $M^{4|4}$

We assume G compact with $\langle\cdot,\cdot\rangle$ a bi-invariant inner product on $\mathfrak{g}$. Extend $\langle\cdot,\cdot\rangle$ to be a *bilinear* form on $\mathfrak{g}_{\mathbb{C}}$. Fix $\tau \in \mathbb{C}$. Then for a constrained connection on $M^{4|4}$ the basic lagrangian density is

$$\boxed{\begin{aligned} \mathcal{L}_2 &= |d^4x|\,\Big\{-\frac{1}{16\pi}\,\mathrm{Im}\Big(d^2\theta\,\tau\,\epsilon^{ab}\langle W_a, W_b\rangle\Big)\Big\} \\ &= |d^4x|\,\frac{1}{2i}\Big\{d^2\theta\,\frac{-\tau}{16\pi}\epsilon^{ab}\langle W_a, W_b\rangle + d^2\bar{\theta}\,\frac{\bar{\tau}}{16\pi}\epsilon^{\dot{a}\dot{b}}\langle \overline{W}_{\dot{a}}, \overline{W}_{\dot{b}}\rangle\Big\}. \end{aligned}} \tag{8.25}$$

Notice that the two terms in $\mathcal{L}_2$ are defined on different cs-manifolds. We compute the contribution of the first to the component lagrangian using (8.20) and (8.21):

(8.26)

$$\begin{aligned} \frac{\tau}{16\pi}\epsilon^{ab} i^* D^2\langle W_a, W_b\rangle &= \frac{\tau}{16\pi}\epsilon^{ab} i^*\Big\{2\langle \mathcal{D}^2 W_a, W_b\rangle + \epsilon^{cd}\langle \mathcal{D}_d W_a, \mathcal{D}_c W_b\rangle\Big\} \\ &= -\frac{\tau}{4\pi}\epsilon^{ab}\epsilon^{\dot{c}\dot{d}} i^*\langle \nabla_{a\dot{c}}\overline{W}_{\dot{d}}, W_b\rangle \\ &\qquad + \frac{\tau}{16\pi}\epsilon^{ab}\epsilon^{cd}\langle F_{da} + \epsilon_{da}\sqrt{-1}E, F_{cb} + \epsilon_{cb}\sqrt{-1}E\rangle \\ &= \frac{\tau}{4\pi}\langle \bar{\lambda}\ \not{D}_A\ \lambda\rangle - \frac{\tau}{16\pi}\epsilon^{ab}\epsilon^{cd}\langle F_{ac}, F_{bd}\rangle + \frac{\tau}{8\pi}|E|^2. \end{aligned}$$

The contribution of the second term in (8.25) is similar and we combine them using (2.68). We fix the standard orientation on $\check{M}^4$ to write the result in terms of differential forms:

$$d^4x\ \check{\mathcal{L}}_2 = d^4x\,\Big\{-\frac{\mathrm{Im}\,\tau}{8\pi}|F_A|^2 + \frac{\mathrm{Im}\,\tau}{4\pi}\langle\bar{\lambda}\ \not{D}_A\ \lambda\rangle + \frac{\mathrm{Im}\,\tau}{8\pi}|E|^2\Big\} + \frac{\mathrm{Re}\,\tau}{8\pi}\langle F\wedge F\rangle. \tag{8.27}$$

Introduce the notation

$$\boxed{\tau = \frac{\theta}{2\pi} + \frac{4\pi i}{g^2}.} \tag{8.28}$$

Then

$$\boxed{d^4x\ \check{\mathcal{L}}_2 = d^4x\ \frac{1}{g^2}\left\{-\frac{1}{2}|F_A|^2 + \langle\bar{\lambda}\ \not{D}_A\ \lambda\rangle + \frac{1}{2}|E|^2\right\} + \frac{\theta}{16\pi^2}\langle F_A \wedge F_A\rangle.} \tag{8.29}$$

This is essentially the super Yang-Mills lagrangian (6.3). We have written the second term in complex notation and dropped an exact term along the way. Note E enters as an auxiliary field; its equation of motion is $E = 0$. The last term is *topological*; the integral is a characteristic number (for connections of finite action). Note that the first term makes sense as a density whereas to integrate the topological term we need an orientation on $\check{M}^4$. Another way to express (8.29) is to write $F = F^+ + F^-$ in terms of its self-dual and anti-self-dual pieces (see (2.64)). Then the lagrangian function is

$$\boxed{\check{\mathcal{L}}_2 = -\frac{\tau}{4\pi}|F_A^+|^2 + \frac{\bar{\tau}}{4\pi}|F_A^-|^2 + \frac{\operatorname{Im}\tau}{4\pi}\langle\bar{\lambda}\ \not{D}_A\lambda\rangle + \frac{\operatorname{Im}\tau}{8\pi}|E|^2.} \tag{8.30}$$

The orientation on $\check{M}^4$ enters here in distinguishing self-dual from anti-self-dual.

We are following standard physics usage in our choice of constants. The closed 4-form $\langle F_A \wedge F_A\rangle$ represents a real characteristic class of $P \to \check{M}^4$; if the bilinear form $\langle\cdot,\cdot\rangle$ on $\mathfrak{g}$ is such that the real characteristic class $\frac{1}{16\pi^2}\langle F_A \wedge F_A\rangle$ comes from a class in $H^4(BG;\mathbb{Z})$, then θ and $\theta + 2\pi$ yield the same quantum theory.

Next, we consider the action of the R-symmetry (2.81). Now $d^2\theta$ has weight 2 under R and from (8.14) we see that W_a has weight -1. It follows that (8.25) is invariant under R. One can also use (8.24) to check that (8.29) is invariant under the R-symmetry. We remark that even though the classical theory has a $U(1)$ R-symmetry, in the quantum theory this is broken to a finite subgroup by an anomaly (which is computed using the index theorem).

We do not analyze this theory in superspace, but rather simply report the equations of motion:

$$\boxed{\begin{aligned}\epsilon^{cd}\nabla_{d\dot{b}}F_{ac} + \epsilon^{\dot{c}\dot{d}}\nabla_{a\dot{d}}F_{\dot{b}\dot{c}} &= [\lambda_a, \bar{\lambda}_{\dot{b}}]\\ \epsilon^{ab}\nabla_{a\dot{c}}\lambda_b &= 0\\ E &= 0\end{aligned}} \tag{8.31}$$

and the supercurrent j_a which corresponds to the supersymmetry transformation Q_a:

$$\boxed{\begin{aligned}j_a = \frac{1}{2}\epsilon^{cd}\epsilon^{\dot{b}\dot{e}}\langle F_{ca}, \bar{\lambda}_{\dot{e}}\rangle\ \frac{\partial}{\partial y^{d\dot{b}}} \otimes |d^4y|\\ + \left\{\frac{1}{2}\epsilon^{\dot{c}\dot{b}}\epsilon^{\dot{d}\dot{e}}\langle\overline{F}_{\dot{c}\dot{d}}, \bar{\lambda}_{\dot{e}}\rangle - \epsilon^{\dot{b}\dot{c}}\langle\sqrt{-1}\,E, \bar{\lambda}_{\dot{c}}\rangle\right\}\ \frac{\partial}{\partial y^{a\dot{b}}} \otimes |d^4y|.\end{aligned}} \tag{8.32}$$

In this formula we take $g = 1$, $\theta = 0$.

§8.3. Gauge theory with matter on $M^{4|4}$

In this subsection we prove Theorem 6.33(ii). Suppose we are given the data in (6.41), but with the additional proviso that X be Hodge. This assumption is only for exposition, as the computation we review is local and done for arbitrary Kähler manifolds. Let $L_\omega \to X$ be the Hodge line bundle, a hermitian line bundle with curvature $\sqrt{-1}\omega$. In this theory a compact Lie group G acts by isometries on X, and we assume given a lift of the G action to L_ω. This is roughly equivalent to choosing a moment map

$$\mu\colon X \longrightarrow \mathfrak{g}^* \tag{8.33}$$

for the action of G on X, and in fact defines a moment map by the formula

$$\mu(\zeta) = -\sqrt{-1}\,\mathrm{vert}_1(\hat{\hat{\zeta}}), \qquad \zeta \in \mathfrak{g}. \tag{8.34}$$

Here $\mu(\zeta)$ is the function on X obtained by pairing (8.33) with ζ; the vector field $\hat{\hat{\zeta}}$ on L_ω is defined by the lifted action; and 'vert_1' indicates the vertical part with respect to the connection on L_ω, taken at points of unit norm. It is easy to check that μ is a moment map. In other words

$$d\mu(\zeta) = -\iota(\hat{\zeta})\omega, \tag{8.35}$$

where $\hat{\zeta}$ is the vector field on X induced by the G action. In case X is Kähler but not Hodge, we must choose a moment map. We also fix a G-invariant holomorphic function $W\colon X \to \mathbb{C}$.

Fix a G bundle $\mathcal{P} \to M^{4|4}$. The superfields in the theory are a constrained connection $\mathcal{A}$ on $\mathcal{P}$ and a chiral section Φ of the associated bundle $X^{\mathcal{P}}$, which we view as an equivariant chiral map

$$\Phi\colon \mathcal{P} \longrightarrow X. \tag{8.36}$$

As in (5.21) there is a global lift to an equivariant chiral map

$$\tilde{\Phi}\colon \mathcal{P} \longrightarrow \Phi^* L_\omega. \tag{8.37}$$

Then the total superspace lagrangian for the theory is $\mathcal{L}_0 + \mathcal{L}_1 + \mathcal{L}_2$, where the individual terms are defined in (5.22), (5.32), and (8.25). We take the coupling constants in the gauge theory to be $g = 1$ and $\theta = 0$. Our task is to work out the expansion in component fields. Note that component fields are defined by (5.7), though now the vector fields which enter the definitions are the $\mathcal{D}_a$ on $\mathcal{P}$—the horizontal lifts of the D_a defined by the connection $\mathcal{A}$.

We re-examine the computation beginning in (5.25). Since this uses the alternative form (5.17) of the lagrangian in terms of a local Kähler potential K, which we assume is G-invariant, our first job is to express a moment map μ in terms of K. Let $\hat{\zeta}$ be the real vector field on X which corresponds to $\zeta \in \mathfrak{g}$. Then an easy computation from (8.35) shows

$$\langle \tilde{\mu}, \zeta \rangle = \mu(\zeta) = -\sqrt{-1}\,\hat{\zeta}^{1,0} \cdot K = \sqrt{-1}\,\hat{\zeta}^{0,1} \cdot K \tag{8.38}$$

defines a moment map, where $\hat\zeta = \hat\zeta^{1,0} + \hat\zeta^{0,1}$ is expressed as a sum of its $(1,0)$ and $(0,1)$ components and $\check\mu\colon X \to \mathfrak{g}$ is obtained from μ using the inner product on $\mathfrak{g}$.

There are several new terms in (5.25)–(5.27) from the nonzero bracket (8.15). First, in the third line of (5.25) there is a new contribution from the bracket term in (5.24):

(8.39)
$$\begin{aligned} i^*\Big(\frac14\epsilon^{ab}\epsilon^{\dot c\dot d}\overline{\mathcal D}_{\dot c}\Big\{2\iota([\mathcal D_b,\nabla_{a\dot d}])\Phi^*\partial K\Big\}\Big) \\ &= -2i^*\epsilon^{\dot c\dot d}\overline{\mathcal D}_{\dot c}\,\iota(\overline{W}_{\dot d})\Phi^*\partial K \\ &= -2i^*\epsilon^{\dot c\dot d}\iota\big([\overline{\mathcal D}_{\dot c},\overline{W}_{\dot d}]\big)\Phi^*\partial K + 2i^*\epsilon^{\dot c\dot d}\iota(\overline{\mathcal D}_{\dot c})\iota(\overline{W}_{\dot d})\Phi^*\bar\partial\partial K \\ &= 4i^*\sqrt{-1}\,\iota\big(-\hat E_\Phi\big)\partial K + 2i^*\epsilon^{\dot c\dot d}\iota(\overline{\mathcal D}_{\dot c}\Phi)\iota\big(-(\widehat{\overline{W}}_\Phi)_{\dot d}\big)\bar\partial\partial K \\ &= 4\langle\phi^*\check\mu, E\rangle + \sqrt2\,\epsilon^{\dot c\dot d}\langle\bar\psi_{\dot c},(\hat{\bar\lambda}_\phi)_{\dot d}\rangle, \end{aligned}$$

where $(\widehat{\overline{W}}_{\dot d})_\Phi$ is the vector field on $X^{\mathcal P}$ induced by $\overline{W}_{\dot d}$ and similarly for E. In the third line we use (5.23) and the fact that $\iota(\overline{\mathcal D}_{\dot c})\Phi^*\partial K$ is a G-invariant function on $\mathcal P$, so its derivative in the fiber direction $\overline{W}_{\dot d}$ vanishes; in the fourth the definition (8.16) of E and the fact that vertical vectors on $\mathcal P$ act with a minus sign (cf. the footnote in §7.3); and in the fifth (8.38) and (5.16), together with an extra minus sign due to the odd variables.[40] From (5.28) and (5.29) we see that the results of (5.25)–(5.27) enter the component lagrangian with a factor $1/4$ and are added to their complex conjugates. Hence (8.39) leads to a contribution

$$2\langle\phi^*\check\mu, E\rangle + \frac{\sqrt2}{4}\Big(\epsilon^{cd}\langle(\hat\lambda_\phi)_c,\psi_d\rangle + \epsilon^{\dot c\dot d}\langle\bar\psi_{\dot c},(\hat{\bar\lambda}_\phi)_{\dot d}\rangle\Big) \tag{8.41}$$

to the component lagrangian. Also, in the second line of (5.27) there is a new term

$$\epsilon^{ab}\epsilon^{\dot c\dot d}\Big\{-\mathcal D_a\,\iota([\overline{\mathcal D}_{\dot c},\nabla_{b\dot d}])\Phi^*\partial K\Big\} = -4\epsilon^{ab}\mathcal D_a\,\iota(W_b)\Phi^*\partial K, \tag{8.42}$$

which by a similar computation contributes

$$-4\langle\phi^*\check\mu, E\rangle - \frac{\sqrt2}{2}\Big(\epsilon^{cd}\langle(\hat\lambda_\phi)_c,\psi_d\rangle + \epsilon^{\dot c\dot d}\langle\bar\psi_{\dot c},(\hat{\bar\lambda}_\phi)_{\dot d}\rangle\Big) \tag{8.43}$$

to the component lagrangian.

As in §7.3 there is also a new term due to a revision of (5.9):

$$\overline{\mathcal D}_{\dot a}\mathcal D^2\Phi = -2\epsilon^{bc}\Big\{\nabla_{b\dot a}\Phi - \frac14 R(\mathcal D_b\Phi,\overline{\mathcal D}_{\dot a}\overline\Phi)\Big\}\mathcal D_c\Phi - 4(\widehat{\overline{W}}_{\dot a})_\Phi, \tag{8.44}$$

This enters the computation in (5.26), where we pick up

$$\frac14\epsilon^{\dot c\dot d}\langle\overline{\mathcal D}_{\dot d}\overline\Phi, 4(\widehat{\overline{W}}_{\dot c})_\Phi\rangle = -\epsilon^{\dot c\dot d}\langle\overline{\mathcal D}_{\dot c}\overline\Phi, (\widehat{\overline{W}}_{\dot d})_\Phi\rangle. \tag{8.45}$$

[40] For even vectors $\xi_1, \bar\xi_2$ of type $(1,0),(0,1)$ respectively, equation (5.16) asserts

$$2\iota(\xi_1)\iota(\bar\xi_2)\bar\partial\partial K = -2\iota(\bar\xi_2)\iota(\xi_1)\bar\partial\partial K = \langle\bar\xi_2,\xi_1\rangle. \tag{8.40}$$

Now substitute $\xi_i = \eta_i\psi_i$, where ζ_i are odd vectors and η_i are auxiliary odd parameters.

Another new term enters into the last line of (5.27) from commuting $\mathcal{D}_a$ past $\nabla_{b\dot{d}}$:

$$\epsilon^{ab}\epsilon^{\dot{c}\dot{d}}\Big\{\frac{1}{2}\langle\overline{\mathcal{D}}_{\dot{c}}\overline{\Phi},[\mathcal{D}_a,\nabla_{b\dot{d}}]\Phi\rangle\Big\} = -2\epsilon^{\dot{c}\dot{d}}\langle\overline{\mathcal{D}}_{\dot{c}}\overline{\Phi},(\widehat{\overline{W}}_{\dot{d}})_\Phi\rangle. \tag{8.46}$$

Taking into account the factor $1/4$ in (5.28), and adding (8.45) and (8.46) to their complex conjugates, we obtain a contribution to the component lagrangian

$$-\frac{3\sqrt{2}}{4}\Big(\epsilon^{cd}\langle(\hat{\lambda}_\phi)_c,\psi_d\rangle + \epsilon^{\dot{c}\dot{d}}\langle\bar{\psi}_{\dot{c}},(\hat{\bar{\lambda}}_\phi)_{\dot{d}}\rangle\Big). \tag{8.47}$$

Combining (8.41), (8.43), and (8.47) we obtain the total new contribution to the component lagrangian:

$$-2\langle\phi^*\check{\mu},E\rangle - \sqrt{2}\Big(\epsilon^{cd}\langle(\hat{\lambda}_\phi)_c,\psi_d\rangle + \epsilon^{\dot{c}\dot{d}}\langle\bar{\psi}_{\dot{c}},(\hat{\bar{\lambda}}_\phi)_{\dot{d}}\rangle\Big) \tag{8.48}$$

The second term appears in (6.42), after using (8.14).

To recover the potential energy term in (6.42) we must eliminate the auxiliary field E. The terms involving E in (8.29) and (8.48) are:

$$\frac{1}{2}|E|^2 - 2\langle\phi^*\check{\mu},E\rangle. \tag{8.49}$$

The classical equation for E is then

$$\boxed{E = 2\phi^*\check{\mu}}, \tag{8.50}$$

and eliminating E we obtain the potential term

$$-2\phi^*|\mu|^2. \tag{8.51}$$

This completes the proof of Theorem 6.33(ii). Note the substitution (8.50) which gives the last equation in (6.43) from the second equation in (8.22).

CHAPTER 9
$N = 2$ Yang-Mills in Dimension 2

The theory we study in this chapter is the reduction to two dimensions of the basic four-dimensional supersymmetric gauge theory studied in §8. Thus we begin in §9.1 with some general remarks about dimensional reduction of ordinary bosonic Yang-Mills. The crucial observation is that in the dimensionally reduced theory we have not only a connection, but also some scalar fields and a nontrivial potential energy. The dimensional reduction of a constrained connection on $M^{4|4}$, which is described in §9.2, may be considered as a connection on $M^{2|4}$ with constraints. But the constraints now allow some nonzero curvature along the odd distribution, unlike the constraint condition in the maximal dimension 4. Recall also that the dimensionally reduced theory has a bigger R-symmetry group. In §9.3 we derive a superspace expression for the Yang-Mills lagrangian.

§9.1. Dimensional reduction of bosonic Yang-Mills

Consider pure Yang-Mills theory on Minkowski space $\check{M}^n$ in n dimensions with gauge group G. Fix a bi-invariant inner product $\langle \cdot, \cdot \rangle$ on $\mathfrak{g}$. The field is a connection A on a principal G bundle $P \to \check{M}^n$, and the lagrangian is

$$L = |d^n x| \, \frac{-1}{2} |F_A|^2. \tag{9.1}$$

Let $x^0, x^1, \ldots, x^{n-1}$ be standard coordinates on $\check{M}^n$ so that the metric is

$$g = (dx^0)^2 - (dx^1)^2 - \cdots - (dx^{n-1})^2. \tag{9.2}$$

Let T be the group of translations in the direction of the last k basis vectors. Let p be the quotient map $p\colon \check{M}^n \to \check{M}^n/T$. The quotient $\check{M}^n/T$ is a Minkowski space of dimension $n-k$. If $\check{M}^{n-k} \subset \check{M}^n$ is defined by $x^{n-k} = \cdots = x^{n-1} = 0$, the map p identifies $\check{M}^{n-k}$ with $\check{M}^n/T$.

In the dimensional reduction of pure Yang-Mills theory, one considers: (a) a principal G bundle P on $\check{M}^n/T \cong \check{M}^{n-k}$, and (b) a T-invariant connection A on its pullback to $\check{M}^n$. Let a be the restriction of A to $\check{M}^{n-k}$. Being T-invariant, the connection A can be written as

$$A = p^*(a) + \sum_{n-k}^{n-1} p^*(\phi_i) dx^i, \tag{9.3}$$

the ϕ_i being sections of the adjoint bundle.

The lagrangian density $\mathcal{L}_n$ on $\check{M}^n$ for A is T-invariant. In the reduced theory, the lagrangian density $\mathcal{L}_{n-k}$ on $\check{M}^{n-k}$ is obtained by restricting $\mathcal{L}_n$ to $\check{M}^{n-k}$ and contracting it against $\partial_{n-k}, \dots, \partial_{n-1}$: for Λ a lattice in T,

$$\int_{\check{M}^{n-k}} \mathcal{L}_{n-k} = \mathrm{vol}(T/\Lambda)^{-1} \int_{\check{M}^n/\Lambda} \mathcal{L}_n. \tag{9.4}$$

The curvatures F_A of A and F_a of a are related by

$$F_A = p^*(F_a) + \nabla_a(\phi_i) dx^i + \sum_{i<j} [\phi_i, \phi_j]\, dx^i \wedge dx^j, \tag{9.5}$$

giving for the lagrangian density of the reduced theory

$$\mathcal{L}_{n-k} = |d^{n-k}x| \left\{ -\frac{1}{2}|F_a|^2 + \frac{1}{2} \sum_{n-k}^{n-1} |\nabla_a(\phi_i)|^2 + \frac{1}{2} \sum_{i<j} |[\phi_i, \phi_j]|^2 \right\}. \tag{9.6}$$

The field content of the reduced theory is a connection and k sections of the adjoint bundle.

Most relevant to us is the reduction to $n-2$ dimensions. Thus consider a connection A invariant by translations in the x^{n-1}, x^{n-2} directions and write

$$A = a + \phi_1 dx^{n-2} + \phi_2 dx^{n-1}. \tag{9.7}$$

A short computation gives for the reduced lagrangian

$$L = |d^{n-2}x| \left\{ -\frac{1}{2}|F_a|^2 + \langle d_a\bar{\sigma}, d_a\sigma \rangle - \frac{1}{2} \| [\sigma, \bar{\sigma}] \|^2 \right\}, \tag{9.8}$$

where in the last line we set

$$\sigma = \frac{\phi_1 + \sqrt{-1}\phi_2}{\sqrt{2}}. \tag{9.9}$$

Thus σ is a section of the *complexified* adjoint bundle. Therefore, the field content of the reduction of pure Yang-Mills by two dimensions is a connection and a *complex* scalar field in the adjoint representation. There is a nonzero potential energy term for the scalar.

Consider the reduction of the bosonic part of the 4-dimensional supersymmetric Yang-Mills lagrangian (8.29) to 2 dimensions. The kinetic term for the connection reduces as in (9.8). The auxiliary field is a scalar, so the reduction is also a scalar. The topological term reduces nontrivially, and the total reduced bosonic lagrangian is

$$|d^2x|\ \frac{1}{g^2}\left\{ -\frac{1}{2}|F_a|^2 + \langle d_a\bar{\sigma}, d_a\sigma \rangle - \frac{1}{2}\| [\sigma, \bar{\sigma}] \|^2 + \frac{1}{2}|E|^2 \right\} - \sqrt{-1}\frac{\theta}{8\pi^2}\langle F_a, [\sigma, \bar{\sigma}] \rangle. \tag{9.10}$$

§9.2. Constrained connections on $M^{2|(2,2)}$

We use the notation established in (2.107)–(2.119) for $M^{2|(2,2)}$.

Let $\mathcal{P} \to M^{4|4}$ be a principal bundle with structure group G. We suppose that the vector fields $\partial_{1\dot{2}}, \partial_{2\dot{1}}$ on $M^{4|4}$ have been lifted to commuting complex conjugate vector fields $\tilde{\partial}_{1\dot{2}}, \tilde{\partial}_{2\dot{1}}$ on $\mathcal{P}$. Let $\mathcal{A}$ be a constrained connection on $\mathcal{P}$ which is invariant under these translations. All of the equations in §8.1 remain valid for $\mathcal{A}$, and we use them freely. The translation invariance is the equation

$$[\tilde{\partial}_{1\dot{2}}, \mathcal{D}_a] = [\tilde{\partial}_{2\dot{1}}, \mathcal{D}_a] = [\tilde{\partial}_{1\dot{2}}, \overline{\mathcal{D}}_{\dot{a}}] = [\tilde{\partial}_{2\dot{1}}, \overline{\mathcal{D}}_{\dot{a}}] = 0. \tag{9.11}$$

Then the vertical vector field

$$\Sigma = \frac{1}{\sqrt{2}}(\nabla_{2\dot{1}} - \tilde{\partial}_{2\dot{1}}) \tag{9.12}$$

is also translation invariant and corresponds to a section Σ of the complexified adjoint bundle. It is the basic invariant of a translationally invariant constrained connection. The Bianchi identity relates Σ to the superfields considered in §8.1. First, we consider Bianchi for $D_2, D_2, \overline{D}_{\dot{1}}$ and $D_2, \overline{D}_{\dot{1}}, \overline{D}_{\dot{1}}$ to deduce

$$\overline{\mathcal{D}}_{\dot{1}}\Sigma = \mathcal{D}_2\Sigma = 0. \tag{9.13}$$

In other words, the restriction of Σ to $M^{2|(2,2)}$ is *twisted chiral.* Then Bianchi for $D_1, \overline{D}_{\dot{1}}, \overline{D}_{\dot{2}}$ and Bianchi for $D_2, \overline{D}_{\dot{1}}, \overline{D}_{\dot{2}}$ imply

$$\begin{aligned} \sqrt{2}W_1 &= -\overline{\mathcal{D}}_{\dot{1}}\overline{\Sigma} \\ \sqrt{2}W_2 &= \ \ \overline{\mathcal{D}}_{\dot{2}}\Sigma. \end{aligned} \tag{9.14}$$

Using (9.14) and the results of §8.1 we can express the curvature $F_{\mathcal{A}}$ completely in terms of Σ.

Restrict the connection $\mathcal{A}$ and all derived superfields to $M^{2|(2,2)} \hookrightarrow M^{4|4}$. Let $i\colon \check{M}^2 \hookrightarrow M^{2|(2,2)}$ denote the inclusion, as usual. The component fields of the two-dimensional reduced theory are

$$\boxed{\begin{aligned} A &= i^*\mathcal{A} \\ \sigma &= i^*\Sigma \\ \lambda_+ = i^*W_+ &= -\frac{1}{\sqrt{2}} i^*\overline{\mathcal{D}}_+\overline{\Sigma} \\ \lambda_- = i^*W_- &= \ \ \frac{1}{\sqrt{2}} i^*\overline{\mathcal{D}}_-\Sigma \end{aligned}} \tag{9.15}$$

We also have

$$\boxed{f_A + [\sigma, \bar{\sigma}] + \sqrt{-1}E = i^*\mathcal{D}_+W_- = \frac{1}{\sqrt{2}} i^*\mathcal{D}_+\overline{\mathcal{D}}_-\Sigma,} \tag{9.16}$$

where the curvature of A is given in (8.19) as

$$F_A = 2f_A\, dy^+ \wedge dy^-. \tag{9.17}$$

The scalar field σ is a section of the complexified adjoint bundle; it is the same field which appears in the previous subsection. So too are f_A, E sections of the adjoint bundle, but both f_A and E are real.

From the point of view of $M^{2|(2,2)}$, we can think of $\mathcal{A}$ as a connection whose curvature is constrained to vanish along the odd distribution *except* for the $D_2, \overline{D}_{\dot{1}}$ direction (and its complex conjugate), where

$$[\mathcal{D}_2, \overline{\mathcal{D}}_{\dot{1}}] = -2\sqrt{2}\,\Sigma. \tag{9.18}$$

There is an important distinction between the abelian and nonabelian cases. In the abelian case the superfield Σ has values in the constant vector space $\mathfrak{g}$ and is unconstrained, except for the requirement that it be twisted chiral. Up to gauge equivalence the connection can be reconstructed from Σ. In the nonabelian case, on the other hand, the superfield Σ is a section of the adjoint bundle, and the twisted chirality condition (9.13) depends on the connection $\mathcal{A}$.

We can compute the transformation laws of the component fields under $\hat{\xi} = \eta^a \tau_{Q_a} + \bar{\eta}^{\dot{a}} \tau_{\overline{Q}_{\dot{a}}}$ directly, or by dimensional reduction from (8.22). The result is

$$\boxed{\begin{aligned}
(\hat{\xi}A)_+ &= -2\eta^-\bar{\lambda}_+ - 2\bar{\eta}^-\lambda_+ \\
(\hat{\xi}A)_- &= \;\; 2\eta^+\bar{\lambda}_- + 2\bar{\eta}^+\lambda_- \\
\hat{\xi}\sigma &= \;\; \sqrt{2}\eta^+\bar{\lambda}_+ - \sqrt{2}\bar{\eta}^-\lambda_- \\
\hat{\xi}\lambda_+ &= -\eta^-(f_A + [\sigma,\bar{\sigma}] - \sqrt{-1}E) - \sqrt{2}\,\eta^+\nabla_+\bar{\sigma} \\
\hat{\xi}\lambda_- &= -\eta^+(f_A + [\sigma,\bar{\sigma}] + \sqrt{-1}E) + \sqrt{2}\,\eta^-\nabla_-\sigma \\
\hat{\xi}f_A &= \;\; \eta^-\nabla_-\bar{\lambda}_+ + \eta^+\nabla_+\bar{\lambda}_- + \bar{\eta}^+\nabla_+\lambda_- + \bar{\eta}^-\nabla_-\lambda_+ \\
\hat{\xi}E &= -\sqrt{-1}(-\eta^+\nabla_+\bar{\lambda}_- + \eta^-\nabla_-\bar{\lambda}_+ + \bar{\eta}^+\nabla_+\lambda_- - \bar{\eta}^-\nabla_-\lambda_+).
\end{aligned}} \tag{9.19}$$

From (9.12) it is easy to see that the R-symmetries J_+, J_- (see (2.117)) act on Σ by

$$\boxed{[J_\pm, \Sigma] = \mp\Sigma.} \tag{9.20}$$

From this and (9.15) we deduce the action on the component fields:

$$\boxed{\begin{aligned}
[J_\pm, A] = [J_\pm, f_A] = [J_\pm, E] = [J_+, \lambda_+] = [J_-, \lambda_-] = 0 \\
[J_\pm, \sigma] = \mp\sigma \\
[J_\pm, \bar{\sigma}] = \pm\bar{\sigma} \\
[J_+, \lambda_-] = -\lambda_- \\
[J_-, \lambda_+] = -\lambda_+.
\end{aligned}} \tag{9.21}$$

§9.3. The reduced Yang-Mills action

We rewrite the action (8.25) in terms of Σ. First, using (9.14) and (9.13) we have

$$\begin{aligned}
\epsilon^{ab}\langle W_a, W_b\rangle &= 2\langle W_+, W_-\rangle \\
&= -\langle \overline{\mathcal{D}}_+\overline{\Sigma}, \overline{\mathcal{D}}_-\Sigma\rangle \\
&= -\overline{D}_+\overline{D}_-\langle\overline{\Sigma}, \Sigma\rangle \\
&= -\overline{D}^2\|\Sigma\|^2.
\end{aligned} \tag{9.22}$$

Let $\check{\mathcal{L}}'_2$ denote the dimensionally reduced component lagrangian function. From the second line of (8.25) we deduce

(9.23)
$$\begin{aligned}\check{\mathcal{L}}'_2 &= \frac{1}{2\sqrt{-1}}\Big\{\int d^2\theta\,\frac{-\tau}{16\pi}\epsilon^{ab}\langle W_a, W_b\rangle + \int d^2\bar\theta\,\frac{\bar\tau}{16\pi}\epsilon^{\dot a\dot b}\langle \overline{W}_{\dot a}, \overline{W}_{\dot b}\rangle\Big\}\\ &= \frac{1}{2\sqrt{-1}}i^*\Big\{-\frac{\tau}{16\pi}D^2\overline{D}^2\|\Sigma\|^2 + \frac{\bar\tau}{16\pi}\overline{D}^2D^2\|\Sigma\|^2\Big\}\\ &= -\frac{\operatorname{Im}\tau}{4\pi}i^*\frac{1}{2}(D^2\overline{D}^2 + \overline{D}^2D^2)\|\Sigma\|^2 - \frac{\operatorname{Re}\tau}{16\pi\sqrt{-1}}i^*\frac{1}{2}(D^2\overline{D}^2 - \overline{D}^2D^2)\|\Sigma\|^2.\end{aligned}$$

Recall from §2.6 that for terms in a lagrangian in $M^{2|(2,2)}$ involving twisted chiral fields only we compute component lagrangians using exactly the combination of derivatives which occurs in the first term of (9.23). The second term is the topological term in (9.10); it does not seem to reduce to an integral $\int d^4\theta$. So for $\operatorname{Re}\tau = 0$ the $N = 2$ superspace Yang-Mills lagrangian in $M^{2|(2,2)}$ is

(9.24)
$$\boxed{\mathcal{L}'_2 = |d^2y|\,d^4\theta\,\frac{-1}{4g^2}\|\Sigma\|^2.}$$

The corresponding component lagrangian is the dimensional reduction of (8.29), which we compute either directly or with the aid of (9.8). After eliminating the auxiliary field E we obtain

(9.25)
$$\boxed{\begin{aligned}\check{\mathcal{L}}'_2 \sim |d^2y|\,\frac{1}{g^2}\Big\{&-\frac{1}{2}|F_A|^2 + \langle d_A\bar\sigma, d_A\sigma\rangle + \langle\bar\lambda\,\not{D}_A\,\lambda\rangle\\ &-\frac{1}{2}\|\,[\sigma,\bar\sigma]\,\|^2 + \sqrt{2}\,\big(\langle\bar\lambda^-, [\bar\sigma, \lambda^+]\rangle + \langle\bar\lambda^+, [\sigma, \lambda^-]\rangle\big)\Big\}.\end{aligned}}$$

The (classical) theory is invariant under the R-symmetries J_+, J_-, which are defined in (2.117). This follows either in superspace from (9.20) and (9.24) or in components from (9.21) and (9.25).

Another type of topological term may be defined for an invariant trace

(9.26)
$$\langle\cdot\rangle\colon \mathfrak{g} \longrightarrow \mathbb{R},$$

and it has a superspace expression. Define

(9.27)
$$\int d\bar\theta^-\,d\theta^+\,\ell = i^*D_+\overline{D}_-\ell.$$

The topological term is then

(9.28)
$$\boxed{\mathcal{L}'_3 = |d^2x|\,\frac{1}{\sqrt{2}}\operatorname{Re}\Big(d\bar\theta^-\,d\theta^+\,\langle\Sigma\rangle\Big).}$$

The component expression is easily seen from (9.16) to be

(9.29)
$$|d^2x|\,\check{\mathcal{L}}'_3 = |d^2x|\,\Big\{\langle f_A\rangle + \sqrt{-1}\langle E\rangle\Big\}.$$

CHAPTER 10
$N = 1$ Yang-Mills in Dimension 6 and $N = 2$ Yang-Mills in Dimension 4

In §10.1 we complete the discussion of the superspace formulation of super Yang-Mills theory in 6 dimensions. As in Chapter 7 and Chapter 8, the main point is to work out the structure of the auxiliary fields and the R-symmetries. The proof that the category of component fields (including the auxiliary fields) is equivalent to the category of superfields is deferred to the next chapter. We did not succeed in finding a manifestly supersymmetric formula for the lagrangian in this case. Section 10.2 is parallel to Chapter 9: We reduce the 6-dimensional pure gauge theory to 4 dimensions, where it has twice the minimal amount of supersymmetry. Neither theories in 4 dimensions with 8 supersymmetries which include matter fields nor theories in 4 dimensions with 16 supersymmetries have superspace formulations which exhibit all of the supersymmetry manifestly. But they can all be written on $M^{4|4}$, where 4 of the supersymmetries are manifest. We do this in §10.3.

§10.1. Constrained connections on $M^{6|(8,0)}$

We specialize the discussion in §6.3 to $M^{6|(8,0)}$. Our first task is to evaluate (6.74). To that end we compute using (2.89)–(2.91):

$$\begin{aligned} g\big(\tilde{\Gamma}(f^{ai}, f^{ck}), \Gamma(f_{bj}, f_{d\ell})\big) &= \frac{1}{2}\epsilon^{ik}\nu^{pqac}\epsilon_{j\ell}\, g(e_{pq}, e_{bd}) \\ &= \epsilon^{ik}\epsilon_{j\ell}\nu^{pqac}\nu_{pqbd} \\ &= 2(\delta^a_b\delta^c_d - \delta^a_d\delta^c_b)\epsilon^{ik}\epsilon_{j\ell}. \end{aligned} \tag{10.1}$$

The curvature term in (6.74) may be expanded using

$$\begin{aligned} \tilde{\Gamma}(f^{ai}, f^{ck}) \otimes \Gamma(f_{ck}, f_{bj}) &= \frac{1}{2}\epsilon^{ik}\nu^{pqac}\epsilon_{kj}\; e_{pq} \otimes e_{cb} \\ &= -\frac{1}{2}\delta^i_j\, \nu^{pqac}\; e_{pq} \otimes e_{cb}, \end{aligned} \tag{10.2}$$

so that the left hand side of (6.74) is

$$-2(-\frac{1}{2}\delta^i_j\nu^{pqac})\mathcal{F}_{(pq)(cb)} = \delta^i_j\,\nu^{pqac}\mathcal{F}_{(pq)(cb)} = \delta^i_j\,\nu^{acpq}\mathcal{F}_{(bc)(pq)}. \tag{10.3}$$

For $a = b$ this vanishes, since ν^{acpq} is symmetric in $(ac), (pq)$ whereas $\mathcal{F}_{(ac)(pq)}$ is skew. Now evaluate (6.74) for $i = j = 1$, then $i = j = 2$, and sum over $a = b$ to learn

$$\mathcal{D}_{ai}\Lambda^{ai} = 0. \tag{10.4}$$

Define

$$E^i_j = -\frac{\sqrt{-1}}{4}\mathcal{D}_{aj}\Lambda^{ai}. \tag{10.5}$$

Then with a bit of work we derive

$$\mathcal{D}_{bj}\Lambda^{ai} = -\frac{1}{2}\tilde{\Gamma}^{\mu(ai)(ck)}\Gamma^{\nu}_{(ck)(bj)}\mathcal{F}_{\mu\nu} + \delta^a_b\sqrt{-1}\,E^i_j, \tag{10.6}$$

which has the desired form (6.76).

To proceed further we remark on some linear algebra. For S_0 of dimension 4, the natural map

$$\textstyle\bigwedge^2(\bigwedge^2 S_0) \to \bigwedge^2 S_0 \otimes \bigwedge^2 S_0 \to \bigwedge^3 S_0 \otimes S_0, \tag{10.7}$$

which is $u\wedge v \mapsto v\otimes v - v\otimes u$ followed by $(x\wedge y)\otimes(z\wedge t) \mapsto (x\wedge y\wedge z)\otimes t - (x\wedge y\wedge t)\otimes z$, gives rise to a short exact sequence

$$0 \to \textstyle\bigwedge^2(\bigwedge^2 S_0) \to \bigwedge^3 S_0 \otimes S_0 \to \bigwedge^4 S_0 \to 0. \tag{10.8}$$

For S_0 given with a volume form, $\bigwedge^3 S_0$ is the dual of S_0, the map to $\bigwedge^4 S_0$ becomes the evaluation map, and we get

$$\textstyle\bigwedge^2(\bigwedge^2 S_0) \cong \{\text{trace zero endomorphisms of } S_0\}. \tag{10.9}$$

This makes it convenient to define

$$f^a_b := \frac{1}{4}\nu^{acde}\mathcal{F}_{(bc)(de)}. \tag{10.10}$$

By (10.9) the tensor f is traceless:

$$f^a_a = 0. \tag{10.11}$$

Also,

$$\mathcal{F}_{(ab)(cd)} = \nu_{ebcd}f^e_a + \nu_{aecd}f^e_b. \tag{10.12}$$

Then using (10.3) we see that (10.6) simplifies to

$$\boxed{\mathcal{D}_{bj}\Lambda^{ai} = \delta^i_j\, f^a_b + \delta^a_b\sqrt{-1}\,E^i_j}. \tag{10.13}$$

We need one additional formula

$$\mathcal{D}_{ai}E^j_k = \sqrt{-1}\,(\epsilon_{ik}\nabla_{ab}\Lambda^{bj} - \delta^j_i\epsilon_{k\ell}\nabla_{ab}\Lambda^{b\ell}), \tag{10.14}$$

which is equivalent to (11.30) and will be explained in Chapter 11.

Now we define the component fields corresponding to a constrained connection. First, we have a connection A on the restriction $P \to \check{M}^6$ of $\mathcal{P}$ to Minkowski space. The odd part of the curvature restricts to a dual spinor field

$$\boxed{i^*\Lambda^{ai} = \lambda^{ai}.} \tag{10.15}$$

The restriction of E^i_j to $\check{M}^6$, which we continue to denote 'E^i_j', is an auxiliary field. Using (2.95), (2.97), and (6.66) we learn that

$$\overline{E^i_j} = E^j_i, \tag{10.16}$$

By (10.4) we see that E is traceless:

$$E^i_i = 0. \tag{10.17}$$

Thus $\left(E^i_j\right)$ has three independent real components; the matrix $\left(\sqrt{-1}\,E^i_j\right)$ is traceless skew-Hermitian (with values in the adjoint bundle). It transforms in the adjoint representation of the $Sp(1)_R$ symmetry (2.102). In other words, the auxiliary fields comprise an imaginary quaternion. The six-dimensional *vector multiplet* is the collection of component fields $\{A, \lambda^{ai}, E^i_j\}$.

Proposition 10.18. *The category of connections $\mathcal{A}$ on $M^{6|(8,0)}$ whose curvature vanishes on the odd distribution τ is equivalent to the category of triples $\{A, \lambda, E\}$ consisting of a connection A on $\check{M}^6$, a dual spinor field λ with values in the adjoint bundle, and an imaginary quaternion scalar field E with values in the adjoint bundle.*

The proof of Proposition 10.18 is presented in Chapter 11.

We derive the supersymmetry transformation laws for the components as in previous sections. Denote the restriction of f^a_b to $\check{M}^6$ by 'f^a_b'. Then the supersymmetry transformation $\eta^{ai}Q_{ai}$ induces the following transformation on component fields:

$$\boxed{\begin{aligned} (\hat{\xi}A)_{pq} &= 2\eta^{ai}\nu_{pqab}\epsilon_{ij}\lambda^{bj} \\ \hat{\xi}\lambda^{ai} &= -\eta^{bi}f^a_b - \eta^{aj}\sqrt{-1}\,E^i_j \\ \hat{\xi}E^i_j &= \eta^{ai}\sqrt{-1}\,(\not{D}\lambda)_{aj} - \eta^{ak}\sqrt{-1}\,\epsilon_{jk}\epsilon^{i\ell}(\not{D}\lambda)_{a\ell}. \end{aligned}} \tag{10.19}$$

and

$$\boxed{\hat{\xi}f^a_b = 2\eta^{ai}(\not{D}\lambda)_{bi} - 2\eta^{ci}\epsilon_{ij}\nabla_{bc}\lambda^{aj} - \delta^a_b\eta^{ci}(\not{D}\lambda)_{ci}.} \tag{10.20}$$

Recall from (2.102) that the R-symmetry group is $SU(2)_R$. The matrix $S = (S^i_j)$ acts on the component fields by

$$\boxed{\begin{aligned} (SA) &= A \\ (S\lambda)^{ai} &= (S^{-1})^i_{i'}\lambda^{ai'} \\ (SE)^i_j &= (S^{-1})^i_{i'}E^{i'}_{j'}S^{j'}_j\,. \end{aligned}} \tag{10.21}$$

§10.2. Reduction to $M^{4|8}$

The story here is analogous to that in Chapter 9, where we reduce Yang-Mills on $M^{4|4}$ to $M^{2|(2,2)}$. For a worked out example, see Problem ASH8 of [I-**Homework**].

We consider a constrained connection $\mathcal{A}$ on a G-bundle $\mathcal{P} \to M^{6|(8,0)}$ which is "constant" in the $\partial_{12}, \partial_{34}$ directions. That is, ∂_{12} and ∂_{34} are lifted to commuting complex conjugate vector fields $\tilde{\partial}_{12}$ and $\tilde{\partial}_{34}$ on $\mathcal{P}$, and the horizontal vector fields $\mathcal{D}_{ai}$ on $\mathcal{P}$ satisfy

$$[\tilde{\partial}_{12}, \mathcal{D}_{ai}] = [\tilde{\partial}_{34}, \mathcal{D}_{ai}] = 0. \tag{10.22}$$

Set

$$\Sigma = \frac{1}{\sqrt{2}}(\nabla_{34} - \tilde{\partial}_{34}). \tag{10.23}$$

Then applying Bianchi to D_{ai}, D_{31}, D_{42} we learn

$$\mathcal{D}_{3i}\Sigma = \mathcal{D}_{4i}\Sigma = 0. \tag{10.24}$$

From Bianchi we also find, after some tedious computation, a formula for Λ^{ai} in terms of derivatives of Σ:

$$\sqrt{2}\,\Lambda^{ai} = \begin{cases} J^a_b \epsilon^{ij} \mathcal{D}_{bj}\Sigma, & a = 1, 2; \\ J^a_b \epsilon^{ij} \mathcal{D}_{bj}\overline{\Sigma}, & a = 3, 4. \end{cases} \tag{10.25}$$

Here J^a_b is the matrix of the pseudoreal structure j described before (2.95). Also, from (10.6) we learn that $\mathcal{D}_{bj}\Lambda^{ai}$ is independent of a, b if $i \neq j$. It follows that

$$\begin{aligned} \mathcal{D}_{11}\mathcal{D}_{21}\Sigma &= \mathcal{D}_{31}\mathcal{D}_{41}\overline{\Sigma} \\ \mathcal{D}_{12}\mathcal{D}_{22}\Sigma &= \mathcal{D}_{32}\mathcal{D}_{42}\overline{\Sigma}. \end{aligned} \tag{10.26}$$

Let $j\colon M^{4|8} \hookrightarrow M^{6|(8,0)}$ be the inclusion, as described in the last paragraph of §2.5. We follow the notation introduced there and also use formula (2.104). Now write $\tilde{\mathcal{A}}, \tilde{D}_{ai}, \tilde{\Sigma}, \tilde{\Lambda}^{ai}, \tilde{E}^i_j$ for the fields on $M^{6|(8,0)}$ to distinguish them from their restrictions to $M^{4|8}$. Set

$$\Sigma = j^*\tilde{\Sigma}. \tag{10.27}$$

Then using (2.104), equation (10.24) asserts that Σ is a *chiral* superfield on $M^{4|8}$ in the sense that

$$\overline{\mathcal{D}}_{\dot{a}(i)}\Sigma = 0, \qquad a = 1, 2, \quad i = 1, 2. \tag{10.28}$$

Formula (10.25) expresses $\Lambda^{ai} = j^*\tilde{\Lambda}^{ai}$ in terms of Σ. We record the result in full:

$$\begin{aligned} \sqrt{2}\,\Lambda^{11} &= -\mathcal{D}^{(2)}_2\Sigma & \sqrt{2}\,\Lambda^{12} &= \mathcal{D}^{(1)}_2\Sigma \\ \sqrt{2}\,\Lambda^{21} &= \mathcal{D}^{(2)}_1\Sigma & \sqrt{2}\,\Lambda^{22} &= -\mathcal{D}^{(1)}_1\Sigma \\ \sqrt{2}\,\Lambda^{31} &= -\overline{\mathcal{D}}_{\dot{2}(1)}\overline{\Sigma} & \sqrt{2}\,\Lambda^{32} &= -\overline{\mathcal{D}}_{\dot{2}(2)}\overline{\Sigma} \\ \sqrt{2}\,\Lambda^{41} &= \overline{\mathcal{D}}_{\dot{1}(1)}\overline{\Sigma} & \sqrt{2}\,\Lambda^{42} &= \overline{\mathcal{D}}_{\dot{1}(2)}\overline{\Sigma} \end{aligned} \tag{10.29}$$

We also work out the auxiliary fields $E^i_j = j^*\tilde{E}^i_j$ using the definition (10.5) together with (10.16), (10.29), and (10.26):

$$\begin{aligned}
E^1_1 &= -\frac{\sqrt{-1}}{2\sqrt{2}}\big(\mathcal{D}^{(1)}_2\mathcal{D}^{(2)}_1\Sigma - \mathcal{D}^{(1)}_1\mathcal{D}^{(2)}_2\Sigma\big)\\
E^1_2 &= -\frac{\sqrt{-1}}{\sqrt{2}}\mathcal{D}^{(2)}_2\mathcal{D}^{(2)}_1\Sigma\\
&= -\frac{\sqrt{-1}}{\sqrt{2}}\overline{\mathcal{D}}_{\dot{2}(1)}\overline{\mathcal{D}}_{\dot{1}(1)}\overline{\Sigma}\\
E^2_1 &= -\frac{\sqrt{-1}}{\sqrt{2}}\mathcal{D}^{(1)}_1\mathcal{D}^{(1)}_2\Sigma\\
&= -\frac{\sqrt{-1}}{\sqrt{2}}\overline{\mathcal{D}}_{\dot{1}(2)}\overline{\mathcal{D}}_{\dot{2}(2)}\overline{\Sigma}\\
E^2_2 &= -E^1_1.
\end{aligned} \tag{10.30}$$

From the point of view of $M^{4|8}$, we can think of $\mathcal{A}$ as a connection whose curvature is constrained to vanish along the odd distribution *except* for the nontrivial curvature

$$[\overline{\mathcal{D}}_{\dot{1}(2)}, \overline{\mathcal{D}}_{\dot{2}(1)}] = -[\overline{\mathcal{D}}_{\dot{2}(2)}, \overline{\mathcal{D}}_{\dot{1}(1)}] = 2\sqrt{2}\,\Sigma \tag{10.31}$$

and the complex conjugate.

As in Chapter 9 there is a sharp distinction between the abelian and nonabelian cases—in the abelian case Σ takes values in the constant vector space $\mathfrak{g}$, the only constraint being (10.28); whereas in the nonabelian case Σ is a section of $\operatorname{ad}\mathcal{P}$ and the constraint (10.28) involves the connection $\mathcal{A}$, which therefore cannot be derived solely from Σ.

Formulas (10.29) and (10.30) are more understandable if we restrict to $M^{4|4} \hookrightarrow M^{4|8}$. (The inclusion $M^{4|4} \hookrightarrow M^{6|(8,0)}$ is discussed towards the end of §2.5.) Let $\hat{\mathcal{A}}, \hat{\Sigma}$ denote the restrictions of $\mathcal{A}, \Sigma$ to $M^{4|4}$. Then $\hat{\mathcal{A}}$ is a vector superfield (cf. §8.1)—a constrained connection on the restriction $\hat{\mathcal{P}}$ of $\mathcal{P}$ to $M^{4|4}$—and (10.28) implies that $\hat{\Sigma}$ is a chiral superfield (cf. §5.1) with values in $\operatorname{ad}\hat{\mathcal{P}}$. The pair $(\hat{\mathcal{A}}, \hat{\Sigma})$ is called an $N = 2$ *vector multiplet* on $M^{4|4}$. We claim, but do not write out a proof, that the category of constrained connections $\mathcal{A}$ on $M^{6|(8,0)}$ which satisfy (10.22) is equivalent to the category of pairs $(\hat{\mathcal{A}}, \hat{\Sigma})$ on $M^{4|4}$.

Let $i\colon \check{M}^4 \hookrightarrow M^{4|8}$ be the inclusion. Motivated by formulas (5.7), and (8.20), (8.4) we define the component fields of the superfields on $M^{4|8}$ to be

$$\boxed{\begin{aligned}
A &= i^*\mathcal{A}\\
\sigma &= i^*\Sigma\\
\lambda_a = i^*\epsilon_{ba}\Lambda^{b1} \qquad &= -\frac{1}{\sqrt{2}}\,i^*\mathcal{D}^{(2)}_a\Sigma\\
\chi_a = i^*\epsilon_{ba}\Lambda^{b2} \qquad &= \frac{1}{\sqrt{2}}\,i^*\mathcal{D}^{(1)}_a\Sigma\\
E = i^*E^1_1 \qquad &= -\frac{\sqrt{-1}}{2\sqrt{2}}\,i^*\big(\mathcal{D}^{(1)}_2\mathcal{D}^{(2)}_1\Sigma - \mathcal{D}^{(1)}_1\mathcal{D}^{(2)}_2\Sigma\big)\\
F = -\frac{\sqrt{-1}}{\sqrt{2}}\,i^*E^2_1 &= \frac{1}{2}\,i^*\mathcal{D}^{(1)}_2\mathcal{D}^{(1)}_1\Sigma.
\end{aligned}} \tag{10.32}$$

The indices a, b in these formulas run over $\{1, 2\}$. Also, we use (10.29), (10.30). Following (8.14) we define the dual spinor field

$$\lambda^a = i^*\Lambda^{a1} = -\frac{1}{\sqrt{2}}\, i^*\epsilon^{ab}\mathcal{D}_b^{(2)}\Sigma. \tag{10.33}$$

These component fields group together into $N = 1$ supermultiplets: (A, λ, E) is an $N = 1$ vector multiplet (§8.1) and (σ, χ, F) is an $N = 1$ chiral multiplet (§5.1). The collection $(A, \sigma, \lambda, \chi, E, F)$ of all component fields is called an $N = 2$ *vector multiplet*. The reader should check that (10.32) is consistent with (5.7) and (8.20). The only nontrivial compatibility check is to see that $\tilde{\Lambda}^{a1}$ $(a = 1, 2)$ on $M^{6|(8,0)}$ restricts on $M^{4|4}$ to be the superfield $\hat{\Lambda}^a$, with both $\tilde{\Lambda}$ and $\hat{\Lambda}$ defined by (6.66); we leave this check to the reader. Note that on $M^{4|4}$ we have

$$\hat{W}_a = -\frac{1}{\sqrt{2}}\, i^*\mathcal{D}_a^{(2)}\hat{\Sigma}. \tag{10.34}$$

On $M^{4|4}$ only half of the supersymmetry of the vector multiplet $(\hat{\mathcal{A}}, \hat{\Sigma})$ is manifest. For the rest we write the vector field $\hat{\xi}'$ on the space of $N = 1$ superfields $\mathcal{F}_{(\hat{\mathcal{A}},\hat{\Sigma})}$ which corresponds to the supersymmetry transformation

$$\eta'^a Q_a^{(2)} + \bar{\eta}'^{\dot{a}}\overline{Q}_{\dot{a}(2)}. \tag{10.35}$$

From the formulas in this section we find

$$\boxed{\begin{aligned} (\hat{\xi}'\hat{\mathcal{A}})_a &= 2\sqrt{2}\,\eta'^b\epsilon_{ba}\overline{\hat{\Sigma}} \\ (\hat{\xi}'\hat{\mathcal{A}})_{a\dot{b}} &= \sqrt{2}\big(\eta'^c\epsilon_{ca}\overline{\mathcal{D}}_{\dot{b}}\overline{\hat{\Sigma}} + \bar{\eta}'^{\dot{c}}\epsilon_{\dot{c}\dot{b}}\mathcal{D}_a\hat{\Sigma}\big) \\ \hat{\xi}'\hat{\Sigma} &= \sqrt{2}\,\eta'^a\hat{W}_a \end{aligned}} \tag{10.36}$$

The penultimate line of (6.79) is used to derive the first two of these formulas.

Next, we write all of the supersymmetry transformations on the component fields on $\check{M}^4$. Let $\hat{\xi}$ be the vector field on the space $\mathcal{F}_{(A,\sigma,\lambda,\chi,E,F)}$ of component fields induced by the supersymmetry transformation

$$\eta^a Q_a^{(1)} + \bar{\eta}^{\dot{a}}\overline{Q}_{\dot{a}(1)} + \eta'^a Q_a^{(2)} + \bar{\eta}'^{\dot{a}}\overline{Q}_{\dot{a}(2)}. \tag{10.37}$$

Then from (10.19) (or (8.22) together with (10.36) and the other formulas of this section) we deduce

(10.38)

$$\boxed{\begin{aligned} (\hat{\xi}A)_{a\dot{b}} &= \ \ 2\eta^c\epsilon_{ca}\bar{\lambda}_{\dot{b}} + 2\bar{\eta}^{\dot{c}}\epsilon_{\dot{c}\dot{b}}\lambda_a + 2\eta'^c\epsilon_{ca}\bar{\chi}_{\dot{b}} + 2\bar{\eta}'^{\dot{c}}\epsilon_{\dot{c}\dot{b}}\chi_a \\ \hat{\xi}\sigma &= -\sqrt{2}\,\eta^a\chi_a + \sqrt{2}\,\eta'^a\lambda_a \\ \hat{\xi}\lambda_a &= -\eta^b F_{ab} + \eta^b\epsilon_{ab}\sqrt{-1}\,E + \sqrt{2}\,\eta'^b\epsilon_{ab}\overline{F} - \sqrt{2}\,\bar{\eta}'^{\dot{b}}\nabla_{a\dot{b}}\sigma \\ \hat{\xi}\chi_a &= -\sqrt{2}\,\eta^b\epsilon_{ab}F + \sqrt{2}\,\bar{\eta}^{\dot{b}}\nabla_{a\dot{b}}\sigma - \eta'^b F_{ab} - \eta'^b\epsilon_{ab}\sqrt{-1}\,E \\ \hat{\xi}E &= \eta^a\sqrt{-1}\,(\not{D}\bar{\lambda})_a - \bar{\eta}^{\dot{a}}\sqrt{-1}\,(\not{D}\lambda)_{\dot{a}} - \eta'^a\sqrt{-1}\,(\not{D}\bar{\chi})_a + \bar{\eta}'^{\dot{a}}\sqrt{-1}\,(\not{D}\lambda)_{\dot{a}} \\ \hat{\xi}F &= \sqrt{2}\,\eta'^a(\not{D}\bar{\lambda})_a - \sqrt{2}\,\bar{\eta}^{\dot{a}}(\not{D}\chi)_{\dot{a}} \end{aligned}}$$

The R-symmetry group of the dimensionally reduced theory is $U(2)$ (see the end of §2.5). The matrix $S = (S^i_j)$ acts on the component fields by

(10.39)

$$\boxed{\begin{aligned} S\colon A &\longmapsto A \\ S\colon \sigma &\longmapsto (\det S)^{-1}\sigma \\ S\colon \begin{pmatrix}\lambda_a\\ \chi_a\end{pmatrix} &\longmapsto S^{-1}\begin{pmatrix}\lambda_a\\ \chi_a\end{pmatrix} \\ S\colon \begin{pmatrix} E & -\sqrt{-1}\sqrt{2}\,\overline{F}\\ \sqrt{-1}\sqrt{2}\,F & -E\end{pmatrix} &\longmapsto S^{-1}\begin{pmatrix} E & -\sqrt{-1}\sqrt{2}\,\overline{F}\\ \sqrt{-1}\sqrt{2}\,F & -E\end{pmatrix} S. \end{aligned}}$$

We remark that it is useful to check (10.38) using equations (5.5), (8.22), and the R-symmetry (10.39) for $S = \left(\begin{smallmatrix}0&1\\1&0\end{smallmatrix}\right)$.

Finally, we write the $N = 2$ Yang-Mills lagrangian in four dimensions in a manifestly supersymmetric way on $M^{4|8}$. As with $M^{4|4}$ (see the beginning of §5.3) there is a "chiral odd integration" on $M^{4|8}$. Namely, if Υ is any superfield on $M^{4|8}$ which is chiral in the sense that $\overline{D}_{\dot a(i)}\Upsilon = 0$, then

$$|d^4x|\int d^4\theta\,\Upsilon = |d^4x|\; i^*D_2^{(1)}D_1^{(1)}D_2^{(2)}D_1^{(2)}\Upsilon \tag{10.40}$$

is a well-defined density on $\check M^4$. Given a connection $\mathcal{A}$ on $M^{4|8}$ of the type we have been considering, with Σ defined as above, set

$$\boxed{\mathcal{L} = |d^4x|\;\mathrm{Im}\Big\{d^4\theta\,\frac{\tau}{32\pi}\langle\Sigma,\Sigma\rangle\Big\}.} \tag{10.41}$$

We claim that $\mathcal{L}$ is a superspace lagrangian which reproduces in components the dimensional reduction of Yang-Mills that we seek. To verify this we transform it into a lagrangian on $N = 1$ superspace $M^{4|4}$. Namely, the component lagrangian function $\check{\mathcal{L}}$ may be written as

$$\begin{aligned}
\check{\mathcal{L}} &= \mathrm{Im}\Big\{\frac{\tau}{32\pi}i^*D_2^{(1)}D_1^{(1)}D_2^{(2)}D_1^{(2)}\langle\Sigma,\Sigma\rangle\Big\} \\
&= \mathrm{Im}\Big\{\frac{\tau}{16\pi}i^*D_2^{(1)}D_1^{(1)}\big(-\langle\mathcal{D}_1^{(2)}\Sigma,\mathcal{D}_2^{(2)}\Sigma\rangle + \langle\mathcal{D}_2^{(2)}\mathcal{D}_1^{(2)}\Sigma,\Sigma\rangle\big)\Big\} \\
&= \mathrm{Im}\Big\{\frac{\tau}{16\pi}i^*D_2^{(1)}D_1^{(1)}\big(-2\langle W_1,W_2\rangle + \langle\overline{D}_{\dot 2(1)}\overline{D}_{\dot 1(1)}\overline{\Sigma},\Sigma\rangle\big)\Big\} \\
&= i^*D_2^{(1)}D_1^{(1)}\,\mathrm{Im}\big\{-\frac{\tau\epsilon^{ab}}{16\pi}\langle W_a,W_b\rangle\big\} + \frac{\mathrm{Im}\,\tau}{16\pi}i^*D_2^{(1)}D_1^{(1)}\overline{D}_{\dot 2(1)}\overline{D}_{\dot 1(1)}\langle\overline{\Sigma},\Sigma\rangle \\
&= \int d^2\theta\;\mathrm{Im}\big\{-\frac{\tau\epsilon^{ab}}{16\pi}\langle W_a,W_b\rangle\big\} + \frac{\mathrm{Im}\,\tau}{4\pi}\int d^4\theta\frac{1}{4}\langle\overline{\Sigma},\Sigma\rangle.
\end{aligned} \tag{10.42}$$

In the third equation we use (10.34) and (10.26) (cf. (10.30)); in the fourth we use the fact that Σ is chiral; and in the fifth we use notions of integration on $M^{4|4}$. For the $d^4\theta$ term we use (2.71), so that the last equality holds only up to exact terms, which we ignore. The component expansion of the first term is (8.29). The component expansion of the second is (5.13), but we must remember the extra

terms explained in §8.3. Recall that for this we eliminate the auxiliary field E using (8.50):

$$\boxed{E = \sqrt{-1}\,[\sigma, \bar{\sigma}].} \tag{10.43}$$

Set

$$\tau = \frac{\theta}{2\pi} + \frac{4\pi\sqrt{-1}}{g^2}. \tag{10.44}$$

The component lagrangian is then a special case of (6.42):

(10.45)

$$\boxed{\begin{aligned} d^4x\,\check{\mathcal{L}} = d^4x\,\frac{1}{g^2}\Big\{ &-\frac{1}{2}|F_A|^2 + \langle d_A\bar{\sigma}, d_A\sigma\rangle + \langle\bar{\lambda}\not{D}_A\lambda\rangle + \langle\bar{\chi}\not{D}_A\chi\rangle - \frac{1}{2}|\,[\sigma,\bar{\sigma}]\,|^2 \\ &-\sqrt{2}\,\big(\langle\bar{\sigma},[\lambda^a,\chi_a]\rangle + \langle[\bar{\lambda}^{\dot{a}},\bar{\chi}_{\dot{a}}],\sigma\rangle\big) + \langle\overline{F},F\rangle\Big\} + \frac{\theta}{16\pi^2}\langle F_A\wedge F_A\rangle. \end{aligned}}$$

The $N = 2$ superspace expression (10.41) is convenient for studying quantum corrections to the classical theory (which are explained in Lecture 18 of [II-**Dynamics of QFT**]). Define the *holomorphic prepotential*

$$\mathcal{F}(\Sigma) = \frac{\tau}{2}\langle\Sigma,\Sigma\rangle. \tag{10.46}$$

Then we rewrite (10.41) as

$$\boxed{\mathcal{L} = |d^4x|\ \mathrm{Im}\Big\{d^4\theta\,\frac{\tau}{16\pi}\mathcal{F}(\Sigma)\Big\}.} \tag{10.47}$$

Using (5.17) we see that the last term of (10.42) corresponds to the Kähler potential

$$K(\overline{\Sigma},\Sigma) = \frac{1}{2}\,\mathrm{Im}\big(\overline{\Sigma}\,\frac{\partial\mathcal{F}}{\partial\Sigma}\big). \tag{10.48}$$

We can replace the classical prepotential $\mathcal{F}$ by any holomorphic function of Σ and still obtain a supersymmetric theory; the relation (10.48) between the prepotential $\mathcal{F}$ and the Kähler potential K is unchanged. The computation of the quantum effective action is then the determination of the function $\mathcal{F}$. Kähler metrics of the form (10.48) are called *special* by physicists, and indeed they exhibit many special properties.[41]

§10.3. More theories on $M^{4|4}$ with extended supersymmetry

We record without proof the form of the general $N = 2$ model in 4 dimensions written on the $N = 1$ superspace $M^{4|4}$. Recall (6.27) that the scalars take values in a hyperkähler manifold X. As in §3.6 we fix a Kähler structure on X. Then

[41] For a mathematical treatment of special Kähler geometry, see: D. S. Freed, *Special Kähler manifolds*, Commun. Math. Phys., to appear, `hep-th/9712042`.

the rest of the hyperkähler structure is encoded in a holomorphic symplectic form. The moment map $\mu_{\mathbb{H}}$ of the G action on X is

$$\mu_{\mathbb{H}} = (\mu_{\mathbb{R}}, \mu_{\text{hol}}), \tag{10.49}$$

where $\mu_{\mathbb{R}}\colon X \to \mathfrak{g}^*$ is the moment map for the Kähler structure and $\mu_{\text{hol}}\colon X \to \mathfrak{g}^*_{\mathbb{C}}$ the complex moment map of the holomorphic symplectic structure. The fields in the model are a constrained connection $\mathcal{A}$ on some principal bundle $\mathcal{P} \to M^{4|4}$ and a chiral superfield Φ with values in the associated fiber bundle $\tilde{X}^{\mathcal{P}}$, where

$$\tilde{X} = X \times \mathfrak{g}_{\mathbb{C}}. \tag{10.50}$$

Note that the $\mathfrak{g}_{\mathbb{C}}$-valued scalars are part of the $N = 2$ vector multiplet, as explained in §10.2. The total Kähler moment map is

$$\mu(\phi, \sigma) = \mu_{\mathbb{R}}(\phi) + \frac{\sqrt{-1}}{2}[\sigma, \bar{\sigma}]. \tag{10.51}$$

There is a nonzero superpotential

$$W = \sqrt{2}\,\tilde{\mu}_{\text{hol}} \tag{10.52}$$

coming from the transpose of the holomorphic moment map:

$$\tilde{\mu}_{\text{hol}}\colon X \times \mathfrak{g}_{\mathbb{C}} \longrightarrow \mathbb{C}. \tag{10.53}$$

We leave the reader to verify that the component lagrangian (6.42) for these data is the dimensional reduction of (6.48). As mentioned at the end of §3.4, if ζ is a vector field on X which preserves the hyperkähler structure, then we can use ζ to add a term to the superpotential. Namely, if η is the holomorphic symplectic form on X for the particular choice of complex structure we have made, we add to W the holomorphic function W_0 which satisfies

$$\partial W_0 = -\iota(\zeta)\eta. \tag{10.54}$$

In other words, W_0 is a holomorphic Hamiltonian function for the infinitesimal symmetry ζ.

The only $N = 4$ model in 4 dimensions is pure Yang-Mills theory, which is reduced from 10 dimensions. In the description on $M^{4|4}$ there is a constrained connection $\mathcal{A}$ and three chiral superfields $\Sigma_1, \Sigma_2, \Sigma_3$ with values in the complex adjoint bundle. The moment map is

$$\mu(\sigma_1, \sigma_2, \sigma_3) = \frac{\sqrt{-1}}{2}\Big\{[\sigma_1, \bar{\sigma}_1] + [\sigma_2, \bar{\sigma}_2] + [\sigma_3, \bar{\sigma}_3]\Big\}, \tag{10.55}$$

and there is a superpotential

$$W(\sigma_1, \sigma_2, \sigma_3) = \sqrt{2}\,\langle \sigma_1, [\sigma_2, \sigma_3]\rangle. \tag{10.56}$$

Note that the $U(1)$ R-symmetry (2.81) is broken by the superpotential (cf. the end of §5.3). This model does have an $SU(4)$ R-symmetry; the $SU(3)$ subgroup which permutes the chiral superfields is manifest in this formulation.[42]

[42] In components the six *real* scalar fields are permuted by $SU(4) \cong \text{Spin}(6)$.

CHAPTER 11
The Vector Multiplet on $M^{6|(8,0)}$

We prove Proposition 10.18, following the treatment in Koller (see References).

§11.1. Complements on $M^{6|(8,0)}$

As in §2.1 we write the relevant spinor representation S, complexified, as $S_{\mathbb{C}} = S_0 \otimes W$. The real structure comes from pseudoreal structures on S_0 and W.

The following lemma gives a convenient description, compatible with the action of $GL(S_0) \times GL(W)$, of the symmetric and exterior powers of $S_{\mathbb{C}}$. For ρ a representation of the symmetric group S_n, we note $\mathrm{Schur}_\rho(U)$ the corresponding Schur functor

$$\mathrm{Schur}_\rho(U) := \mathrm{Hom}_{S_n}(\rho, U^{\otimes n}). \tag{11.1}$$

One has

$$\bigoplus \mathrm{Schur}_\rho(V) \otimes \rho \xrightarrow{\sim} V^{\otimes n}, \tag{11.2}$$

the sum running over representatives of the isomorphism classes of irreducible representations of S_n. The map (11.2) is an isomorphism of representations of $GL(V) \times S_n$.

Lemma 11.3. *Let U and V be vector spaces. One has*

$$\bigoplus \mathrm{Schur}_\rho(U) \otimes \mathrm{Schur}_{\rho^\vee}(V) \xrightarrow{\sim} \mathrm{Sym}^n(U \otimes V) \tag{11.4}$$

$$\bigoplus \mathrm{Schur}_\rho(U) \otimes \mathrm{Schur}_{\rho^\vee \otimes \mathrm{sgn}}(V) \xrightarrow{\sim} \bigwedge^n (U \otimes V) \tag{11.5}$$

Note that ρ and $\rho^\vee$ are isomorphic. In (11.5) sgn is the sign character of S_n.

Proof. Apply (11.2) to U and V, and take the coinvariants (resp. co-anti-invariants) by S_n.

On $M^{6|(8,0)}$ the tangent bundle is a direct sum $V \oplus \tau$. As vector bundles the structural odd distribution τ is the constant bundle S^*, parity changed, and V is the constant bundle real part of $\bigwedge^2 S_0$. In particular, each s in $S_{\mathbb{C}}^*$ defines a complex odd vector field D_s in $\tau_{\mathbb{C}}$. The bases $\{f_a\}$ of S_0^* and $\{e_i\}$ of W^* define a moving

frame (see §2.5) consisting of the D_{ai} and ∂_{ab} ($a < b$). For $\partial_{ba} := -\partial_{ab}$ one has $D_{ai} = D_{f_a \otimes e_i}$ and

$$[D_{ai}, D_{bj}] = -2\epsilon_{ij}\partial_{ab}. \tag{11.6}$$

For x in $\mathrm{Sym}^2(S^*_{\mathbb{C}})$ define $D(x)$ by

$$D(s't') = [D_{s'}, D_{t'}]. \tag{11.7}$$

By the lemma,

$$\mathrm{Sym}^2(S^*_{\mathbb{C}}) = \bigwedge^2 S^*_0 \otimes \bigwedge^2 W^* \ \oplus \ \mathrm{Sym}^2(S^*_0) \otimes \mathrm{Sym}^2(W^*), \tag{11.8}$$

and (11.6) gives that $D(x)$ vanishes for x in the second factor. By polarization in w this amounts to saying that for w in W^*, we have $[D_{s_1 \otimes w}, D_{s_2 \otimes w}] = 0$. For $w \neq 0$ fixed in W^*, the $D_{s \otimes w}$ ($s \in S^*_0$) span an integrable distribution τ_w; its leaves are affine spaces with vector space of translations $S_0 \otimes w$, parity changed. Of course, τ_w is a complex distribution so that it makes sense to speak of its leaves only after complexification, at least in the odd direction (cs spaces). One can as well say that for $s \neq 0$ fixed in S^*_0, the $D_{s \otimes w}$ ($w \in W^*$) span an integrable distribution τ_s, with leaves affine spaces with vector space of translations $s \otimes W^*$.

The intersection of $s \otimes W^*$ and of $S^*_0 \otimes w$ is the line spanned by $s \otimes w$. This line also gives rise to an integrable distribution, with leaves affine spaces of dimension $0|1$, intersections of a leaf of τ_s and a leaf of τ_w.

We can now use the volume forms ν and ϵ to define differential operators

$$D^4_w := \nu^{1234} D_{f_1 \otimes w} D_{f_2 \otimes w} D_{f_3 \otimes w} D_{f_4 \otimes w} \tag{11.9}$$

$$D^2_s := \epsilon^{12} D_{s \otimes e_1} D_{s \otimes e_2}. \tag{11.10}$$

They can be viewed as integrals on the leaves of τ_w (resp. τ_s).

Lemma 11.11. $D^4_w D^2_s = D^2_s D^4_w = 0$.

Proof. It suffices to check it for w and s not zero; if we take a basis with $f_1 = s$ and $e_1 = w$, the lemma reduces to $D^2_{s \otimes w} = 0$.

Variant. D^4_w produces a function constant along the leaves of τ_w. The operator D^2_s is right divisible by $D_{s \otimes w}$. As $s \otimes w$ is in τ_w, the operator D^2_s annuls the image of D^4_w. Permuting the roles of S^*_0 and W^*, one similarly proves that $D^4_w D^2_s = 0$.

§11.2. Constrained connections

Let G be a Lie group, with Lie algebra $\mathfrak{g}$. We think to connections ∇ on a G-torsor $\mathcal{P}$ as follows: For each representation R of G, the principal bundle $\mathcal{P}$ gives rise to a vector bundle $R^{\mathcal{P}}$, and ∇ gives connections on the $R^{\mathcal{P}}$, functorially in R and compatibly with the tensor product.

From this point of view, the curvature F is the $\mathfrak{g}^{\mathcal{P}}$-valued 2-form such that for each R, the function $-i_X i_Y F$ acts on $R^{\mathcal{P}}$ (by the action of $\mathfrak{g}^{\mathcal{P}}$ on $R^{\mathcal{P}}$ deduced from the action of $\mathfrak{g}$ on R) by $[\nabla_X, \nabla_Y] - \nabla_{[X,Y]}$. The formula

$$-\iota(X)\iota(Y)F = [\nabla_X, \nabla_Y] - \nabla_{[X,Y]} \tag{11.12}$$

differs by a sign from the one which holds when ∇_X is viewed as a horizontal vector field on $\mathcal{P}$ lifting X. (See (6.75) for an example of the latter formula.)

A constrained connection is one whose curvature vanishes in the τ-direction. Let us write $\mathcal{D}_s$ for $\nabla_{\mathcal{D}_s}$ and $\mathcal{D}_{ai}$ for $\mathcal{D}_{f_a \otimes e_i}$. The constraint is that, in parallel to (11.6),

$$[\mathcal{D}_{ai}, \mathcal{D}_{bj}] = -2\epsilon_{ij}\nabla_{ab}. \tag{11.13}$$

A constrained connection is uniquely determined by its restriction to τ. For this restriction, here is what the constraint becomes. For x in $\mathrm{Sym}^2(S^*_{\mathbb{C}})$, define $\mathcal{D}(x)$ by $\mathcal{D}(s't') = [\mathcal{D}_{s'}, \mathcal{D}_{t'}]$; the constraint is that $\mathcal{D}(x)$ vanishes for x in the second factor (11.8) of $\mathrm{Sym}^2(S_{\mathbb{C}})$. This amounts to saying that for w in W^*, the connection is integrable along the leaves of τ_w. It also amounts to saying that it is integrable along the leaves of τ_s ($s \in S^*_0$). If we define $\mathcal{D}^4_w$ and $\mathcal{D}^4_s$ by (11.9) and (11.10), with D replaced by $\mathcal{D}$, we still have (in any $V^{\mathcal{P}}$)

$$\mathcal{D}^4_w \mathcal{D}^2_s = \mathcal{D}^2_s \mathcal{D}^4_w = 0, \tag{11.14}$$

with the same proof.

§11.3. An auxiliary Lie algebra

Let $\mathcal{L} = \bigoplus_{\nu=0}^{\infty} \mathcal{L}^\nu$ be the graded super Lie algebra (with parity = degree modulo 2) generated in degree 1 by $S^*_0 \otimes W^*$, and with relations given by

$$\mathrm{Sym}^2(S^*_0) \otimes \mathrm{Sym}^2(W^*) \subset \mathrm{Sym}^2(S^*_0 \otimes W^*). \tag{11.15}$$

For any constrained connection, the $\mathcal{D}(s)$ obey the relations of $\mathcal{L}$: for any representation R of G, the Lie algebra $\mathcal{L}$ acts on the local sections of $R^{\mathcal{P}}$. In degree one, we get the $\mathcal{D}(s)$; in degree two, the ∇_{ab}: the connection in the direction of V; in degree three, the components $F_{ab,ck} = [\nabla_{ab}, \mathcal{D}_{ck}]$ of the curvature, in the moving frame of §2.5; in degree ≥ 3 the action of elements of $\mathcal{L}$ is given by sections of $\mathfrak{g}^{\mathcal{P}}$, iterated covariant derivatives of the components of the curvature.

As a preliminary to a description in component fields of constrained connections, we now analyze $\mathcal{L}$ in degree ≤ 5. In degree 1, we have a basis $\mathcal{D}_{ai}$. In degree 2 we have $\mathcal{L}^2 \cong \bigwedge^2 S^*_0$, with basis the ∇_{ab} ($a < b$) defined by (11.13).

Lemma 11.16. *The iterated bracket* $[[\mathcal{D}_{ai}, \mathcal{D}_{bj}], \mathcal{D}_{ck}] = -2\epsilon_{ij}[\nabla_{ab}, \mathcal{D}_{ck}]$ *is antisymmetric in* (a, b, c).

Proof. By (11.13), the iterated bracket B is antisymmetric in (a, b) and in (i, j). As i, j, k take only two values, two of them are equal. We consider the cases $i = j$, $j = k$ and $i = k$ in turn:

$i = j$: one has $B = 0$;
$i = k$: $B = [\mathcal{D}_{ai}, [\mathcal{D}_{bj}, \mathcal{D}_{ck}]]$, which is antisymmetric in (b, c);
$j = k$: B is similarly antisymmetric in (a, c).

In all cases, antisymmetry in (a, b, c) follows.

By the lemma, $[\nabla_{ab}, \mathcal{D}_{ck}]$ defines a morphism $\bigwedge^3 S_0^* \otimes W^* \to \mathcal{L}^3$. As S_0^* is of dimension 4, the volume form of S_0 identifies $\bigwedge^3 S_0^*$ to the dual S_0 of S_0^*. Hence we have a map

$$\Lambda\colon S_0 \otimes W \longrightarrow \mathcal{L}^3, \tag{11.17}$$

where we take

$$[\mathcal{D}_{ak}, \nabla_{bc}] = 2\epsilon_{kj}\nu_{abcd}\Lambda^{dj}. \tag{11.18}$$

We find it convenient to lower the index on Λ by

$$\Lambda_k^d = \epsilon_{kj}\Lambda^{dj}, \tag{11.19}$$

and so

$$[\mathcal{D}_{ak}, \nabla_{bc}] = 2\nu_{abcd}\Lambda_k^d. \tag{11.20}$$

For any x in $S_0 \otimes W^*$, one easily checks that $\epsilon_{ij}\nu_{abcd}x_k^d$, averaged by cyclic permutations of (ac, bj, ck), vanishes. This gives the Jacobi identity required to check that (11.17) is an isomorphism.

Degree 4

The bracket $[\quad]\colon \mathcal{L}^2 \otimes \mathcal{L}^2 \to \mathcal{L}^4$ defines a map

$$\begin{aligned} \textstyle\bigwedge^2\bigwedge^2 S_0^* &\longrightarrow \mathcal{L}^4 \\ (f_a \wedge f_b) \wedge (f_c \wedge f_d) &\longmapsto [\nabla_{ab}, \nabla_{cd}]. \end{aligned} \tag{11.21}$$

The iterated exterior power $\bigwedge^2\bigwedge^2 S_0^*$ is the orthogonal Lie algebra of $\bigwedge^2 S_0^*$, provided with the symmetric bilinear form $h_{ab,cd} = \nu_{abcd}$. The exceptional isomorphism $SL(S_0^*) \xrightarrow{\sim} \mathrm{Spin}(\bigwedge^2 S_0^*)$ identifies it with the Lie algebra of traceless endomorphisms of S_0^*. The map is $f_a^b \mapsto F_{ab,cd}$, with

$$F_{ab,cd} = -\nu_{abef}(\delta_c^e f_d^f - \delta_d^e f_c^f), \tag{11.22}$$

or, moving up the vector index ab by $h^{ab,cd} = \nu^{abcd}$ and multiplying by a suitable factor, we have

$$F_{cd}^{ab} = \text{ antisymmetrization in } (a,b) \text{ and in } (c,d) \text{ of } \delta_c^a f_d^b. \tag{11.23}$$

Note that F, given by this formula, is antisymmetric in (ab, cd) if and only if f is traceless. In particular, we obtain a traceless f_b^a in $\mathcal{L}^4$ with

$$[\nabla_{ab}, \nabla_{cd}] = -\nu_{abef}\left(\delta_c^e f_d^f - \delta_d^e f_c^f\right). \tag{11.24}$$

The degree 4 part of $\mathcal{L}$ is generated by the $[\mathcal{D}_{ai}, \Lambda_j^b]$. They are related to the $F_{ab,cd} := [\nabla_{ab}, \nabla_{cd}]$ by

$$\begin{aligned} -2\epsilon_{k\ell}F_{ab,cd} &= [\nabla_{ab}, [\mathcal{D}_{ck}, \mathcal{D}_{d\ell}]] \\ &= [[\nabla_{ab}, \mathcal{D}_{ck}], \mathcal{D}_{d\ell}] && \text{symmetrized in } (ck, d\ell) \\ &= 2\nu_{abce}[\mathcal{D}_{d\ell}, \Lambda_k^e] && \text{symmetrized in } (ck, d\ell) \\ &= 2\nu_{abef}\delta_c^e[\mathcal{D}_{d\ell}, \Lambda_k^f] && \text{symmetrized in } (ck, d\ell). \end{aligned} \tag{11.25}$$

Let us extract from this formula the parts symmetric and antisymmetric in (k,ℓ). Antisymmetrization gives

$$-4\epsilon_{k\ell}F_{ab,cd} = \nu_{abef}\delta^e_c[\mathcal{D}_{d\ell},\Lambda^f_k] \quad \text{antisymmetrized in } (c,d),\ (k,\ell), \tag{11.26}$$

in other words,

$$4\epsilon_{k\ell}f^a_b = \mathcal{D}_{b\ell}\Lambda^a_k \qquad \text{antisymmetrized in } (k,\ell). \tag{11.27}$$

Symmetrization in (k,ℓ) gives the vanishing of the symmetrization in (c,d), (k,ℓ) and antisymmetrization in (a,b) of $\delta^a_c\mathcal{D}_{d\ell}\Lambda^b_k$. Let us fix k, ℓ and let D^b_d be the symmetrization of $\mathcal{D}_{d\ell}\Lambda^b_k$ in (k,ℓ). For $a=c=1$, $b=2$, $d=1$ we get $D^2_1=0$. This holding in any basis, D^b_d is a multiple of δ^b_a. Hence for some $\mathcal{E}_{ij}$, we have[43]

$$\delta^a_b\mathcal{E}_{ij} = \mathcal{D}_{bj}\Lambda^a_i \qquad \text{symmetrized in } (i,j). \tag{11.29}$$

The morphism from $\bigwedge^2\bigwedge^2 S^*_0 \oplus \mathrm{Sym}^2(W^*)$ to $\mathcal{L}^4$ is an isomorphism. Indeed, if we define $\mathcal{L}^4$ as being $\bigwedge^2\bigwedge^2 S^*_0 \oplus \mathrm{Sym}^2(W^*)$ and the bracket by (11.27), (11.29), we get a Lie algebra. The only new Jacobi identity to check is (11.25), from which (11.27) and (11.29) were derived.

Degree ≥ 5

From the forthcoming description (Proposition 11.33) of constrained connections in components, one can deduce that the part of degree ≥ 2 of the super Lie algebra $\mathcal{L}$ is free, with generators the ∇_{ab}, Λ^a_i and $\mathcal{E}_{ij}$. To prepare for Proposition 11.33, we show here that it is generated by those elements. If $\mathcal{L}_1$ is the subalgebra generated by the ∇_{ab}, Λ^a_i and $\mathcal{E}_{ij}$, it suffices to check that a $\mathcal{D}_{ai}$, applied to any of the generators, is again in $\mathcal{L}_1$. It only remains to show that the $[\mathcal{D}_{ai},\mathcal{E}_{jk}]$ are linear combinations of the $[\nabla_{cd},\Lambda^e_\ell]$. The formula

$$\mathcal{D}_{ai}\mathcal{E}_{jk} = C\epsilon_{ij}\nabla_{ab}\Lambda^b_k \qquad \text{symmetrized in } (j,k) \tag{11.30}$$

for some constant C is the only possible formula compatible with the grading, the actions of $SL(S^*_0)$, $SL(W)$, and with the symmetry in (j,k). (Further computation shows $C=-2$.) Here, we will content ourselves to check that, modulo the $[\nabla,\Lambda]$, all

$$[\mathcal{D}_{ai},[\mathcal{D}_{bj},[\mathcal{D}_{ck},[\mathcal{D}_{d\ell},\mathcal{D}_{em}]]]] = 4\epsilon_{\ell m}\nu_{cdef}[\mathcal{D}_{ai},[\mathcal{D}_{bj},\Lambda^f_k]] \tag{11.31}$$

vanish.

If we antisymmetrize (11.31) in (j,k), we get by (11.27) a $[\mathcal{D},[\nabla,\nabla]]$, which Jacobi turns into a $[\nabla,[\nabla,\mathcal{D}]]$, i.e., a $[\nabla,\Lambda]$. Modulo the $[\nabla,\Lambda]$, this gives the symmetry of (11.31) in (j,k) and, by (11.29), its antisymmetry in (b,c,d,e).

[43] $\mathcal{E}_{ij}$ is related to the E^i_j of Chapter 10 by

$$\mathcal{E}_{ij} = 2\sqrt{-1}\,\epsilon_{ik}E^k_j. \tag{11.28}$$

If a quantity x_{ijk} is symmetric in (j,k), to prove its vanishing, it suffices to check the vanishing of its symmetrization y_{ijk} in (i,j). Indeed, the symmetrization of y in (j,k) gives x_{ijk} back, up to a multiple of the complete symmetrization (of x or y, this amounts to the same):

$$y_{ijk} + y_{jki} = x_{ijk} + (x_{ikj} + x_{jik} + x_{kij}). \tag{11.32}$$

To prove the vanishing of (11.31) modulo the $[\nabla, \Lambda]$, it hence suffices to consider its symmetrization $(11.31)^S$ in (i,j). In $(11.31)^S$, if we symmetrize in (a,b), we get 0: a $[[\mathcal{D}_{ai}, \mathcal{D}_{bj}] \dots]$ symmetrized in (i,j). We conclude that $(11.31)^S$ is antisymmetric in (a,b). Modulo the $[\nabla, \Lambda]$, it is also antisymmetric in (b,c,d,e), and as $\dim S_0^* = 4 < 5$, its vanishing follows.

§11.4. Components of constrained connections

In classical differential geometry, if N is a subvariety of M and $(\mathcal{P}, \nabla)$ a torsor with connection on M, the restriction of $(\mathcal{P}, \nabla)$ to a formal neighborhood of N is determined up to unique isomorphism by (a) the restriction (P, ∇) of $(\mathcal{P}, \nabla)$ to N, and (b) the value on N of all iterated covariant derivatives of the components of the curvature of P. The same holds in the super case. In addition, if N is of codimension $0|s$ in M, the formal neighborhood of N in M is just M itself.

If the codimension is $0|1$, the value on N of the curvature is enough. Indeed, take local coordinates (θ^0, x^i), where the x^i can be even or odd, for which N is defined by $\theta^0 = 0$. A trivialization of P on N extends uniquely to a trivialization s of $\mathcal{P}$ for which $\nabla_0 s = 0$ on N. Given a trivialization of P, the connection on P is given by a $\mathfrak{g}$-valued 1-form α, while $(\mathcal{P}, \nabla)$ is given by a $\mathfrak{g}$-valued 1-form β on M, such that $\langle \partial_0, \beta \rangle = 0$ on N. Let us write $\beta = d\theta^0\, \theta^0 b + \sum dx^i (c^i + \theta^0 d^i)$, with b, c, d functions of the x^i. Then, α gives the c^i, the components $F_{0,i}$ of the curvature, restricted to N, gives the d^i and F_{00} restricted to N gives b.

To handle the case of codimension $0|s$, one chooses $N \subset N_1 \subset \dots \subset M$, each of codimension $0|1$ in the next, and one uses induction on s: on N_i, after recovering $(\mathcal{P}, \nabla)|N_i$, one recovers from the data on N the data on N_i of all covariant derivatives of the components of the curvature, restricted to N_i.

Let $(\mathcal{P}, \nabla)$ be a constrained G-torsor on $M^{6|(8,0)}$. From §11.3, we obtain from it a $\mathfrak{g}^{\mathcal{P}}$-valued spinor Λ_i^a, and $\mathfrak{g}^{\mathcal{P}}$-valued auxiliary fields $\mathcal{E}_{ij}$ By restriction to Minkowski space $\check{M}^6$, we obtain a torsor with connection (P, ∇) on M^6, $\mathfrak{g}^P$-valued spinors λ_i^a and $\mathfrak{g}^P$-valued auxiliary fields E_{ij}. By §11.3, all covariant derivatives of the components of $\mathcal{P}$, restricted to M^6, can be expressed from ∇, λ and $\mathcal{E}$. It follows that $(\mathcal{P}, \nabla)$ is determined up to unique isomorphism by (P, ∇, λ, E). More precisely, one has the following rephrasing of Proposition 10.18.

Proposition 11.33. *The functor $(\mathcal{P}, \nabla) \mapsto (P, \nabla, \lambda, E)$ is an equivalence of categories.*

For a spelling out of what this means, we refer to the text following Proposition 7.5. We will check Proposition 11.33 in the complex setting; the real case results from it as in the proof of Proposition 8.7.

Proof. We have to check that any (P, ∇, λ, E) comes from some constrained $(\mathcal{P}, \nabla)$. The flat P with $\lambda, E = 0$ corresponds to $\mathcal{P}$ flat. We now construct vector fields on the space of connections, tangent to the constrained ones, to get from there to any

(P, ∇, λ, E). Rather, we work with the space of connections along τ. A deformation of a connection $\mathcal{D}$ in the τ direction is given by $\mathfrak{g}^{\mathcal{P}}$-valued $\delta\mathcal{D}_{ai}$. It is tangent to constrained connections if the symmetrization in (a,b) and (i,j) of $[\delta\mathcal{D}_{ai}, \mathcal{D}_{bj}]$ vanishes.

We will need the following operators acting on $\mathfrak{g}^{\mathcal{P}}$:

$$\begin{aligned} &\mathcal{D}_{ab} = \tfrac{1}{2!}\epsilon^{ij}\mathcal{D}_{ai}\mathcal{D}_{bj}, && \text{symmetrized in } (a,b) \\ &\mathcal{D}_{ijk\ell} = \tfrac{1}{4!}\epsilon^{abcd}\mathcal{D}_{ai}\mathcal{D}_{bj}\mathcal{D}_{ck}\mathcal{D}_{d\ell}, && \text{symmetrized in } (i,j,k,\ell). \end{aligned} \tag{11.34}$$

They polarize the $\mathcal{D}_s^2$ and $\mathcal{D}_w^4$ of (11.14), and from (11.14) we get that

$$\mathcal{D}_{ab}\mathcal{D}_{ijk\ell} = \mathcal{D}_{ijk\ell}\mathcal{D}_{ab} = 0 \tag{11.35}$$

(for their action on sections of $\mathfrak{g}^{\mathcal{P}}$).

Fix $\mathfrak{g}^{P}$-valued fields V^{ij} and consider

$$\delta\mathcal{D}_{ai} := \epsilon^{jj'}\mathcal{D}_{aj}\mathcal{D}_{ij'k\ell}V^{k\ell}. \tag{11.36}$$

If we start from a constrained $\mathcal{D}$, we have

$$\begin{aligned} &[\mathcal{D}_{ai}, \delta\mathcal{D}_{bj}] && \text{symmetrized in } (a,b),\ (i,j) \\ &\quad = \epsilon^{kk'}\mathcal{D}_{ai}\mathcal{D}_{bk}\mathcal{D}_{jk'\ell m}V^{\ell m} && \text{symmetrized in } (a,b) \end{aligned} \tag{11.37}$$

and the (a,b)-symmetrization of $\mathcal{D}_{ai}\mathcal{D}_{bk}$ is alternating in (i,k): it is $2\epsilon_{ik}\mathcal{D}_{ab}$. We get

$$\ldots = -2\mathcal{D}_{ab}\mathcal{D}_{ji\ell m}V^{\ell m} = 0.$$

The $\delta\mathcal{D}_{ai}$ is the promised vector field, tangent to the constrained $\mathcal{D}$.

Let (θ) be the ideal defining M^6 in $M^{6|(8,0)}$. To conclude, it remains to check that:

(a) for V in $(\theta)^6$, the variation of ∇_{ab} on $\breve{M}$ depends only on V modulo $(\theta)^7$, and the ∇_{ab} are changed freely;
(b) for V in $(\theta)^7$, the variation of λ_i^a depends only on V modulo $(\theta)^8$, and the λ_i^a are changed freely;
(c) for V in $(\theta)^8$, the auxiliary fields E are changed freely.

The variation of $[\mathcal{D}_{ai}, \mathcal{D}_{bj}]$ (resp. $[\mathcal{D}_{ai}, [\mathcal{D}_{bj}, \mathcal{D}_{ck}]]$, resp. $[\mathcal{D}_{ai}, [\mathcal{D}_{bj}, [\mathcal{D}_{ck}, \mathcal{D}_{d\ell}]]]$) is given by 6 (resp. 7, resp. 8) operators $\mathcal{D}$ applied to V. For V in $(\theta)^6$ (resp. $(\theta)^7$, resp. $(\theta)^8$), this variation, restricted to M^6, depends only on V modulo the next power of (θ) and, to compute it, one may replace each $\mathcal{D}_{ai}$ by ∂_{ai}. The resulting simplified computation is left to the reader.

APPENDIX
Sign Manifesto

§1. Standard mathematical conventions

- *We apply the sign rule relentlessly.*

This means that when passing from ordinary algebra to $\mathbb{Z}/2$-graded, or super, algebra we pick up a sign $(-1)^{|a|\,|b|}$ when permuting homogeneous elements a, b of parity $|a|, |b|$. Structure maps (multiplications, Lie brackets, inner products, ...) are even.

For example, consider a graded complex vector space $V = V^0 \oplus V^1$. A hermitian inner product $\langle \cdot, \cdot \rangle$ satisfies, among other properties,

$$\langle v_1, v_2 \rangle = (-1)^{|v_1|\,|v_2|}\overline{\langle v_2, v_1 \rangle}, \qquad v_1, v_2 \in V \text{ homogeneous.} \tag{A.1}$$

and from the evenness of the inner product it follows that V^0 is orthogonal to V^1. From (A.1) we deduce that $\langle v, v \rangle$ is real for v even and pure imaginary for v odd. The adjoint T^* of a homogeneous linear operator $T\colon V \to V$ is characterized by

$$\langle Tv_1, v_2 \rangle = (-1)^{|T|\,|v_1|}\langle v_1, T^* v_2 \rangle. \tag{A.2}$$

Skew-adjoint operators form a super Lie algebra.

- *Symmetry groups act on the left.*

For example, if $\mathfrak{g}$ is a Lie algebra, then an action of $\mathfrak{g}$ on a vector space V is a *homomorphism* $\mathfrak{g} \to \mathrm{End}(V)$. Brackets are preserved. On the other hand, an action of $\mathfrak{g}$ on a manifold M is an *antihomomorphism* $\mathfrak{g} \to \mathrm{VectorFields}(M)$. The reversal of sign comes from the rule $\xi f = \frac{d}{dt}\exp(t\xi)^* f$ and the fact that $f \mapsto u^* f$ is a *right* action of diffeomorphisms (u) on functions (f).

§2. Choices

- *A hermitian inner product on a complex vector space V is conjugate linear in the first variable:*

$$\langle \lambda_1 v_1, \lambda_2 v_2 \rangle = \overline{\lambda_1}\lambda_2 \langle v_1, v_2 \rangle, \qquad \lambda_i \in \mathbb{C}, \quad v_i \in V. \tag{A.3}$$

- *If $V = V^0 \oplus V^1$ is a super Hilbert space, then*

$$-i\langle v, v\rangle \geq 0, \qquad v \in V^1. \tag{A.4}$$

- *We pass from self-adjoint operators to skew-adjoint operators using multiplication by $-i$:*

$$T \text{ self-adjoint} \longleftrightarrow -iT \text{ skew-adjoint.} \tag{A.5}$$

- *The Lorentz metric g on n dimensional Minkowski space has signature $(1, n-1)$:*

$$\text{Signature}(g) = + - - - \cdots \tag{A.6}$$

- *The quantum hamiltonian $\hat{H}$ is minus the operator which corresponds to infinitesimal time translation:*

$$\hat{H} = -\hat{P}_0. \tag{A.7}$$

§3. Rationale

The first choice (A.3) is not the usual one in mathematics, but it has its merits. For example, since linear operators act on the left, it makes sense to have the commuting scalar multiplication act on the right. In mathematics we do follow this convention for modules over noncommutative rings. With right scalar multiplication (A.3) reads: $\langle v_1\lambda_1, v_2\lambda_2\rangle = \overline{\lambda_1}\langle v_1, v_2\rangle\lambda_2$. We do not adopt this convention for scalar multiplication, but do adopt (A.3). Physicists like (A.3) in view of Dirac's notation '$\langle v_1|T|v_2\rangle$' for '$\langle v_1, Tv_2\rangle$'. Comment: In computations it is often more convenient and safer to work with a bilinear form rather than a sesquilinear form, and so to write the sesquilinear inner product as '$\langle\overline{v_1}, v_2\rangle$'.

From a mathematical point of view it is more natural to quantize with *skew*-adjoint operators, since they form a Lie algebra. We use (A.5) to convert to *self*-adjoint operators, whose real eigenvalues correspond to physical measurements.

The sign choice in (A.6) leads to the usual bosonic lagrangian (A.20) with a plus sign in front of the kinetic energy.

In (A.7) we assume that time translation is a symmetry of a quantum theory, so that the infinitesimal generator is represented by a self-adjoint operator $\hat{P}_0$ on the quantum (super)Hilbert space. The minus sign gives the standard answer for the hamiltonian of a classical free particle.

§4. Notation

Throughout $i = \sqrt{-1}$.

Let M denote n dimensional affine Minkowski space with associated vector space of translations V and future timelike cone $C \subset V$. We fix linear coordinates $x^0, \ldots, x^{n-1}$ with respect to which the metric is

$$g = g_{\mu\nu}\, dx^\mu \otimes dx^n = (dx^0)^2 - \cdots - (dx^{n-1})^2, \tag{A.8}$$

and the cone is

$$C = \{x : \langle x, x\rangle \geq 0 \text{ and } x^0 \geq 0\}. \tag{A.9}$$

Let $\{e_\mu\}$ be the corresponding basis of V and ∂_μ the corresponding vector field on M. The standard density on M is

$$|d^n x|\, d^s\theta = |dx^0 \dots dx^{n-1}|. \tag{A.10}$$

Let S be a real spin representation. Fix a basis $\{f^a\}$ of S and dual basis $\{f_a\}$ of S^*. Then there are symmetric pairings

$$\begin{aligned} \Gamma\colon S^* \otimes S^* &\longrightarrow V \\ \tilde{\Gamma}\colon S \otimes S &\longrightarrow V. \end{aligned} \tag{A.11}$$

We write

$$\begin{aligned} \Gamma(f_a, f_b) &= \Gamma^\mu_{ab} e_\mu, \\ \tilde{\Gamma}(f^a, f^b) &= \tilde{\Gamma}^{\mu ab} e_\mu, \end{aligned} \tag{A.12}$$

where as usual we sum over repeated indices if one is upstairs and the other is downstairs. We raise and lower indices using the metric. The pairings (A.11) are assumed to satisfy the Clifford relation

$$\tilde{\Gamma}^{\mu ab}\Gamma^\nu_{bc} + \tilde{\Gamma}^{\nu ab}\Gamma^\mu_{bc} = 2g^{\mu\nu}\delta^a_c \tag{A.13}$$

and the positivity condition

$$\Gamma(s^*, s^*) \in C \qquad \text{for all } s^* \in S^*. \tag{A.14}$$

For $v \in C^o$, the form $\langle v, \Gamma(s^*, s^*)\rangle$ is then positive definite. From (A.13) and (A.14) it follows that

$$\tilde{\Gamma}(s, s) \in C \qquad \text{for all } s \in S. \tag{A.15}$$

In a classical field theory we work with a space of fields $\mathcal{F}$, where $f \in \mathcal{F}$ is some sort of function on M. An infinitesimal symmetry is a vector field ξ on $\mathcal{F}$ which preserves the lagrangian in a certain sense. Corresponding to ξ is a Noether current J_ξ, which is a twisted $(n-1)$-form on M. The Noether charge Q_ξ is the integral of J_ξ over a time slice. We usually consider the current and charge only on the space of classical solutions $\mathcal{M}$, which carries a closed 2-form ω. The infinitesimal symmetry and Noether charge are related by

$$dQ_\xi = -\iota(\xi)\omega. \tag{A.16}$$

For $(\mathcal{M}, \omega)$ symplectic, (A.16) can be rewritten

$$\xi f = \{Q_\xi, f\} \tag{A.17}$$

for f a function on $\mathcal{M}$.

Quantization is, in principle, a map

$$Q \longmapsto \hat{Q} \tag{A.18}$$

from functions on $\mathcal{M}$ to operators on a complex Hilbert space $\mathcal{H}$. We assume

$$\overline{\hat{Q}} = \hat{Q}^*, \tag{A.19}$$

so that real functions map to self-adjoint operators.

Let P_μ be infinitesimal translation in the Poincaré algebra and Q_a the odd generator of the supersymmetry algebra.[44] Let $\hat{P}_\mu, \hat{Q}_a$ be the corresponding quantum operators.

[44]The notational conflict between the supersymmetry generator and the Noether charge is too ingrained to correct.

§5. Consequences of §2 on other signs

- *The kinetic lagrangian for a scalar field $\phi\colon M \to \mathbb{R}$ is*

$$L = \frac{1}{2}|d\phi|^2 \, |d^n x| \, d^s\theta = \frac{1}{2} g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi \, |d^n x| \, d^s\theta. \tag{A.20}$$

The sign of this term is a consequence of (A.6); it is the main rationale for preferring (A.6) over the other choice.

- *Suppose $V = V^0 \oplus V^1$ is a graded hermitian vector space and T an odd skew-adjoint operator. Then*

$$i[T, T] \geq 0. \tag{A.21}$$

Observe from (A.2), (A.3) and (A.4) that an odd skew-adjoint operator has eigenvalues on the line $i^{-1/2}\mathbb{R} \subset \mathbb{C}$.

- *The bracket in the supersymmetry algebra is*

$$[Q_a, Q_b] = -2\Gamma^\mu_{ab} P_\mu. \tag{A.22}$$

Because we use left group actions, upon quantization we expect a homomorphism from the supersymmetry algebra to *skew*-adjoint operators. Using (A.5) we see that the sign in (A.22) leads to

$$[-i\hat{Q}_a, -i\hat{Q}_b] = -2\Gamma^\mu_{ab}(-i\hat{P}_\mu). \tag{A.23}$$

Setting $a = b$ we see from (A.21) that $-2\Gamma^\mu_{aa}\hat{P}_\mu > 0$ for all a. From (A.14) we see that $\Gamma^\mu_{aa}P_\mu$ has nonnegative norm in V. Except possibly in dimension 2, the positive cone generated by $\{\Gamma^\mu_{aa}P_\mu\}_a$ includes P_0, and so the sign choice in (A.22) renders the hamiltonian nonnegative (rather than nonpositive), in view of (A.7).

- *The vector field $\hat{\xi}_{P_\mu}$ on $\mathcal{F}$ corresponding to infinitesimal translation P_μ is*

$$\hat{\xi}_{P_\mu} f = -\partial_\mu f, \qquad f \in \mathcal{F}. \tag{A.24}$$

This follows since a diffeomorphism $\varphi\colon M \to M$ acts on functions by $(\varphi^{-1})^*$.

- *If $\mathfrak{g}$ is a Lie algebra of infinitesimal symmetries, then the vector fields ξ_λ ($\lambda \in \mathfrak{g}$) on the space of fields $\mathcal{F}$ satisfy*

$$[\xi_{\lambda_1}, \xi_{\lambda_2}] = -\xi_{[\lambda_1,\lambda_2]}. \tag{A.25}$$

- *The Noether currents satisfy*[45]

$$\{j_{\lambda_1}, j_{\lambda_2}\} = -j_{[\lambda_1,\lambda_2]}. \tag{A.26}$$

[45] The bracketing operation on Noether currents is defined in [I-**Classical Fields**, §2.6].

- *The Noether charges satisfy*

$$\{Q_{\lambda_1}, Q_{\lambda_2}\} = -Q_{[\lambda_1,\lambda_2]}. \tag{A.27}$$

- *The quantum operators satisfy*

$$[-i\hat{Q}_{\lambda_1}, -i\hat{Q}_{\lambda_2}] = -i\hat{Q}_{[\lambda_1,\lambda_2]}. \tag{A.28}$$

Equations (A.25)–(A.27) follow from the fact that $\mathfrak{g} \to \text{VectorFields}(\mathcal{F})$ is an antihomorphism and the standard equations for Poisson brackets which follow from (A.16). Equation (A.28) says that $\mathfrak{g} \to \text{End}(\mathcal{H})$ is a homomorphism to skew-adjoint operators, where we use (A.5).

- *The self-adjoint quantum operators $\hat{Q}_1, \hat{Q}_2$ which correspond to classical functions Q_1, Q_2 on $\mathcal{M}$ satisfy*

$$[\hat{Q}_1, \hat{Q}_2] = -i\hbar\{Q_1, Q_2\}\hat{} \quad \textit{modulo } O(\hbar^2). \tag{A.29}$$

To the extent that (A.29) holds exactly, it says that the map

$$Q \longmapsto \frac{-i}{\hbar}\hat{Q} \tag{A.30}$$

to skew-adjoint operators is an antihomorphism. The sign in (A.30) is dictated by (A.5). The desire to have an *anti*homomorphism is dictated by (A.27) and (A.28), and this determines the sign in (A.29).

- *The Schrödinger equation for the evolution of a state ψ is*

$$\frac{\partial\psi}{\partial t} = \frac{-i}{\hbar}\hat{H}\psi. \tag{A.31}$$

Evolution through time t for a static hamiltonian $\hat{H}$ is $e^{-it\hat{H}/\hbar}$.

The sign follows from (A.7) and (A.5).

- *Let η_1^a, η_2^b be odd parameters and $\hat{\xi}_i$ the even vector field on $\mathcal{F}$ corresponding to $\eta_i^a Q_a$. Then*

$$[\hat{\xi}_1, \hat{\xi}_2]f = 2\eta_1^a\eta_2^b\Gamma_{ab}^\mu\partial_\mu f, \qquad f \in \mathcal{F}. \tag{A.32}$$

To see this, observe from (A.22) that in the abstract supersymmetry algebra we have

$$[\eta_1^a Q_a, \eta_2^b Q_b] = 2\eta_1^a\eta_2^b\Gamma_{ab}^\mu P_\mu. \tag{A.33}$$

Then (A.32) follows from (A.25) and (A.24); the minus signs in these two equations cancel.

- *If ψ_1, ψ_2 are complex classical odd quantities, then complex conjugation satisfies*

$$\overline{\psi_1\psi_2} = \overline{\psi_1}\,\overline{\psi_2}. \tag{A.34}$$

This is a consequence of the sign rule if we assume that $\psi \mapsto \overline{\psi}$ is a $*$-*operation* and ψ_1 (super)commutes with ψ_2. (A $*$ operation satisfies $(ab)^* = (-1)^{|a|\,|b|}\, b^* a^*$.) Notice that the classical statement (A.34) is consistent with the quantum statement (A.19), since the adjoint operation on linear operators is also a $*$ operation. Notice that the product of real commuting odd quantities is real.

- *The kinetic lagrangian for a spinor field $\psi\colon M \to S$ is*

$$L = \frac{1}{2}\psi \not{D}\psi \; |d^n x|\, d^s\theta = \frac{1}{2}\tilde{\Gamma}^{\mu a b}\psi_a \partial_\mu \psi_b \; |d^n x|\, d^s\theta. \tag{A.35}$$

- *The kinetic lagrangian for a dual spinor field $\lambda\colon M \to S^*$ is*

$$L = \frac{1}{2}\lambda \not{D}\lambda \; |d^n x|\, d^s\theta = \frac{1}{2}\Gamma^{\mu}_{ab}\lambda^a \partial_\mu \lambda^b \; |d^n x|\, d^s\theta. \tag{A.36}$$

The spinor fields are odd. In view of (A.34), the lagrangians (A.35) and (A.36) are real, as they must be in Minkowski space. It is easiest to check the sign in classical mechanics ($n = 1$). Then from (A.35) we deduce[46] the classical Poisson bracket

$$\{\psi, \psi\} = -1. \tag{A.37}$$

Upon quantization we know the corresponding operators satisfy (A.29). The sign in (A.37) is compatible with (A.21), and this means that the sign in (A.35) is correct.

- *The energy-momentum tensor is minus the Noether current of P_μ. The supercurrent is minus the Noether current of Q_a.*

This is a definition and follows by superPoincaré invariance from the definition (A.7) of the hamiltonian. It means that the charges computed from the energy-momentum tensor are energy and *minus* momentum.

§6. Differential forms

- *When computing with differential forms on superspace, we use a bigraded point of view.*[47]

Objects have a "cohomological" degree, corresponding to a classical (that is, non-super) degree, and a parity. The permutation of objects of parity p_1, p_2 and cohomological degree d_1, d_2 introduces two signs: a classical sign $(-1)^{d_1 d_2}$ and an additional factor $(-1)^{p_1 p_2}$.

- *On $\mathbb{R}^{p|q}$ with coordinates $t^1, \dots t^p, \theta^1, \dots \theta^q$ we have the following table of parities and cohomological degrees:*

[46] See Problem FP2 of [I-**Homework**] for a derivation.
[47] See the appendix to Chapter 1 of [I-**Supersymmetry**].

quantity	type	parity ($\mathbb{Z}/2\mathbb{Z}$)	coh deg ($\mathbb{Z}$)
t^μ	even coordinate	0	0
θ^a	odd coordinate	1	0
$\partial/\partial t^\mu$, $\iota(\partial/\partial t^\mu)$	even vector field	0	-1
$\partial/\partial\theta^a$, $\iota(\partial/\partial\theta^a)$	odd vector field	1	-1
$\mathrm{Lie}(\xi)$	Lie derivative	$p(\xi)$	0
$dt^\mu, \epsilon(dt^\mu)$	even 1-form	0	1
$d\theta^a, \epsilon(d\theta^a)$	odd 1-form	1	1
$d^p t\, d^q\theta$	berezinian	$q \pmod 2$	p
$\lvert d^p t\rvert\, d^q\theta$	density	$q \pmod 2$	0
$\iota(\xi_r)\dots\iota(\xi_1)\ d^p t\, d^q\theta$	integral form	$\sum p(\xi_i)+q \pmod 2$	$p-r$
$\iota(\xi_r)\dots\iota(\xi_1)\ \lvert d^p t\rvert\, d^q\theta$	integral density	$\sum p(\xi_i)+q \pmod 2$	$-r$

- *For X a (super)manifold the canonical pairing of vectors and 1-forms is written with the vector on the left.*

Therefore, by the sign rule, for a tangent vector ξ and a 1-form α there is a sign[48] when passing from the canonical pairing $\iota(\xi)\alpha$ to $(-1)^{p(\xi)p(\alpha)}\alpha(\xi)$, where p is the parity.

- *For integration over odd variables we have*

$$\int d\theta\ \theta = 1$$
$$\int d\theta^2 d\theta^1\ \theta^1\theta^2 = \int d\theta^2 \left(\int d\theta^1\ \theta^1\right)\theta^2 = 1.$$

- *For any vector field ξ we have the Cartan formula*

$$\mathrm{Lie}(\xi) = [d, \iota(\xi)]. \tag{A.38}$$

Both sides of (A.38) act on differential or integral forms.

§7. Miscellaneous signs

- *Let X be a smooth manifold, ξ a vector field on X, φ_t the one-parameter group of diffeomorphisms generated, and T a tensor field. Then*

$$\mathrm{Lie}(\xi)T = \frac{d}{dt}\Big|_{t=0} \varphi_t^* T = \frac{d}{dt}\Big|_{t=0} (\varphi_{-t})_* T. \tag{A.39}$$

- *On a Kähler manifold X the Riemannian metric g, Kähler form ω, complex structure J, and a local Kähler potential K are related by the equations*

$$\begin{aligned} \omega(\xi_1,\xi_2) &= g(J\xi_1,\xi_2) \\ \omega &= i\,\partial\bar\partial K. \end{aligned} \tag{A.40}$$

[48] See [I-**Supersymmetry**, §3.3].

- *Suppose f is a (suitable) function on a vector space V of dimension n. Then f and its Fourier transform $\hat{f}$ on V^* are related by*

$$\begin{aligned} \hat{f}(k) &= \frac{1}{(2\pi)^{n/2}} \int_V e^{-i\langle k,x\rangle} f(x)\, |d^n x|\, d^s\theta, \\ f(x) &= \frac{1}{(2\pi)^{n/2}} \int_{V^*} e^{+i\langle k,x\rangle} \hat{f}(k)\, |d^n k|. \end{aligned} \tag{A.41}$$

- *Suppose $\mathfrak{g}$ is the Lie algebra of a real Lie group G. Then the complexified Lie algebra $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g} \otimes \mathbb{C}$ carries a conjugation whose set of real points is $\mathfrak{g}$.*

For example, starting with the unitary group $G = U(n)$, we obtain the conjugation $A \mapsto -A^*$ on complex $n \times n$ matrices. (A^* is the conjugate transpose of A.)

REFERENCES

Most of the material discussed here was developed in the physics literature in the '70s and '80s. It is hopeless for us to reference all of the original papers. Our more modest goal is to give the reader an entry into the literature and to point out a few more specialized references we used.

Some standard physics books on the subject (alphabetical by author) are:

P. G. O. Freund, *Introduction to supersymmetry*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1986.

S. J. Gates, M. T. Grisaru, M. Roček, W. Siegel, *Superspace, One thousand and one lessons in supersymmetry*, Frontiers in Physics, vol. 58, Benjamin/Cummings Publishing Co. Inc., Reading, Mass., 1983.

M. F. Sohnius, *Introducing supersymmetry*, Phys. Rep. **128** (1985), 39–204.

J. Wess, J. Bagger, *Supersymmetry and supergravity*, Princeton Series in Physics, Princeton University Press, Princeton, NJ, 1992.

P. West, *Introduction to supersymmetry and supergravity*, World Scientific Publishing Co., Inc., Teaneck, NJ, 1990.

An expository introduction to some of this material may be found in

D. S. Freed, *Five Lectures on Supersymmetry*, American Mathematical Society, 1999 (to appear).

The superspace formulation of 6-dimensional Yang-Mills (Chapters 10 and 11) does not seem to be widely known, but it can be found in the literature. See, for example:

P. S. Howe, G. Sierra, P. K. Townsend, *Supersymmetry in six dimensions*, Nucl. Phys. B **221** (1983), 331–348.

J. Koller, *A six-dimensional superspace approach to extended superfields*, Nucl. Phys. B **222** (1983), 319–337.